

# A Polymer in a Multi-Interface Medium

Francesco Caravenna

Università degli Studi di Padova

Università di Roma 3 ~ July 21, 2009

# References

- ▶ [CP1] F. Caravenna and N. Pétrélis  
*A polymer in a multi-interface medium*  
AAP (2009), to appear
- ▶ [CP2] F. Caravenna and N. Pétrélis  
*Depinning of a polymer in a multi-interface medium*  
preprint (2009) [arXiv.org: 0901.2902]

# Outline

## 1. Introduction

What is a polymer?

Interaction with the environment

## 2. The model and the main results

Definition

The free energy

Path results

## 3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach

# Outline

## 1. Introduction

What is a polymer?

Interaction with the environment

## 2. The model and the main results

Definition

The free energy

Path results

## 3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach

# What is a polymer?

A **polymer** is a large molecule composed of repeating smaller units, called **monomers**, linked together to form a chain.


Polymer configurations  $\longleftrightarrow$  Trajectories of a random process





# Copolymer and pinning at an interface


A polymer interacting with **two solvents** and with the **interface** that separates them:





# Copolymer and pinning at an interface


A polymer interacting with **two solvents** and with the **interface** that separates them:



- **Copolymer** interaction with the solvents

# Copolymer and pinning at an interface


A polymer interacting with **two solvents** and with the **interface** that separates them:



- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface

# Copolymer and pinning at an interface


A polymer interacting with **two solvents** and with the **interface** that separates them:



- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

# Copolymer and pinning at an interface

A polymer interacting with **two solvents** and with the **interface** that separates them:




- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

**Localization vs. delocalization? Phase transitions?**

# Copolymer and pinning at an interface

A polymer interacting with **two solvents** and with the **interface** that separates them:




- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

**Localization** vs. **delocalization**? **Phase transitions**?

Recent results: very good comprehension (survey: [Giacomin '07])


# Multi-interface media

More general environments: a **multi-interface** medium



# Multi-interface media


More general environments: a **multi-interface** medium



- **Copolymer** case studied by [den Hollander & Wüthrich 04]: some path results for  $\log \log N \ll T_N \ll \log N$  ( $N$  = polymer size)

# Multi-interface media

More general environments: a **multi-interface** medium



- ▶ **Copolymer** case studied by [den Hollander & Wüthrich 04]: some path results for  $\log \log N \ll T_N \ll \log N$  ( $N$  = polymer size)
- ▶ We focus on the **pinning** case: **homogeneous** interaction (attractive or repulsive), **general**  $T_N$ . **Path behavior**?




# Polymer in a slit

Recent physical literature:

Polymer **confined** between two walls and **interacting** with them

- ▶ [Brak, Owkzarek, Rechnitzer, Whittington; J Phys A 2005]
- ▶ [Martin, Orlandini, Owkzarek, Rechnitzer, Whittington; J Phys A 2007]
- ▶ [Owkzarek, Prellbert, Rechnitzer; J Phys A 2008]




# Polymer in a slit

Recent physical literature:


Polymer **confined** between two walls and **interacting** with them

- ▶ [Brak, Owkzarek, Rechnitzer, Whittington; J Phys A 2005]
- ▶ [Martin, Orlandini, Owkzarek, Rechnitzer, Whittington; J Phys A 2007]
- ▶ [Owkzarek, Prellbert, Rechnitzer; J Phys A 2008]



Attraction/repulsion of interfaces by polymers

# Polymer in a slit



# Outline

## 1. Introduction

What is a polymer?

Interaction with the environment

## 2. The model and the main results

Definition

The free energy

Path results

## 3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach

# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

- ▶ Polymer size  $N \in 2\mathbb{N}$  (number of monomers)

# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

- ▶ Polymer size  $N \in 2\mathbb{N}$  (number of monomers)
- ▶ Interaction strength  $\delta \in \mathbb{R}$  ( $> 0$  attractive,  $< 0$  repulsive)

# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

- ▶ Polymer size  $N \in 2\mathbb{N}$  (number of monomers)
- ▶ Interaction strength  $\delta \in \mathbb{R}$  ( $> 0$  attractive,  $< 0$  repulsive)
- ▶ Interface distance  $T \in 2\mathbb{N}$  (interfaces  $\equiv T\mathbb{Z}$ )



# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

- ▶ Polymer size  $N \in 2\mathbb{N}$  (number of monomers)
- ▶ Interaction strength  $\delta \in \mathbb{R}$  ( $> 0$  attractive,  $< 0$  repulsive)
- ▶ Interface distance  $T \in 2\mathbb{N}$  (interfaces  $\equiv T\mathbb{Z}$ )

Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP} := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T}$$

# Definition of the model

Ingredients:

- ▶ Simple symmetric random walk  $S = \{S_n\}_{n \geq 0}$  on  $\mathbb{Z}$ :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$


with  $\{X_i\}_i$  i.i.d. and  $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$ .

- ▶ Polymer size  $N \in 2\mathbb{N}$  (number of monomers)
- ▶ Interaction strength  $\delta \in \mathbb{R}$  ( $> 0$  attractive,  $< 0$  repulsive)
- ▶ Interface distance  $T \in 2\mathbb{N}$  (interfaces  $\equiv T\mathbb{Z}$ )

Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP}(S) := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T} = \frac{1}{Z_{N,\delta}^T} \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\},$$

# Definition of the model



Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP}(S) := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T} = \frac{1}{Z_{N,\delta}^T} \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\},$$

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$   
(**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$   
(**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

Some questions on  $\mathbf{P}_{N,\delta}^{T_N}$  for large  $N$

- What is the typical size of  $S_N$ ?

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$  (**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

Some questions on  $\mathbf{P}_{N,\delta}^{T_N}$  for large  $N$

- ▶ What is the typical size of  $S_N$ ?
- ▶ How many times does the polymer visit the interfaces?

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$  (**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

Some questions on  $\mathbf{P}_{N,\delta}^{T_N}$  for large  $N$

- ▶ What is the typical size of  $S_N$ ?
- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?



# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$  (**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

Some questions on  $\mathbf{P}_{N,\delta}^{T_N}$  for large  $N$

- ▶ What is the typical size of  $S_N$ ?
- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?
- ▶ What is the interplay between  $\delta$  and  $T = \{T_N\}_N$ ?

# Interpretation and some questions

The law  $\mathbf{P}_{N,\delta}^T$  describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of  $\mathbf{P}_{N,\delta}^T$  as  $N \rightarrow \infty$   
(**thermodynamic limit**), for **fixed**  $\delta \in \mathbb{R}$  and for **arbitrary**  $T = T_N$

Some questions on  $\mathbf{P}_{N,\delta}^{T_N}$  for large  $N$

- ▶ What is the typical size of  $S_N$ ?
- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?
- ▶ What is the interplay between  $\delta$  and  $T = \{T_N\}_N$ ?

Penalisation of a random walk

# The free energy

The **free energy**  $\phi(\delta, \{T_n\}_n)$  encodes the exponential asymptotic behavior of the **partition function**  $Z_{N,\delta}^{T_N}$  as  $N \rightarrow \infty$ :

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

# The free energy

The **free energy**  $\phi(\delta, \{T_n\}_n)$  encodes the exponential asymptotic behavior of the **partition function**  $Z_{N,\delta}^{T_N}$  as  $N \rightarrow \infty$ :

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[ \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right] = E[\exp \{\delta L_N\}]$$

$L_N := \#\{i \leq N : S_i \in T\mathbb{Z}\}$  **number of visits** to the interfaces

# The free energy

The **free energy**  $\phi(\delta, \{T_n\}_n)$  encodes the exponential asymptotic behavior of the **partition function**  $Z_{N,\delta}^{T_N}$  as  $N \rightarrow \infty$ :

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[ \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right] = E[\exp \{\delta L_N\}]$$

$L_N := \#\{i \leq N : S_i \in T\mathbb{Z}\}$  **number of visits** to the interfaces

$\phi$  is a **generating function**: if  $\phi'(\delta, \{T_n\}_n)$  exists

$$\phi'(\delta, \{T_n\}_n) = \lim_{N \rightarrow \infty} \mathbf{E}_{N,\delta}^{T_N} \left( \frac{L_N}{N} \right) \quad \boxed{L_N \sim \phi'(\delta, \{T_n\}_n) \cdot N}$$

# The free energy

The **free energy**  $\phi(\delta, \{T_n\}_n)$  encodes the exponential asymptotic behavior of the **partition function**  $Z_{N,\delta}^{T_N}$  as  $N \rightarrow \infty$ :

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[ \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right] = E[\exp \{\delta L_N\}]$$

$L_N := \#\{i \leq N : S_i \in T\mathbb{Z}\}$  **number of visits** to the interfaces

$\phi$  is a **generating function**: if  $\phi'(\delta, \{T_n\}_n)$  exists

$$\phi'(\delta, \{T_n\}_n) = \lim_{N \rightarrow \infty} \mathbf{E}_{N,\delta}^{T_N} \left( \frac{L_N}{N} \right) \quad \boxed{L_N \sim \phi'(\delta, \{T_n\}_n) \cdot N}$$

If  $\phi(\delta, \{T_n\}_n)$  is non-analytic in  $\delta \in \mathbb{R}$  there is a **phase transition**

# The free energy: characterization

We may assume that  $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$  exists

# The free energy: characterization

We may assume that  $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$  exists

Let  $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$  and  $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$   
(Also  $T = \infty$ : first return to zero of the SSRW)



# The free energy: characterization

We may assume that  $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$  exists

Let  $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$  and  $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$   
 (Also  $T = \infty$ : first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \phi(\delta, T_\infty) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

# The free energy: characterization

We may assume that  $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$  exists

Let  $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$  and  $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$   
 (Also  $T = \infty$ : first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \phi(\delta, T_\infty) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

If  $T_N \rightarrow T_\infty < \infty$  (same as  $T_N \equiv T_\infty$ )

►  $\phi(\delta, T_\infty)$  is analytic on  $\mathbb{R}$ : no phase transitions

# The free energy: characterization

We may assume that  $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$  exists

Let  $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$  and  $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$   
 (Also  $T = \infty$ : first return to zero of the SSRW)

## Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \phi(\delta, T_\infty) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

If  $T_N \rightarrow T_\infty < \infty$  (same as  $T_N \equiv T_\infty$ )

- ▶  $\phi(\delta, T_\infty)$  is analytic on  $\mathbb{R}$ : no phase transitions
- ▶  $\phi'(\delta, T_\infty) > 0$  for every  $\delta \in \mathbb{R}$ : positive density of contacts

$$L_N \sim \phi'(\delta, T_\infty) \cdot N \quad (\text{conj. diffusive behavior of } S_N)$$

# The free energy: further results

If  $T_N \rightarrow \infty$

► Phase transition (only) at  $\delta = 0$

► If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \longrightarrow L_N = o(N)$

# The free energy: further results

If  $T_N \rightarrow \infty$

► Phase transition (only) at  $\delta = 0$

- If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \longrightarrow L_N = o(N)$
- If  $\delta > 0$  then  $\phi'(\delta, \infty) > 0 \longrightarrow L_N \sim \phi'(\delta, \infty) \cdot N$

# The free energy: further results

If  $T_N \rightarrow \infty$

- ▶ Phase transition (only) at  $\delta = 0$ 
  - ▶ If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \rightarrow L_N = o(N)$
  - ▶ If  $\delta > 0$  then  $\phi'(\delta, \infty) > 0 \rightarrow L_N \sim \phi'(\delta, \infty) \cdot N$
- ▶ Every  $\{T_n\}_n \rightarrow \infty$  yields the same free energy as if  $T_n \equiv \infty$  (homogeneous pinning model)  $\rightarrow$  same density of visits

# The free energy: further results

If  $T_N \rightarrow \infty$

- ▶ Phase transition (only) at  $\delta = 0$ 
  - ▶ If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \rightarrow L_N = o(N)$
  - ▶ If  $\delta > 0$  then  $\phi'(\delta, \infty) > 0 \rightarrow L_N \sim \phi'(\delta, \infty) \cdot N$
- ▶ Every  $\{T_n\}_n \rightarrow \infty$  yields the same free energy as if  $T_n \equiv \infty$  (homogeneous pinning model)  $\rightarrow$  same density of visits
- ▶ Same path behavior? NO!

# The free energy: further results

If  $T_N \rightarrow \infty$

- ▶ Phase transition (only) at  $\delta = 0$ 
  - ▶ If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \rightarrow L_N = o(N)$
  - ▶ If  $\delta > 0$  then  $\phi'(\delta, \infty) > 0 \rightarrow L_N \sim \phi'(\delta, \infty) \cdot N$
- ▶ Every  $\{T_n\}_n \rightarrow \infty$  yields the same free energy as if  $T_n \equiv \infty$  (homogeneous pinning model)  $\rightarrow$  same density of visits
- ▶ Same path behavior? NO!
- ▶ If  $\delta < 0$  then  $Z_{N,\delta}^{T_N} = \exp(o(N))$ .



# The free energy: further results

If  $T_N \rightarrow \infty$

- ▶ Phase transition (only) at  $\delta = 0$ 
  - ▶ If  $\delta \leq 0$  then  $\phi(\delta, \infty) = \phi'(\delta, \infty) \equiv 0 \rightarrow L_N = o(N)$
  - ▶ If  $\delta > 0$  then  $\phi'(\delta, \infty) > 0 \rightarrow L_N \sim \phi'(\delta, \infty) \cdot N$
- ▶ Every  $\{T_n\}_n \rightarrow \infty$  yields the same free energy as if  $T_n \equiv \infty$  (homogeneous pinning model)  $\rightarrow$  same density of visits
- ▶ Same path behavior? NO!
- ▶ If  $\delta < 0$  then  $Z_{N,\delta}^{T_N} = \exp(o(N))$ . In fact

$$Z_{N,\delta}^{T_N} \approx \frac{(\text{const.})}{N^{3/2}} f\left(\frac{\sqrt{N}}{T_N}\right) g\left(\frac{N^{1/3}}{T_N}\right),$$

improving known results for the polymer in a slit.

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )

$$\frac{S_N}{C_\delta (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}} \Rightarrow N(0, 1)$$

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$  then  $S_N \asymp T_N$

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$  then  $S_N \asymp T_N$

$$\frac{S_N}{T_N} \Rightarrow S_\Gamma \quad \Gamma \sim \text{Poisson}(f(\delta, \zeta))$$



# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$  then  $S_N \asymp T_N$
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow +\infty$  then  $S_N \asymp O(1)$

# Path results: the attractive case $\delta > 0$

Assume  $\delta > 0$  and  $T_N \rightarrow \infty$ . Since  $\phi'(\delta, \infty) > 0$ , the polymer visits the interfaces a **positive fraction of times**:  $L_N \sim \phi'(\delta, \infty) \cdot N$ .

What is the typical size of  $S_N$ ?

Not.:  $S_N \asymp \alpha_N$  means  $S_N/\alpha_N$  is tight and  $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$


## Theorem ([CP1])

For every  $\delta > 0$  there exists  $c_\delta > 0$  such that under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$  then  $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$   
(note that  $T_N \ll S_N \ll \sqrt{N}$ )
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$  then  $S_N \asymp T_N$
- ▶ If  $T_N - \frac{1}{c_\delta} \log N \rightarrow +\infty$  then  $S_N \asymp O(1)$

$$\lim_{L \rightarrow \infty} \sup_{N \in 2\mathbb{N}} \mathbf{P}_{N,\delta}^{T_N}(|S_N| > L) = 0$$

# Path results: the attractive case $\delta > 0$



- Sub-diffusive scaling ( $T_N \rightarrow \infty$ )
- Transition at  $T_N \approx \log N$

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

## Theorem ([CP2])

For every  $\delta < 0$  we have under  $\mathbf{P}_{N,\delta}^{T_N}$ :

► If  $T_N \ll N^{1/3}$  then  $S_N \asymp \sqrt{N/T_N} \gg T_N$

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

## Theorem ([CP2])

For every  $\delta < 0$  we have under  $\mathbf{P}_{N,\delta}^{T_N}$ :

► If  $T_N \ll N^{1/3}$  then  $S_N \asymp \sqrt{N/T_N} \gg T_N$

$$c_1 P(a < Z \leq b) \leq \mathbf{P}_{N,\delta}^{T_N} \left( a < \frac{S_N}{C_\delta \sqrt{\frac{N}{T_N}}} \leq b \right) \leq c_2 P(a < Z \leq b)$$

with  $Z \sim N(0, 1)$



# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

## Theorem ([CP2])

For every  $\delta < 0$  we have under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N \ll N^{1/3}$  then  $S_N \asymp \sqrt{N/T_N} \gg T_N$
- ▶ If  $(\text{const.})N^{1/3} \leq T_N \leq (\text{const.})\sqrt{N}$  then  $S_N \asymp T_N$

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

## Theorem ([CP2])

For every  $\delta < 0$  we have under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N \ll N^{1/3}$  then  $S_N \asymp \sqrt{N/T_N} \gg T_N$
- ▶ If  $(\text{const.})N^{1/3} \leq T_N \leq (\text{const.})\sqrt{N}$  then  $S_N \asymp T_N$
- ▶ If  $T_N \gg \sqrt{N}$  then  $S_N \asymp \sqrt{N}$

# Path results: the repulsive case $\delta < 0$

Assume  $\delta < 0$  and  $T_N \rightarrow \infty$ . The polymer visits the interfaces a null fraction of times:  $L_N = o(N)$ .

What is the typical size of  $S_N$ ?

Notation:  $S_N \asymp \alpha_N$  as above •  $a_n \ll b_n \iff a_n/b_n \rightarrow 0$


## Theorem ([CP2])

For every  $\delta < 0$  we have under  $\mathbf{P}_{N,\delta}^{T_N}$ :

- ▶ If  $T_N \ll N^{1/3}$  then  $S_N \asymp \sqrt{N/T_N} \gg T_N$
- ▶ If  $(\text{const.})N^{1/3} \leq T_N \leq (\text{const.})\sqrt{N}$  then  $S_N \asymp T_N$
- ▶ If  $T_N \gg \sqrt{N}$  then  $S_N \asymp \sqrt{N}$

- ▶ If  $T_N \sim (\text{const.})N^{1/3} \longrightarrow O(1)$  visited interfaces.
- ▶ If  $T_N \gg N^{1/3} \longrightarrow 1$  visited interface,  $L_N = O(1)$ .

# Path results: the repulsive case $\delta < 0$



- Sub-diffusive if  $1 \ll T_N \ll \sqrt{N}$
- Transitions  $T_N \approx N^{1/3}, \sqrt{N}$

# Outline

## 1. Introduction

What is a polymer?

Interaction with the environment

## 2. The model and the main results

Definition

The free energy

Path results

## 3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach


# An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero)  $\longrightarrow$  for  $\delta > 0$  it is **localized at the interface**

# An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero)  $\rightarrow$  for  $\delta > 0$  it is **localized at the interface**


$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$



# An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero)  $\rightarrow$  for  $\delta > 0$  it is **localized at the interface**

$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$




- If  $T_N \gg \log N$  **nothing changes**: polymer localized at zero



# An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero)  $\rightarrow$  for  $\delta > 0$  it is **localized at the interface**

$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$




- ▶ If  $T_N \gg \log N$  nothing changes: polymer localized at zero
- ▶ If  $T_N \ll \log N$  it is worth to visit different interfaces:  
**sub-diffusive behavior**

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?

# An heuristic view: the repulsive case $\delta < 0$


How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?



Under the **simple random walk** law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?




Under the simple random walk law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

Under the **polymer measure**  $\mathbf{P}_{N,\delta}^T$  with  $\delta < 0$   $\hat{\tau}_1^T \approx T^3$  (repulsion)

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?




Under the simple random walk law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

Under the polymer measure  $\mathbf{P}_{N,\delta}^T$  with  $\delta < 0$   $\hat{\tau}_1^T \approx T^3$  (repulsion)

- ▶ if  $T_N^3 \ll N$  infinitely many interfaces:  $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?




Under the simple random walk law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

Under the polymer measure  $\mathbf{P}_{N,\delta}^T$  with  $\delta < 0$   $\hat{\tau}_1^T \approx T^3$  (repulsion)

- ▶ if  $T_N^3 \ll N$  **infinitely many** interfaces:  $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if  $T_N^3 \gg N$  **confinement**:  $|S_i| < T_N$  for all  $i \leq N$ :

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?




Under the simple random walk law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

Under the polymer measure  $\mathbf{P}_{N,\delta}^T$  with  $\delta < 0$   $\hat{\tau}_1^T \approx T^3$  (repulsion)

- ▶ if  $T_N^3 \ll N$  **infinitely many** interfaces:  $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if  $T_N^3 \gg N$  **confinement**:  $|S_i| < T_N$  for all  $i \leq N$ :
  - $S_N \approx T_N$  (if  $T_N \leq \sqrt{N}$ )

# An heuristic view: the repulsive case $\delta < 0$

How long is the time  $\hat{\tau}_1^T$  to reach level  $\pm T$ ?



Under the simple random walk law  $\hat{\tau}_1^T \approx T^2$  (diffusivity)

Under the polymer measure  $\mathbf{P}_{N,\delta}^T$  with  $\delta < 0$   $\hat{\tau}_1^T \approx T^3$  (repulsion)

- ▶ if  $T_N^3 \ll N$  **infinitely many** interfaces:  $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if  $T_N^3 \gg N$  **confinement**:  $|S_i| < T_N$  for all  $i \leq N$ :
  - $S_N \approx T_N$  (if  $T_N \leq \sqrt{N}$ )
  - $S_N \approx \sqrt{N}$  (if  $T_N \gg \sqrt{N}$ )



# A renewal theory approach

Let  $\tau_1^T, \tau_2^T, \tau_3^T \dots$  be the points at which  $S_n$  visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

# A renewal theory approach

Let  $\tau_1^T, \tau_2^T, \tau_3^T \dots$  be the points at which  $S_n$  visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

Under the **simple random walk** law  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is a classical **renewal process** with explicit law

$$q_T(n) := P(\tau_1^T = n) \approx \frac{1}{(\min\{n, T^2\})^{3/2}} e^{-\frac{\pi^2}{2T^2}n}$$

# A renewal theory approach

Let  $\tau_1^T, \tau_2^T, \tau_3^T \dots$  be the points at which  $S_n$  visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

Under the **simple random walk** law  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is a classical **renewal process** with explicit law

$$q_T(n) := P(\tau_1^T = n) \approx \frac{1}{(\min\{n, T^2\})^{3/2}} e^{-\frac{\pi^2}{2T^2}n}$$

Note that under  $P$

$$\tau_1^T \approx \begin{cases} O(1) \text{ with probab. } 1 - \frac{1}{T} & [q_T(n) \approx \frac{1}{n^{3/2}}] \\ O(T^2) \text{ with probab. } \frac{1}{T} & [q_T(n) \approx \frac{1}{T^3} e^{-\frac{\pi^2}{2T^2}n}] \end{cases}$$

# A renewal theory approach

Under the **polymer measure**  $\mathbf{P}_{N,\delta}^T$  the process  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is not even time-homogeneous ...

# A renewal theory approach

Under the **polymer measure**  $\mathbf{P}_{N,\delta}^T$  the process  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is not even time-homogeneous ... however for large  $N$  it is nearly a **renewal process** with a different law  $\mathcal{P}_{\delta,T}$

$$K_{\delta,T}(n) := \mathcal{P}_{\delta,T}(\tau_1^T = n) = e^\delta P(\tau_1^T = n) e^{-\phi(\delta,T)n}$$

# A renewal theory approach

Under the **polymer measure**  $\mathbf{P}_{N,\delta}^T$  the process  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is not even time-homogeneous ... however for large  $N$  it is nearly a **renewal process** with a different law  $\mathcal{P}_{\delta,T}$

$$K_{\delta,T}(n) := \mathcal{P}_{\delta,T}(\tau_1^T = n) = e^\delta P(\tau_1^T = n) e^{-\phi(\delta,T)n}$$

For  $\delta > 0$ , we have  $\phi(\delta, T) \rightarrow \phi(\delta, \infty) > 0$  as  $T \rightarrow \infty$ , hence

$$K_{\delta,T}(n) \approx e^{-\phi(\delta,\infty)n}$$

# A renewal theory approach

Under the **polymer measure**  $\mathbf{P}_{N,\delta}^T$  the process  $\{\tau_n^T\}_{n \in \mathbb{N}}$  is not even time-homogeneous ... however for large  $N$  it is nearly a **renewal process** with a different law  $\mathcal{P}_{\delta,T}$

$$K_{\delta,T}(n) := \mathcal{P}_{\delta,T}(\tau_1^T = n) = e^\delta P(\tau_1^T = n) e^{-\phi(\delta,T)n}$$

For  $\delta > 0$ , we have  $\phi(\delta, T) \rightarrow \phi(\delta, \infty) > 0$  as  $T \rightarrow \infty$ , hence

$$K_{\delta,T}(n) \approx e^{-\phi(\delta,\infty)n}$$

For  $\delta < 0$ , we have  $\phi(\delta, T) \approx -\frac{\pi^2}{2T^2} + \frac{C_\delta}{T^3}$  as  $T \rightarrow \infty$ , hence

$$\tau_1^T \approx \begin{cases} O(1) \text{ with probab. } e^\delta & [K_{\delta,T}(n) \approx \frac{1}{n^{3/2}}] \\ O(T^3) \text{ with probab. } 1 - e^\delta & [K_{\delta,T}(n) \approx \frac{1}{T^3} e^{-\frac{C_\delta}{T^3}n}] \end{cases}$$

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^T \implies \mathcal{P}_{\delta,T}$ )



# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^{T_N} \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^T \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

1. Study  $\tau^{T_N} \cap [0, N]$  under  $\mathcal{P}_{\delta,T_N}$

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^{T_N} \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

1. Study  $\tau^{T_N} \cap [0, N]$  under  $\mathcal{P}_{\delta,T_N}$
2. Transfer the results to  $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$  that is to  $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$  (hard part)

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^{T_N} \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

1. Study  $\tau^{T_N} \cap [0, N]$  under  $\mathcal{P}_{\delta,T_N}$
2. Transfer the results to  $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$  that is to  $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$  (hard part)
3. Remove the conditioning on  $\{N \in \tau^{T_N}\}$

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^{T_N} \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

1. Study  $\tau^{T_N} \cap [0, N]$  under  $\mathcal{P}_{\delta,T_N}$
2. Transfer the results to  $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$  that is to  $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$  (hard part)
3. Remove the conditioning on  $\{N \in \tau^{T_N}\}$

► Good estimates on  $q_T(n)$  and on the free energy  $\phi(\delta, T)$

# Strategy of the proof

The law of  $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$  is the same

under  $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$  and  $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

$\mathcal{P}_{\delta,T}$  does not depend explicitly on  $N$  ( $\mathbf{P}_{N,\delta}^{T_N} \implies \mathcal{P}_{\delta,T}$ )

However we want to study  $\mathbf{P}_{N,\delta}^{T_N}$  with  $T_N$  varying with  $N$

1. Study  $\tau^{T_N} \cap [0, N]$  under  $\mathcal{P}_{\delta,T_N}$
2. Transfer the results to  $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$  that is to  $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$  (hard part)
3. Remove the conditioning on  $\{N \in \tau^{T_N}\}$

- ▶ Good estimates on  $q_T(n)$  and on the free energy  $\phi(\delta, T)$
- ▶ Uniform renewal theorems

Thanks.