

A Polymer in a Multi-Interface Medium

Francesco Caravenna

`caravenna@math.unipd.it`

Università degli Studi di Padova

Humboldt Universität zu Berlin ~ November 12, 2008

References

- ▶ [CP1] F. Caravenna and N. Pétrélis
A polymer in a multi-interface medium
Preprint (2007), arXiv.org: 0712.3426
- ▶ [CP2] F. Caravenna and N. Pétrélis
Depinning of a polymer in a multi-interface medium
In preparation.

Outline

1. Introduction

What is a polymer?

Polymers and probability

2. The model and the main results

Definition

The free energy

Path results

3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach

Outline

1. Introduction

What is a polymer?

Polymers and probability

2. The model and the main results

Definition

The free energy

Path results

3. Techniques and ideas from the proof

Some heuristics


A renewal theory approach

What is a polymer?

A **polymer** is a large molecule composed of repeating smaller units, called **monomers**, linked together to form a chain.


What is a polymer?

A **polymer** is a large molecule composed of repeating smaller units, called **monomers**, linked together to form a chain.



What is a polymer?

A **polymer** is a large molecule composed of repeating smaller units, called **monomers**, linked together to form a chain.




Typical examples:

- ▶ DNA, RNA
- ▶ Proteins
- ▶ Plastics

Important research topic in chemistry, physics, biology...

What is a polymer?

A **polymer** is a large molecule composed of repeating smaller units, called **monomers**, linked together to form a chain.



Typical examples:

- ▶ DNA, RNA
- ▶ Proteins
- ▶ Plastics

Important research topic in chemistry, physics, biology...

...and **mathematics** too

Polymers and probability


Polymer configurations \longleftrightarrow Trajectories of a random process

Polymers and probability

Polymer configurations \longleftrightarrow Trajectories of a random process

Self-avoiding walks

Simple symmetric random walk on \mathbb{Z}^d
conditioned to visit each site at most
once \longrightarrow very difficult!




Polymers and probability

Polymer configurations \longleftrightarrow Trajectories of a random process


Self-avoiding walks

Simple symmetric random walk on \mathbb{Z}^d
conditioned to visit each site at most
once \longrightarrow very difficult!




Directed walks

Processes with a deterministic component: (n, S_n) where S_n is the simple symmetric random walk on \mathbb{Z}^{d-1}
 → tractable models




Interaction with the environment

A polymer interacting with **two solvents** and with the **interface** that separates them:



Interaction with the environment


A polymer interacting with **two solvents** and with the **interface** that separates them:



- **Copolymer** interaction with the solvents

Interaction with the environment


A polymer interacting with **two solvents** and with the **interface** that separates them:



- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface

Interaction with the environment


A polymer interacting with **two solvents** and with the **interface** that separates them:



- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

Interaction with the environment

A polymer interacting with **two solvents** and with the **interface** that separates them:




- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

Localization vs. delocalization? Phase transitions?

Interaction with the environment

A polymer interacting with **two solvents** and with the **interface** that separates them:




- ▶ **Copolymer** interaction with the solvents
- ▶ **Pinning** interaction with the interface
- ▶ Both interactions may be **inhomogeneous** and **disordered**

Localization vs. delocalization? Phase transitions?

Recent results: very good comprehension (survey: [Giacomin '07])


Multi-interface media

More general environments: a **multi-interface** medium



Multi-interface media


More general environments: a **multi-interface** medium



- **Copolymer** case studied by [den Hollander & Wüthrich 04]: some path results for $\log \log N \ll T_N \ll \log N$ (N = polymer size)

Multi-interface media


More general environments: a **multi-interface** medium



- ▶ **Copolymer** case studied by [den Hollander & Wüthrich 04]: some path results for $\log \log N \ll T_N \ll \log N$ (N = polymer size)
- ▶ We focus on the **pinning** case: **homogeneous** interaction (attractive or repulsive), **general** T_N . **Path behavior?**

Multi-interface media

More general environments: a **multi-interface** medium



- **Copolymer** case studied by [den Hollander & Wüthrich 04]: some path results for $\log \log N \ll T_N \ll \log N$ (N = polymer size)
- We focus on the **pinning** case: **homogeneous** interaction (attractive or repulsive), **general** T_N . **Path behavior?**

[Copolymer in an **emulsion**: den Hollander, P  tr  lis, Whittington, W  thrich]

Outline

1. Introduction

What is a polymer?

Polymers and probability

2. The model and the main results

Definition

The free energy

Path results

3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

- ▶ Polymer size $N \in 2\mathbb{N}$ (number of monomers)

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

- ▶ Polymer size $N \in 2\mathbb{N}$ (number of monomers)
- ▶ Interaction strength $\delta \in \mathbb{R}$ (> 0 attractive, < 0 repulsive)

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

- ▶ Polymer size $N \in 2\mathbb{N}$ (number of monomers)
- ▶ Interaction strength $\delta \in \mathbb{R}$ (> 0 attractive, < 0 repulsive)
- ▶ Interface distance $T \in 2\mathbb{N}$ (interfaces $\equiv T\mathbb{Z}$)

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$

with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

- ▶ Polymer size $N \in 2\mathbb{N}$ (number of monomers)
- ▶ Interaction strength $\delta \in \mathbb{R}$ (> 0 attractive, < 0 repulsive)
- ▶ Interface distance $T \in 2\mathbb{N}$ (interfaces $\equiv T\mathbb{Z}$)

Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP} := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T}$$

Definition of the model

Ingredients:

- ▶ Simple symmetric random walk $S = \{S_n\}_{n \geq 0}$ on \mathbb{Z} :

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n,$$


with $\{X_i\}_i$ i.i.d. and $P(X_i = +1) = P(X_i = -1) = \frac{1}{2}$.

- ▶ Polymer size $N \in 2\mathbb{N}$ (number of monomers)
- ▶ Interaction strength $\delta \in \mathbb{R}$ (> 0 attractive, < 0 repulsive)
- ▶ Interface distance $T \in 2\mathbb{N}$ (interfaces $\equiv T\mathbb{Z}$)

Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP}(S) := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T} = \frac{1}{Z_{N,\delta}^T} \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\},$$

Definition of the model



Polymer measure:

$$\frac{d\mathbf{P}_{N,\delta}^T(S)}{dP}(S) := \frac{\exp(H_{N,\delta}^T(S))}{Z_{N,\delta}^T} = \frac{1}{Z_{N,\delta}^T} \exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\},$$

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of $\mathbf{P}_{N,\delta}^T$ as $N \rightarrow \infty$
(**thermodynamic limit**), for **fixed** $\delta \in \mathbb{R}$ and for **arbitrary** $T = T_N$

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of $\mathbf{P}_{N,\delta}^T$ as $N \rightarrow \infty$ (**thermodynamic limit**), for **fixed** $\delta \in \mathbb{R}$ and for **arbitrary** $T = T_N$

Some questions on $\mathbf{P}_{N,\delta}^{T_N}$ for large N

- How many times does the polymer visit the interfaces?

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of $\mathbf{P}_{N,\delta}^T$ as $N \rightarrow \infty$
(**thermodynamic limit**), for **fixed** $\delta \in \mathbb{R}$ and for **arbitrary** $T = T_N$

Some questions on $\mathbf{P}_{N,\delta}^{T_N}$ for large N

- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of $\mathbf{P}_{N,\delta}^T$ as $N \rightarrow \infty$ (**thermodynamic limit**), for **fixed** $\delta \in \mathbb{R}$ and for **arbitrary** $T = T_N$

Some questions on $\mathbf{P}_{N,\delta}^{T_N}$ for large N

- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?
- ▶ What is the typical size of S_N ?

Interpretation and some questions

The law $\mathbf{P}_{N,\delta}^T$ describes the **statistical distribution** of the configurations of the polymer, given the external conditions

Interested in the properties of $\mathbf{P}_{N,\delta}^T$ as $N \rightarrow \infty$ (**thermodynamic limit**), for **fixed** $\delta \in \mathbb{R}$ and for **arbitrary** $T = T_N$

Some questions on $\mathbf{P}_{N,\delta}^{T_N}$ for large N

- ▶ How many times does the polymer visit the interfaces?
- ▶ How many different interfaces are visited?
- ▶ What is the typical size of S_N ?
- ▶ What is the interplay between δ and $T = \{T_N\}_N$?

The free energy

The **free energy** $\phi(\delta, \{T_n\}_n)$ encodes the exponential asymptotic behavior of the **partition function** $Z_{N,\delta}^{T_N}$ as $N \rightarrow \infty$:

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

The free energy

The **free energy** $\phi(\delta, \{T_n\}_n)$ encodes the exponential asymptotic behavior of the **partition function** $Z_{N,\delta}^{T_N}$ as $N \rightarrow \infty$:

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[\exp \left\{ H_{N,\delta}^{T_N} \right\} \right] = E \left[\exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right]$$

The free energy

The **free energy** $\phi(\delta, \{T_n\}_n)$ encodes the exponential asymptotic behavior of the **partition function** $Z_{N,\delta}^{T_N}$ as $N \rightarrow \infty$:

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[\exp \left\{ H_{N,\delta}^{T_N} \right\} \right] = E \left[\exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right]$$

Why should one look at the free energy? Introducing the **number of visits** to the interfaces $L_N := \#\{i \leq N : S_i \in T\mathbb{Z}\}$ we have

$$\frac{\partial}{\partial \delta} \phi(\delta, \{T_n\}_n) = \lim_{N \rightarrow \infty} \mathbf{E}_{N,\delta}^{T_N} \left(\frac{1}{N} L_N \right) \quad \text{if } \frac{\partial}{\partial \delta} \phi \text{ exists}$$

The free energy

The **free energy** $\phi(\delta, \{T_n\}_n)$ encodes the exponential asymptotic behavior of the **partition function** $Z_{N,\delta}^{T_N}$ as $N \rightarrow \infty$:

$$\phi(\delta, \{T_n\}_n) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\delta}^{T_N} \quad (\text{super-additivity} + \dots)$$

$$Z_{N,\delta}^{T_N} := E \left[\exp \left\{ H_{N,\delta}^{T_N} \right\} \right] = E \left[\exp \left\{ \delta \sum_{i=1}^N \mathbf{1}_{\{S_i \in T\mathbb{Z}\}} \right\} \right]$$

Why should one look at the free energy? Introducing the **number of visits** to the interfaces $L_N := \#\{i \leq N : S_i \in T\mathbb{Z}\}$ we have

$$\frac{\partial}{\partial \delta} \phi(\delta, \{T_n\}_n) = \lim_{N \rightarrow \infty} \mathbf{E}_{N,\delta}^{T_N} \left(\frac{1}{N} L_N \right) \quad \text{if } \frac{\partial}{\partial \delta} \phi \text{ exists}$$

If $\phi(\delta, \{T_n\}_n)$ is non-analytic in $\delta \in \mathbb{R}$ there is a **phase transition**

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

Let $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$ and $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$
(Also $T = \infty$: first return to zero of the SSRW)

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

Let $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$ and $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$
 (Also $T = \infty$: first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

Let $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$ and $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$
 (Also $T = \infty$: first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

► If $T_\infty < \infty$ then $\phi(\delta, T_\infty)$ is analytic: **no phase transitions**

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

Let $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$ and $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$
 (Also $T = \infty$: first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

- ▶ If $T_\infty < \infty$ then $\phi(\delta, T_\infty)$ is analytic: **no phase transitions**
- ▶ If $T_\infty = \infty$ phase transition (only) at $\delta = 0$

The free energy

We may assume that $\lim_{N \rightarrow \infty} T_N =: T_\infty \in 2\mathbb{N} \cup \{+\infty\}$ exists

Let $\tau_1^T := \inf \{n > 0 : S_n \in \{\pm T, 0\}\}$ and $Q_T(\lambda) := E(e^{-\lambda \tau_1^T})$
(Also $T = \infty$: first return to zero of the SSRW)

Theorem ([CP1])

$$\phi(\delta, \{T_n\}_n) = \begin{cases} (Q_{T_\infty})^{-1}(e^{-\delta}) & \text{if } T_\infty < +\infty \\ (Q_\infty)^{-1}(e^{-\delta} \wedge 1) & \text{if } T_\infty = +\infty \end{cases}$$

- ▶ If $T_\infty < \infty$ then $\phi(\delta, T_\infty)$ is analytic: **no phase transitions**
- ▶ If $T_\infty = \infty$ phase transition (only) at $\delta = 0$
- ▶ Every $\{T_n\}_n \rightarrow \infty$ yields **the same free energy** as if $T_n \equiv \infty$
(homogeneous pinning model) \longrightarrow **same density of visits**

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)

$$\frac{S_N}{C_\delta (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}} \Rightarrow N(0, 1)$$

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$

$$\frac{S_N}{T_N} \implies S_\Gamma \quad \Gamma \sim \text{Poisson}(f(\delta, \zeta))$$

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$

Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow +\infty$ then $S_N \asymp O(1)$ and $\hat{L}_N \rightarrow 0$

Path results: the attractive case $\delta > 0$

Assume $\delta > 0$ and $T_N \rightarrow \infty$. Since $\phi'(\delta, \infty) > 0$, the polymer visits the interfaces a **positive fraction of times**: $L_N \sim \phi'(\delta, \infty) \cdot N$.

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Not.: $S_N \asymp \alpha_N$ means S_N/α_N is tight and $\mathbf{P}_{N,\delta}^{T_N}(|S_N/\alpha_N| \geq \varepsilon) \geq \varepsilon$


Theorem ([CP1])

For every $\delta > 0$ there exists $c_\delta > 0$ such that under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow -\infty$ then $S_N \asymp (e^{-\frac{c_\delta}{2} T_N} T_N) \sqrt{N}$ and $\hat{L}_N \asymp N/e^{c_\delta T_N}$ (note that $T_N \ll S_N \ll \sqrt{N}$ and $\hat{L}_N \rightarrow \infty$)
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow \zeta \in \mathbb{R}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$
- ▶ If $T_N - \frac{1}{c_\delta} \log N \rightarrow +\infty$ then $S_N \asymp O(1)$ and $\hat{L}_N \rightarrow 0$


$$\lim_{L \rightarrow \infty} \sup_{N \in 2\mathbb{N}} \mathbf{P}_{N,\delta}^{T_N}(|S_N| > L) = 0$$

Path results: the attractive case $\delta > 0$



- Sub-diffusive scaling ($T_N \rightarrow \infty$)
- Transition at $T_N \approx \log N$

Path results: the attractive case $\delta > 0$



- Sub-diffusive scaling ($T_N \rightarrow \infty$)
- Transition at $T_N \approx \log N$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Theorem ([CP2])

For every $\delta < 0$ we have under $\mathbf{P}_{N,\delta}^{T_N}$:

► If $T_N \ll N^{1/3}$ then $S_N \asymp \sqrt{N/T_N}$ and $\hat{L}_N \asymp N/(T_N)^3$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Theorem ([CP2])

For every $\delta < 0$ we have under $\mathbf{P}_{N,\delta}^{T_N}$:

► If $T_N \ll N^{1/3}$ then $S_N \asymp \sqrt{N/T_N}$ and $\hat{L}_N \asymp N/(T_N)^3$

$$c_1 P(a < Z \leq b) \leq \mathbf{P}_{N,\delta}^{T_N} \left(a < \frac{S_N}{C_\delta \sqrt{\frac{N}{T_N}}} \leq b \right) \leq c_2 P(a < Z \leq b)$$

with $Z \sim N(0, 1)$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Theorem ([CP2])

For every $\delta < 0$ we have under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N \ll N^{1/3}$ then $S_N \asymp \sqrt{N/T_N}$ and $\hat{L}_N \asymp N/(T_N)^3$
- ▶ If $T_N \sim (\text{const.})N^{1/3}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?

Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Theorem ([CP2])

For every $\delta < 0$ we have under $\mathbf{P}_{N,\delta}^{T_N}$:

- ▶ If $T_N \ll N^{1/3}$ then $S_N \asymp \sqrt{N/T_N}$ and $\hat{L}_N \asymp N/(T_N)^3$
- ▶ If $T_N \sim (\text{const.})N^{1/3}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$
- ▶ If $N^{1/3} \ll T_N \leq \sqrt{N}$ then $S_N \asymp T_N$ and $\hat{L}_N \rightarrow 0$

Path results: the repulsive case $\delta < 0$

What happens if $\delta < 0$? ($T_N \rightarrow \infty$ for convenience)

The polymer visits the interfaces a **null fraction** $o(N)$ of times

How many jumps \hat{L}_N to **different** interfaces? Typical size of S_N ?


Notation: $S_N \asymp \alpha_N$ as above • $a_n \ll b_n \iff a_n/b_n \rightarrow 0$

Theorem ([CP2])

For every $\delta < 0$ we have under $\mathbf{P}_{N,\delta}^{T_N}$:


- ▶ If $T_N \ll N^{1/3}$ then $S_N \asymp \sqrt{N/T_N}$ and $\hat{L}_N \asymp N/(T_N)^3$
- ▶ If $T_N \sim (\text{const.})N^{1/3}$ then $S_N \asymp T_N$ and $\hat{L}_N \asymp O(1)$
- ▶ If $N^{1/3} \ll T_N \leq \sqrt{N}$ then $S_N \asymp T_N$ and $\hat{L}_N \rightarrow 0$
- ▶ If $T_N \gg \sqrt{N}$ then $S_N \asymp \sqrt{N}$ and $\hat{L}_N \rightarrow 0$

Path results: the repulsive case $\delta < 0$



- Sub-diffusive if $1 \ll T_N \ll \sqrt{N}$
- Transitions $T_N \approx N^{1/3}, \sqrt{N}$

Path results: the repulsive case $\delta < 0$



- Sub-diffusive if $1 \ll T_N \ll \sqrt{N}$
- Transitions $T_N \approx N^{1/3}, \sqrt{N}$

Outline

1. Introduction

What is a polymer?

Polymers and probability

2. The model and the main results

Definition

The free energy

Path results

3. Techniques and ideas from the proof

Some heuristics

A renewal theory approach


An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero) \longrightarrow for $\delta > 0$ it is **localized at the interface**

An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero) \rightarrow for $\delta > 0$ it is **localized at the interface**


$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$



An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero) \rightarrow for $\delta > 0$ it is **localized at the interface**

$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$




- If $T_N \gg \log N$ **nothing changes**: polymer localized at zero

An heuristic view: the attractive case $\delta > 0$

Consider the **simple homogeneous pinning model** (one interface at zero) \rightarrow for $\delta > 0$ it is **localized at the interface**

$$S_N = O(1) \quad \max_{0 \leq i \leq N} |S_i| \asymp \log N$$




- ▶ If $T_N \gg \log N$ nothing changes: polymer localized at zero
- ▶ If $T_N \ll \log N$ it is worth to visit different interfaces:
sub-diffusive behavior

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?

An heuristic view: the repulsive case $\delta < 0$


How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?



Under the **simple random walk** law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?




Under the simple random walk law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

Under the **polymer measure** $\mathbf{P}_{N,\delta}^T$ with $\delta < 0$ $\hat{\tau}_1^T \approx T^3$ (repulsion)

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?




Under the simple random walk law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

Under the polymer measure $\mathbf{P}_{N,\delta}^T$ with $\delta < 0$ $\hat{\tau}_1^T \approx T^3$ (repulsion)

- ▶ if $T_N^3 \ll N$ infinitely many interfaces: $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?




Under the simple random walk law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

Under the polymer measure $\mathbf{P}_{N,\delta}^T$ with $\delta < 0$ $\hat{\tau}_1^T \approx T^3$ (repulsion)

- ▶ if $T_N^3 \ll N$ **infinitely many** interfaces: $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if $T_N^3 \gg N$ **confinement**: $|S_i| < T_N$ for all $i \leq N$:

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?




Under the simple random walk law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

Under the polymer measure $\mathbf{P}_{N,\delta}^T$ with $\delta < 0$ $\hat{\tau}_1^T \approx T^3$ (repulsion)

- ▶ if $T_N^3 \ll N$ **infinitely many** interfaces: $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if $T_N^3 \gg N$ **confinement**: $|S_i| < T_N$ for all $i \leq N$:
 - $S_N \approx T_N$ (if $T_N \leq \sqrt{N}$)

An heuristic view: the repulsive case $\delta < 0$

How long is the time $\hat{\tau}_1^T$ to reach level $\pm T$?



Under the simple random walk law $\hat{\tau}_1^T \approx T^2$ (diffusivity)

Under the polymer measure $\mathbf{P}_{N,\delta}^T$ with $\delta < 0$ $\hat{\tau}_1^T \approx T^3$ (repulsion)

- ▶ if $T_N^3 \ll N$ **infinitely many** interfaces: $S_N \approx T_N \sqrt{\frac{N}{T_N^3}} = \sqrt{\frac{N}{T_N}}$
- ▶ if $T_N^3 \gg N$ **confinement**: $|S_i| < T_N$ for all $i \leq N$:
 - $S_N \approx T_N$ (if $T_N \leq \sqrt{N}$)
 - $S_N \approx \sqrt{N}$ (if $T_N \gg \sqrt{N}$)

A renewal theory approach

Let $\tau_1^T, \tau_2^T, \tau_3^T \dots$ be the points at which S_n visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

A renewal theory approach

Let $\tau_1^T, \tau_2^T, \tau_3^T \dots$ be the points at which S_n visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

Under the **simple random walk** law $\{\tau_n^T\}_{n \in \mathbb{N}}$ is a classical **renewal process** with explicit law

$$q_T(n) := P(\tau_1^T = n) \approx \frac{c}{n^{3/2}} e^{-\frac{\pi^2}{2T^2} n}$$

A renewal theory approach

Let $\tau_1^T, \tau_2^T, \tau_3^T \dots$ be the points at which S_n visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

Under the **simple random walk** law $\{\tau_n^T\}_{n \in \mathbb{N}}$ is a classical **renewal process** with explicit law

$$q_T(n) := P(\tau_1^T = n) \approx \frac{c}{n^{3/2}} e^{-\frac{\pi^2}{2T^2} n}$$

Under the **polymer measure** $\mathbf{P}_{N,\delta}^T$ the process $\{\tau_n^T\}_{n \in \mathbb{N}}$ is not even time-homogeneous ...

A renewal theory approach

Let $\tau_1^T, \tau_2^T, \tau_3^T \dots$ be the points at which S_n visits an interface

$$\tau_{k+1}^T := \inf \{n > \tau_k^T : S_n - S_{\tau_k^T} \in \{-T, 0, T\}\} \quad (T \text{ is fixed})$$

Under the **simple random walk** law $\{\tau_n^T\}_{n \in \mathbb{N}}$ is a classical **renewal process** with explicit law

$$q_T(n) := P(\tau_1^T = n) \approx \frac{c}{n^{3/2}} e^{-\frac{\pi^2}{2T^2}n}$$

Under the **polymer measure** $\mathbf{P}_{N,\delta}^T$ the process $\{\tau_n^T\}_{n \in \mathbb{N}}$ is not even time-homogeneous ... however for large N it is nearly a **renewal process** with a different law $\mathcal{P}_{\delta,T}$

$$K_{\delta,T}(n) := \mathcal{P}_{\delta,T}(\tau_1^T = n) = e^\delta P(\tau_1^T = n) e^{-\phi(\delta,T)n}$$

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ does not depend explicitly on N .

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ does not depend explicitly on N .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ does not depend explicitly on N .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

1. Study $\tau^{T_N} \cap [0, N]$ under \mathcal{P}_{δ,T_N}

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ does not depend explicitly on N .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

1. Study $\tau^{T_N} \cap [0, N]$ under \mathcal{P}_{δ,T_N}
2. Transfer the results to $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$ that is to $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$ (hard part)

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ does not depend explicitly on N .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

1. Study $\tau^{T_N} \cap [0, N]$ under \mathcal{P}_{δ,T_N}
2. Transfer the results to $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$ that is to $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$ (hard part)
3. Remove the conditioning on $\{N \in \tau^{T_N}\}$

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ **does not depend explicitly on N** .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

1. Study $\tau^{T_N} \cap [0, N]$ under \mathcal{P}_{δ,T_N}
2. Transfer the results to $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$ that is to $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$ (**hard part**)
3. Remove the conditioning on $\{N \in \tau^{T_N}\}$

► Good estimates on $q_T(n)$ and on the free energy $\phi(\delta, T)$

Strategy of the proof

More precisely, the law of $\tau^T \cap [0, N] = \{\tau_1^T, \dots, \tau_{L_N}^T\}$ is the same

under $\mathbf{P}_{N,\delta}^T(\cdot \mid N \in \tau^T)$ and $\mathcal{P}_{\delta,T}(\cdot \mid N \in \tau^T)$

Important feature: $\mathcal{P}_{\delta,T}$ **does not depend explicitly on N** .

However we want to study $\mathbf{P}_{N,\delta}^{T_N}$ with T_N varying with N

1. Study $\tau^{T_N} \cap [0, N]$ under \mathcal{P}_{δ,T_N}
2. Transfer the results to $\mathcal{P}_{\delta,T_N}(\cdot \mid N \in \tau^{T_N})$ that is to $\mathbf{P}_{N,\delta}^{T_N}(\cdot \mid N \in \tau^{T_N})$ (**hard part**)
3. Remove the conditioning on $\{N \in \tau^{T_N}\}$

- ▶ Good estimates on $q_T(n)$ and on the free energy $\phi(\delta, T)$
- ▶ **Uniform renewal theorems**

Thanks.