

Young and Rough Equations in Infinite Dimension

Davide Addona

University of Parma

"The Mathematics of Subjective Probability"
Topics in Economics, Statistics and Probability

Milano, 8th-10th September, 2025

Special session "Stochastic Analysis"

Joint works with Luca Lorenzi (Parma) and Gianmario Tessitore (Bicocca)



The equation

We consider the following differential equation in a Banach space E

$$dY(t) = AY(t)dt + \sigma(Y(t))dX(t), \quad t \in (0, 1], \quad Y(0) = y_0 \in E \quad (\text{RDE})$$



The equation

We consider the following differential equation in a Banach space E

$$dY(t) = AY(t)dt + \sigma(Y(t))dX(t), \quad t \in (0, 1], \quad Y(0) = y_0 \in E \quad (\text{RDE})$$

- ★ $A : D(A) \subseteq E \rightarrow E$ is the **generator** of a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$
- ★ $\sigma : E \rightarrow E$ is a sufficiently smooth function
- ★ $X : [0, 1] \rightarrow \mathbb{R}$ is a α -Hölder continuous function for some $\alpha \in (\frac{1}{3}, 1)$



The equation

We consider the following differential equation in a Banach space E

$$dY(t) = AY(t)dt + \sigma(Y(t))dX(t), \quad t \in (0, 1], \quad Y(0) = y_0 \in E \quad (\text{RDE})$$

- ★ $A : D(A) \subseteq E \rightarrow E$ is the **generator** of a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$
- ★ $\sigma : E \rightarrow E$ is a sufficiently smooth function
- ★ $X : [0, 1] \rightarrow \mathbb{R}$ is a **α -Hölder** continuous function for some $\alpha \in (\frac{1}{3}, 1)$

GOALS: existence, uniqueness, regularity properties for the mild solution Y to (RDE)



The equation

We consider the following differential equation in a Banach space E

$$dY(t) = AY(t)dt + \sigma(Y(t))dX(t), \quad t \in (0, 1], \quad Y(0) = y_0 \in E \quad (\text{RDE})$$

★ $A : D(A) \subseteq E \rightarrow E$ is the generator of a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$

★ $\sigma : E \rightarrow E$ is a sufficiently smooth function

★ $X : [0, 1] \rightarrow \mathbb{R}$ is a α -Hölder continuous function for some $\alpha \in (\frac{1}{3}, 1)$

GOALS: existence, uniqueness, regularity properties for the mild solution Y to (RDE) formally given by

$$Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s), \quad t \in [0, 1] \quad (\text{MILD})$$





Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$



Motivation & Problems

Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$

W Brownian Motion \Rightarrow integral formulation of the SDE: $Z(t) = \xi + \int_0^t \sigma(Z(s))dW(s)$



Motivation & Problems

Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$

W Brownian Motion \Rightarrow integral formulation of the SDE: $Z(t) = \xi + \int_0^t \sigma(Z(s))dW(s)$

How can we define the stochastic integral $\int_0^t \sigma(Z(s))dW(s)$?



Motivation & Problems

Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$

W Brownian Motion \Rightarrow integral formulation of the SDE: $Z(t) = \xi + \int_0^t \sigma(Z(s))dW(s)$

How can we define the stochastic integral $\int_0^t \sigma(Z(s))dW(s)$?

- probabilistic definition of $\int_0^t \sigma(Z(s))dW(s)$: Itô-Stratonovich



Motivation & Problems

Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$

W Brownian Motion \Rightarrow integral formulation of the SDE: $Z(t) = \xi + \int_0^t \sigma(Z(s))dW(s)$

How can we define the stochastic integral $\int_0^t \sigma(Z(s))dW(s)$?

- probabilistic definition of $\int_0^t \sigma(Z(s))dW(s)$: Itô-Stratonovich
- pathwise definition of $\int_0^t \sigma(Z_\omega(s))dW_\omega(s)$, $\omega \in \Omega$???



Motivation & Problems

Consider the SDE: $Z(t) = \sigma(Z(t))dW(t)$, $t \in (0, 1]$, $Z(0) = \xi \in \mathbb{R}$

W Brownian Motion \Rightarrow integral formulation of the SDE: $Z(t) = \xi + \int_0^t \sigma(Z(s))dW(s)$

How can we define the stochastic integral $\int_0^t \sigma(Z(s))dW(s)$?

- probabilistic definition of $\int_0^t \sigma(Z(s))dW(s)$: Itô-Stratonovich
- pathwise definition of $\int_0^t \sigma(Z_\omega(s))dW_\omega(s)$, $\omega \in \Omega$???

\mathbb{P} -a.s. $\omega \in \Omega$:

- W_ω has not finite 1-variation
- W_ω (and $\sigma(Z_\omega)$) α -Hölder continuous for every $\alpha < \frac{1}{2}$
- W_ω has finite p -variation for every $p > 2$



We set $I(f, g) := \int_0^1 f(t)\dot{g}(t)dt$, $f, g \in C^1([0, 1])$



We set $I(f, g) := \int_0^1 f(t)\dot{g}(t)dt$, $f, g \in C^1([0, 1])$

Negative result (Lyons '91): \nexists a Banach space $\mathcal{B} \subseteq C([0, 1])$ such that

- $C^\alpha \subseteq \mathcal{B}$ for every $\alpha < \frac{1}{2}$
- I extends to a continuous map $I : \mathcal{B} \times \mathcal{B} \rightarrow C([0, 1])$



We set $I(f, g) := \int_0^{\cdot} f(t) \dot{g}(t) dt$, $f, g \in C^1([0, 1])$

Negative result (Lyons '91): \nexists a Banach space $\mathcal{B} \subseteq C([0, 1])$ such that

- $C^\alpha \subseteq \mathcal{B}$ for every $\alpha < \frac{1}{2}$
- I extends to a continuous map $I : \mathcal{B} \times \mathcal{B} \rightarrow C([0, 1])$

Positive result (Young '36, Kondurar '37): if $\alpha + \beta > 1$, then

- I extends to a continuous map $I_Y : C^\alpha([0, 1]) \times C^\beta([0, 1]) \rightarrow C^\beta([0, 1])$
- $|I_Y(f, g)(s, t) - f(s)(g(t) - g(s))| \leq C_{\alpha+\beta} [f]_\alpha [g]_\beta |t - s|^{\alpha+\beta}$
- $I_Y(f, g)(s, t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} f(u)(g(v) - g(u))$, $0 \leq s < t \leq 1$, P partition of $[s, t]$



We set $I(f, g) := \int_0^1 f(t)\dot{g}(t)dt$, $f, g \in C^1([0, 1])$

Negative result (Lyons '91): \nexists a Banach space $\mathcal{B} \subseteq C([0, 1])$ such that

- $C^\alpha \subseteq \mathcal{B}$ for every $\alpha < \frac{1}{2}$
- I extends to a continuous map $I : \mathcal{B} \times \mathcal{B} \rightarrow C([0, 1])$

Positive result (Young '36, Kondurar '37): if $\alpha + \beta > 1$, then

- I extends to a continuous map $I_Y : C^\alpha([0, 1]) \times C^\beta([0, 1]) \rightarrow C^\beta([0, 1])$
- $|I_Y(f, g)(s, t) - f(s)(g(t) - g(s))| \leq C_{\alpha+\beta} [f]_\alpha [g]_\beta |t - s|^{\alpha+\beta}$
- $I_Y(f, g)(s, t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} f(u)(g(v) - g(u))$, $0 \leq s < t \leq 1$, P partition of $[s, t]$

Consequences: let $\mathcal{B} = C^\alpha([0, 1])$, $\alpha \in (\frac{1}{2}, 1)$.

- I extends to a continuous map $I_Y : \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$: **Young Integral**
- What about the case $\alpha \leq \frac{1}{2}$?



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1$, $F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$

$$\left[F(X(r)) - F(X(s)) = F'(X(s))(X(r) - X(s)) + \frac{1}{2}F''(\xi)(X(r) - X(s))^2 \right]$$



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$

$$\left[F(X(r)) - F(X(s)) = F'(X(s))(X(r) - X(s)) + \frac{1}{2}F''(\xi)(X(r) - X(s))^2 \right]$$

$$= F'(X(s)) \int_s^t (X(r) - X(s))dX(r) + \text{reminder}$$



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$

$$\left[F(X(r)) - F(X(s)) = F'(X(s))(X(r) - X(s)) + \frac{1}{2}F''(\xi)(X(r) - X(s))^2 \right]$$

$$= F'(X(s)) \int_s^t (X(r) - X(s))dX(r) + \text{reminder}$$

$$\int_s^t (X(r) - X(s))dX(r) = \int_s^t X(r)dX(r) - X(s)(X(t) - X(s))$$



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$

$$\left[F(X(r)) - F(X(s)) = F'(X(s))(X(r) - X(s)) + \frac{1}{2}F''(\xi)(X(r) - X(s))^2 \right]$$

$$= F'(X(s)) \int_s^t (X(r) - X(s))dX(r) + \text{reminder}$$

$$\int_s^t (X(r) - X(s))dX(r) = \int_s^t X(r)dX(r) - X(s)(X(t) - X(s))$$

$$\alpha \in \left(\frac{1}{2}, 1\right) \Rightarrow \int_s^t X(r)dX(r) = I_Y(X, X)(s, t) \text{ Young Integral}$$



Integration of 1-forms (Folmer '81, Lyons '98)

Assume $X \in C^1, F \in C^2$. Then

$$\int_s^t F(X(r))dX(r) - F(X(s))(X(t) - X(s)) = \int_s^t (F(X(r)) - F(X(s)))dX(r)$$

$$\left[F(X(r)) - F(X(s)) = F'(X(s))(X(r) - X(s)) + \frac{1}{2}F''(\xi)(X(r) - X(s))^2 \right]$$

$$= F'(X(s)) \int_s^t (X(r) - X(s))dX(r) + \text{reminder}$$

$$\int_s^t (X(r) - X(s))dX(r) = \int_s^t X(r)dX(r) - X(s)(X(t) - X(s))$$

$$\alpha \in \left(\frac{1}{2}, 1\right) \Rightarrow \int_s^t X(r)dX(r) = I_Y(X, X)(s, t) \text{ Young Integral}$$

$$\alpha \leq \frac{1}{2} \Rightarrow \int_s^t X(r)dX(r) \text{ defines } \int_s^t F(X(r))dX(r)$$



Hölder Continuous Rough Paths (Lyons '98)



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$

$$\bullet X : [0, 1] \rightarrow \mathbb{R}, \quad \sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty \quad (X \in C^\alpha([0, 1]; \mathbb{R}))$$



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$

- $X : [0, 1] \rightarrow \mathbb{R}$, $\sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty$ ($X \in C^\alpha([0, 1]; \mathbb{R})$)
- $\mathbb{X} : [0, 1]_{<}^2 \rightarrow \mathbb{R}$, $\|\mathbb{X}\|_{2\alpha} := \sup_{0 \leq s < t \leq 1} \frac{|\mathbb{X}(s, t)|}{|t - s|^{2\alpha}} < \infty$ ($\mathbb{X} \in C^{2\alpha}([0, 1]_{<}^2; \mathbb{R})$)



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$

- $X : [0, 1] \rightarrow \mathbb{R}$, $\sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty$ ($X \in C^\alpha([0, 1]; \mathbb{R})$)
- $\mathbb{X} : [0, 1]_{<}^2 \rightarrow \mathbb{R}$, $\|\mathbb{X}\|_{2\alpha} := \sup_{0 \leq s < t \leq 1} \frac{|\mathbb{X}(s, t)|}{|t - s|^{2\alpha}} < \infty$ ($\mathbb{X} \in C^{2\alpha}([0, 1]_{<}^2; \mathbb{R})$)
- $\mathbb{X}(s, u) - \mathbb{X}(s, t) - \mathbb{X}(t, u) = (X(t) - X(s))(X(u) - X(t))$ (Chen's Relation)



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$

- $X : [0, 1] \rightarrow \mathbb{R}$, $\sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty$ ($X \in C^\alpha([0, 1]; \mathbb{R})$)
- $\mathbb{X} : [0, 1]_{<}^2 \rightarrow \mathbb{R}$, $\|\mathbb{X}\|_{2\alpha} := \sup_{0 \leq s < t \leq 1} \frac{|\mathbb{X}(s, t)|}{|t - s|^{2\alpha}} < \infty$ ($\mathbb{X} \in C^{2\alpha}([0, 1]_{<}^2; \mathbb{R})$)
- $\mathbb{X}(s, u) - \mathbb{X}(s, t) - \mathbb{X}(t, u) = (X(t) - X(s))(X(u) - X(t))$ (Chen's Relation)

What is the meaning of \mathbb{X} ?



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of a α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s < t\})$

- $X : [0, 1] \rightarrow \mathbb{R}$, $\sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty$ ($X \in C^\alpha([0, 1]; \mathbb{R})$)
- $\mathbb{X} : [0, 1]_{<}^2 \rightarrow \mathbb{R}$, $\|\mathbb{X}\|_{2\alpha} := \sup_{0 \leq s < t \leq 1} \frac{|\mathbb{X}(s, t)|}{|t - s|^{2\alpha}} < \infty$ ($\mathbb{X} \in C^{2\alpha}([0, 1]_{<}^2; \mathbb{R})$)
- $\mathbb{X}(s, u) - \mathbb{X}(s, t) - \mathbb{X}(t, u) = (X(t) - X(s))(X(u) - X(t))$ (Chen's Relation)

What is the meaning of \mathbb{X} ?

$\alpha > \frac{1}{2} \Rightarrow \mathbb{X}(s, t) := \int_s^t (X(r) - X(s))dX(r)$, $0 \leq s \leq t \leq 1$: Young Integral



Hölder Continuous Rough Paths (Lyons '98)

Given $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, the space $\mathcal{C}^\alpha([0, 1]; \mathbb{R})$ of α -Hölder rough paths is given by the couples $\mathbf{X} = (X; \mathbb{X})$ such that $([0, 1]_{<}^2 := \{(s, t) \in [0, 1]^2 : s \leq t\})$

- $X : [0, 1] \rightarrow \mathbb{R}$, $\sup_{0 \leq s < t \leq 1} \frac{|X(t) - X(s)|}{|t - s|^\alpha} < \infty$ ($X \in C^\alpha([0, 1]; \mathbb{R})$)
- $\mathbb{X} : [0, 1]_{<}^2 \rightarrow \mathbb{R}$, $\|\mathbb{X}\|_{2\alpha} := \sup_{0 \leq s < t \leq 1} \frac{|\mathbb{X}(s, t)|}{|t - s|^{2\alpha}} < \infty$ ($\mathbb{X} \in C^{2\alpha}([0, 1]_{<}^2; \mathbb{R})$)
- $\mathbb{X}(s, u) - \mathbb{X}(s, t) - \mathbb{X}(t, u) = (X(t) - X(s))(X(u) - X(t))$ (Chen's Relation)

What is the meaning of \mathbb{X} ?

$\alpha > \frac{1}{2} \Rightarrow \mathbb{X}(s, t) := \int_s^t (X(r) - X(s))dX(r)$, $0 \leq s \leq t \leq 1$: Young Integral

$\alpha \in (\frac{1}{3}, \frac{1}{2}] \Rightarrow \mathbb{X}(s, t)$ possible choice for " $\int_s^t (X(r) - X(s))dX(r)$ ": infinitely many!





Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$



Rough Integration

$X \in C^1([0, 1])$, $F \in C_b^2(\mathbb{R})$, $0 \leq s < t \leq 1$,

Integral: $\int_s^t F(X(r))dX(r)$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) \pm F(X(s))(X(t) - X(s))$$



Rough Integration

$X \in C^1([0, 1])$, $F \in C_b^2(\mathbb{R})$, $0 \leq s < t \leq 1$,

Integral: $\int_s^t F(X(r))dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s)))dX(r)$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s))) dX(r)$$

$$\text{Taylor: } F(X(t)) - F(X(s)) = F'(X(s))(X(t) - X(s)) + \frac{1}{2}F''(\xi)(X(t) - X(s))^2$$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s))) dX(r)$$

$$\text{Taylor: } F(X(t)) - F(X(s)) = F'(X(s))(X(t) - X(s)) + \frac{1}{2}F''(\xi)(X(t) - X(s))^2$$

$$\text{Reminder: } R(s, t) = O(|X(t) - X(s)|^2) = F'(X(s))\mathbb{X}(s, t) + O(|X(t) - X(s)|^3)$$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s))) dX(r)$$

$$\text{Taylor: } F(X(t)) - F(X(s)) = F'(X(s))(X(t) - X(s)) + \frac{1}{2} F''(\xi)(X(t) - X(s))^2$$

$$\text{Reminder: } R(s, t) = O(|X(t) - X(s)|^2) = F'(X(s))\mathbb{X}(s, t) + O(|X(t) - X(s)|^3)$$

$$X \in C^\alpha([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1:$$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s))) dX(r)$$

$$\text{Taylor: } F(X(t)) - F(X(s)) = F'(X(s))(X(t) - X(s)) + \frac{1}{2}F''(\xi)(X(t) - X(s))^2$$

$$\text{Reminder: } R(s, t) = O(|X(t) - X(s)|^2) = F'(X(s))\mathbb{X}(s, t) + O(|X(t) - X(s)|^3)$$

$$X \in C^\alpha([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1:$$

★ If $\alpha > \frac{1}{2}$ then $R(s, t) = O(|t - s|^{2\alpha})$ and the limit

$$\lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} F(X(u))(X(v) - X(u)) = I_Y(F(X), X) \text{ exists (Young Integral)}$$



Rough Integration

$$X \in C^1([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1,$$

$$\text{Integral: } \int_s^t F(X(r)) dX(r) = F(X(s))(X(t) - X(s)) + \int_s^t (F(X(r)) - F(X(s))) dX(r)$$

$$\text{Taylor: } F(X(t)) - F(X(s)) = F'(X(s))(X(t) - X(s)) + \frac{1}{2} F''(\xi)(X(t) - X(s))^2$$

$$\text{Reminder: } R(s, t) = O(|X(t) - X(s)|^2) = F'(X(s))\mathbb{X}(s, t) + O(|X(t) - X(s)|^3)$$

$$X \in C^\alpha([0, 1]), F \in C_b^2(\mathbb{R}), 0 \leq s < t \leq 1:$$

★ If $\alpha > \frac{1}{2}$ then $R(s, t) = O(|t - s|^{2\alpha})$ and the limit

$$\lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} F(X(u))(X(v) - X(u)) = I_Y(F(X), X) \text{ exists (Young Integral)}$$

★ If $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ then $R(s, t) = F'(X(s))\mathbb{X}(s, t) + O(|t - s|^{3\alpha})$ and the limit exists:

$$\lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} [F(X(u))(X(v) - X(u)) + F'(X(u))\mathbb{X}(u, v)]$$

Lyons '98: $\mathbf{X} = (X, \mathbb{X})$ α -Hölder rough path, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $F \in C_b^2(\mathbb{R})$. The rough integral

$$I_R(F(\mathbf{X}), \mathbf{X})(s, t) = \lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} [F(X(u))(X(v) - X(u)) + F'(X(u))\mathbb{X}(u, v)]$$

exists for every $0 \leq s < t \leq 1$.



Lyons '98: $\mathbf{X} = (X, \mathbb{X})$ α -Hölder rough path, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $F \in C_b^2(\mathbb{R})$. The rough integral

$$I_R(F(\mathbf{X}), \mathbf{X})(s, t) = \lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} [F(X(u))(X(v) - X(u)) + F'(X(u))\mathbb{X}(u, v)]$$

exists for every $0 \leq s < t \leq 1$. Moreover, $I_R(F(\mathbf{X}), \mathbf{X})$ satisfies

$$|I_R(F(\mathbf{X}), \mathbf{X})(s, t) - F(X(s))(X(t) - X(s)) - F'(X(s))\mathbb{X}(s, t)| \leq C(F, \mathbf{X})|t - s|^{3\alpha}$$



Lyons '98: $\mathbf{X} = (X, \mathbb{X})$ α -Hölder rough path, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $F \in C_b^2(\mathbb{R})$. The rough integral

$$I_R(F(\mathbf{X}), \mathbf{X})(s, t) = \lim_{|P(s, t)| \rightarrow 0} \sum_{[u, v] \in P(s, t)} [F(X(u))(X(v) - X(u)) + F'(X(u))\mathbb{X}(u, v)]$$

exists for every $0 \leq s < t \leq 1$. Moreover, $I_R(F(\mathbf{X}), \mathbf{X})$ satisfies

$$|I_R(F(\mathbf{X}), \mathbf{X})(s, t) - F(X(s))(X(t) - X(s)) - F'(X(s))\mathbb{X}(s, t)| \leq C(F, \mathbf{X})|t - s|^{3\alpha}$$

Lyons solves a rough differential equation of the form

$$dY(t) = \sigma(Y(t))d\mathbf{X}(t), \quad Y(0) = y_0 \text{ (i.e., } Y(t) = y_0 + I_R(\sigma(Y), \mathbf{X})(0, t))$$

but the concept of solution and the techniques exploited are complicated and involved. Another (but equivalent) definition of solution and proof of the existence and uniqueness of solutions can be obtained by means of *Sewing Lemma* and *Controlled Rough Paths*, introduced in '04 and generalized to the mild situation in '10 by Gubinelli.



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that
$$\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$

★ **Chen's Relation**: $(\delta_2 \mathbb{X})(s, t, u) = (\delta_1 X)(s, t)(\delta_1 X)(t, u)$, $(s, t, u) \in [0, 1]_{<}^3$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$

★ **Chen's Relation**: $(\delta_2 \mathbb{X})(s, t, u) = (\delta_1 X)(s, t)(\delta_1 X)(t, u)$, $(s, t, u) \in [0, 1]_{<}^3$

$$C_2^{\alpha, \beta}([0, 1]) := \{A \in C^\alpha([0, 1]_{<}^2) : \delta_2 A \in C^\beta([0, 1]_{<}^3)\}$$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$

★ **Chen's Relation**: $(\delta_2 \mathbb{X})(s, t, u) = (\delta_1 X)(s, t)(\delta_1 X)(t, u)$, $(s, t, u) \in [0, 1]_{<}^3$

$$C_2^{\alpha, \beta}([0, 1]) := \{A \in C^\alpha([0, 1]_{<}^2) : \delta_2 A \in C^\beta([0, 1]_{<}^3)\}$$

Sewing Lemma (Gubinelli '04): Let $0 < \alpha \leq 1 < \beta$. There exists a unique continuous linear map $I : C_2^{\alpha, \beta}([0, 1]) \rightarrow C^\alpha([0, 1])$ such that $IA(0) = 0$ and

- $|(\delta_1 IA)(s, t) - A(s, t)| \leq C(\beta, A)|t - s|^\beta$, $(s, t) \in [0, 1]_{<}^2$
- $IA(t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} A(u, v)$, P partition of $[0, t]$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$

★ **Chen's Relation**: $(\delta_2 \mathbb{X})(s, t, u) = (\delta_1 X)(s, t)(\delta_1 X)(t, u)$, $(s, t, u) \in [0, 1]_{<}^3$

$$C_2^{\alpha, \beta}([0, 1]) := \{A \in C^\alpha([0, 1]_{<}^2) : \delta_2 A \in C^\beta([0, 1]_{<}^3)\}$$

Sewing Lemma (Gubinelli '04): Let $0 < \alpha \leq 1 < \beta$. There exists a unique continuous linear map $I : C_2^{\alpha, \beta}([0, 1]) \rightarrow C^\alpha([0, 1])$ such that $IA(0) = 0$ and

$$|(\delta_1 IA)(s, t) - A(s, t)| \leq C(\beta, A)|t - s|^\beta, \quad (s, t) \in [0, 1]_{<}^2$$

$$IA(t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} A(u, v), \quad P \text{ partition of } [0, t]$$

★ **Example**: $A(s, t) = f(s)(\delta_1 g)(s, t) \Rightarrow (\delta_2 A)(s, t, u) = -(\delta_1 f)(s, t)(\delta_1 g)(t, u)$



Sewing Lemma

Let $N \in \mathbb{N}$, $\alpha, \beta \geq 0$, $[0, 1]_{<}^N := \{(t_1, \dots, t_N) \in [0, 1]^N : 0 \leq t_1 \leq \dots \leq t_N \leq 1\}$

★ $C^\beta([0, 1]_{<}^N)$: functions $A : [0, 1]_{<}^N \rightarrow \mathbb{R}$ such that $\sup_{0 \leq t_1 < \dots < t_N \leq 1} \frac{|A(t_1, \dots, t_N)|}{|t_N - t_1|^\beta} < \infty$

★ $\delta_N : C^0([0, 1]_{<}^N) \rightarrow C^0([0, 1]_{<}^{N+1})$ defined as (for $N = 1, 2$, $(s, t, u) \in [0, 1]_{<}^3$)

$$(\delta_1 A)(s, t) = A(t) - A(s), \quad (\delta_2 A)(s, t, u) = A(s, u) - A(s, t) - A(t, u)$$

★ **Chen's Relation**: $(\delta_2 \mathbb{X})(s, t, u) = (\delta_1 X)(s, t)(\delta_1 X)(t, u)$, $(s, t, u) \in [0, 1]_{<}^3$

$$C_2^{\alpha, \beta}([0, 1]) := \{A \in C^\alpha([0, 1]_{<}^2) : \delta_2 A \in C^\beta([0, 1]_{<}^3)\}$$

Sewing Lemma (Gubinelli '04): Let $0 < \alpha \leq 1 < \beta$. There exists a unique continuous linear map $I : C_2^{\alpha, \beta}([0, 1]) \rightarrow C^\alpha([0, 1])$ such that $IA(0) = 0$ and

$$|(\delta_1 IA)(s, t) - A(s, t)| \leq C(\beta, A)|t - s|^\beta, \quad (s, t) \in [0, 1]_{<}^2$$

$$IA(t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} A(u, v), \quad P \text{ partition of } [0, t]$$

★ **Example**: $A(s, t) = f(s)(\delta_1 g)(s, t) \Rightarrow (\delta_2 A)(s, t, u) = -(\delta_1 f)(s, t)(\delta_1 g)(t, u)$

It $f \in C^\alpha, g \in C^\beta, \alpha + \beta > 1$, then $IA = I_\gamma(f, g)$





Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$



Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$

Example: If $A(s, t) = F(X(s))(\delta_1 X)(s, t) + F'(X(s))\mathbb{X}(s, t)$, then

$$(\delta_2 A)(s, t, u) = -R(s, t)(\delta_1 X)(t, u) - (\delta_1 F'(X))(s, t)\mathbb{X}(t, u)$$

If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, then $R(s, t) = O(|t - s|^{2\alpha})$ and $IA = I_R(F(X), \mathbf{X})$



Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$

Example: If $A(s, t) = F(X(s))(\delta_1 X)(s, t) + F'(X(s))\mathbb{X}(s, t)$, then

$$(\delta_2 A)(s, t, u) = -R(s, t)(\delta_1 X)(t, u) - (\delta_1 F'(X))(s, t)\mathbb{X}(t, u)$$

If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, then $R(s, t) = O(|t - s|^{2\alpha})$ and $IA = I_R(F(X), \mathbf{X})$

How can we extend (beyond $F(X)$) the class of “functions under the integral sign”?



Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$

Example: If $A(s, t) = F(X(s))(\delta_1 X)(s, t) + F'(X(s))\mathbb{X}(s, t)$, then

$$(\delta_2 A)(s, t, u) = -R(s, t)(\delta_1 X)(t, u) - (\delta_1 F'(X))(s, t)\mathbb{X}(t, u)$$

If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, then $R(s, t) = O(|t - s|^{2\alpha})$ and $IA = I_R(F(X), \mathbf{X})$

How can we extend (beyond $F(X)$) the class of “functions under the integral sign”?

Replacing $F(X)$ and $F'(X)$ with objects with the same properties!



Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$

Example: If $A(s, t) = F(X(s))(\delta_1 X)(s, t) + F'(X(s))\mathbb{X}(s, t)$, then

$$(\delta_2 A)(s, t, u) = -R(s, t)(\delta_1 X)(t, u) - (\delta_1 F'(X))(s, t)\mathbb{X}(t, u)$$

If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, then $R(s, t) = O(|t - s|^{2\alpha})$ and $IA = I_R(F(X), \mathbb{X})$

How can we extend (beyond $F(X)$) the class of “functions under the integral sign”?

Replacing $F(X)$ and $F'(X)$ with objects with the same properties!

Gubinelli '04: $\mathcal{D}_X^{2\alpha}(0, 1)$ is the space of couples $(Y, Y') \in C^\alpha([0, 1]) \times C^\alpha([0, 1])$:

$$R^Y(s, t) := (\delta_1 Y)(s, t) - Y'(s)(\delta_1 X)(s, t) = O(|t - s|^{2\alpha})$$

Y is controlled by X if $\exists Y'$ such that $(Y, Y') \in \mathcal{D}_X^{2\alpha}$, Y' Gubinelli derivative of Y



Controlled Rough Paths

For $F \in C_b^2$, We set $R(s, t) := (\delta_1 F(X))(s, t) - F'(X(s))(\delta_1 X)(s, t) \sim (X(t) - X(s))^2$

Example: If $A(s, t) = F(X(s))(\delta_1 X)(s, t) + F'(X(s))\mathbb{X}(s, t)$, then

$$(\delta_2 A)(s, t, u) = -R(s, t)(\delta_1 X)(t, u) - (\delta_1 F'(X))(s, t)\mathbb{X}(t, u)$$

If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, then $R(s, t) = O(|t - s|^{2\alpha})$ and $IA = I_R(F(X), \mathbf{X})$

How can we extend (beyond $F(X)$) the class of “functions under the integral sign”?

Replacing $F(X)$ and $F'(X)$ with objects with the same properties!

Gubinelli '04: $\mathcal{D}_X^{2\alpha}(0, 1)$ is the space of couples $(Y, Y') \in C^\alpha([0, 1]) \times C^\alpha([0, 1])$:

$$R^Y(s, t) := (\delta_1 Y)(s, t) - Y'(s)(\delta_1 X)(s, t) = O(|t - s|^{2\alpha})$$

Y is controlled by X if $\exists Y'$ such that $(Y, Y') \in \mathcal{D}_X^{2\alpha}$, Y' Gubinelli derivative of Y

Gubinelli '04: If $(X, \mathbb{X}) \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ and $(Y, Y') \in \mathcal{D}_X^{2\alpha}$, then

$$I_R(Y, \mathbf{X})(s, t) = \lim_{|P| \rightarrow 0} \sum_{[u, v] \in P} (Y(u)(\delta_1 X)(u, v) + Y'(u)\mathbb{X}(u, v))$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\int_s^t f(r)dX(r) = \int_s^t (f(r) \pm f(s))dX(r)$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned} \int_s^t f(r)dX(r) &= \int_s^t (f(r) \pm f(s))dX(r) \\ &= f(s)(X(t) - X(s)) + \int_s^t (f(r) - f(s))dX(r) \end{aligned}$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned} \int_s^t e^{(t-r)A}f(r)dX(r) &= \int_s^t e^{(t-r)A}(f(r) \pm f(s))dX(r) \\ &= f(s)(X(t) - X(s)) + \int_s^t e^{(t-r)A}(f(r) - f(s))dX(r) \end{aligned}$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned} \int_s^t e^{(t-r)A}f(r)dX(r) &= \int_s^t e^{(t-r)A}(f(r) \pm e^{(r-s)A}f(s))dX(r) \\ &= f(s)(X(t) - X(s)) + \int_s^t e^{(t-r)A}(f(r) - e^{(r-s)A}f(s))dX(r) \end{aligned}$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned}\int_s^t e^{(t-r)A}f(r)dX(r) &= \int_s^t e^{(t-r)A}(f(r) \pm e^{(r-s)A}f(s))dX(r) \\ &= e^{(t-s)A}f(s)(X(t) - X(s)) + \int_s^t e^{(t-r)A}(f(r) - e^{(r-s)A}f(s))dX(r)\end{aligned}$$



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned}\int_s^t e^{(t-r)A}f(r)dX(r) &= \int_s^t e^{(t-r)A}(f(r) \pm e^{(r-s)A}f(s))dX(r) \\ &= e^{(t-s)A}f(s)(X(t) - X(s)) + \int_s^t e^{(t-r)A}(f(r) - e^{(r-s)A}f(s))dX(r)\end{aligned}$$

★ We have new increments $\delta_{S,N}$:



Mild Integration: the Young Case

$A : D(A) \subseteq E \rightarrow E$ generates a semigroup $(e^{tA}) \subseteq \mathcal{L}(E)$, $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$

The equation: $dY(t) = AY(t)dt + \sigma(Y(t))dX(t)$, $t \in (0, 1]$, $Y(0) = y_0 \in E$

The mild solution: $Y(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))dX(s)$, $t \in [0, 1]$

★ We aim at defining $\int_s^t e^{(t-r)A}f(r)dX(r)$, $X \in C^\alpha$, for a class of $f : [0, 1] \rightarrow E$:

$$\begin{aligned}\int_s^t e^{(t-r)A}f(r)dX(r) &= \int_s^t e^{(t-r)A}(f(r) \pm e^{(r-s)A}f(s))dX(r) \\ &= e^{(t-s)A}f(s)(X(t) - X(s)) + \int_s^t e^{(t-r)A}(f(r) - e^{(r-s)A}f(s))dX(r)\end{aligned}$$

★ We have new increments $\delta_{S,N}$:

$$(\delta_{S,1}B)(s, t) := B(t) - e^{(t-s)A}B(s), \quad (\delta_{S,2}B)(s, t, u) = B(s, u) - B(t, u) - e^{(u-t)A}B(s, t)$$



$$C_{S,2}^{\alpha,\beta}([0,1];E) := \{B \in C^\alpha([0,1]_{<}^2;E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3;E)\}$$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1} I_S B)(s, t) - B(s, t)| \leq C(\beta, B) |t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v)$, P partition of $[0, t]$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1} I_S B)(s, t) - B(s, t)| \leq C(\beta, B) |t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v)$, P partition of $[0, t]$

If $B(s, t) = e^{(t-s)A} f(s) (\delta_1 X)(s, t)$, then $(\delta_{S,2} B)(s, t, u) = -(\delta_{S,1} f)(s, t) (\delta_1 X)(t, u)$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1} I_S B)(s, t) - B(s, t)| \leq C(\beta, B) |t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v)$, P partition of $[0, t]$

If $B(s, t) = e^{(t-s)A} f(s) (\delta_1 X)(s, t)$, then $(\delta_{S,2} B)(s, t, u) = -(\delta_{S,1} f)(s, t) (\delta_1 X)(t, u)$

$I_S B(t) = \int_0^t e^{(t-s)A} f(s) dX(s)''$ when $\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$ with $\alpha + \eta > 1$

How to find a mild solution? Fixed point for $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ given by

$$\Gamma(f)(t) = e^{tA} y_0 + I_S(\sigma(f))(t), \quad t \in [0, 1],$$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1} I_S B)(s, t) - B(s, t)| \leq C(\beta, B) |t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v)$, P partition of $[0, t]$

If $B(s, t) = e^{(t-s)A} f(s) (\delta_1 X)(s, t)$, then $(\delta_{S,2} B)(s, t, u) = -(\delta_{S,1} f)(s, t) (\delta_1 X)(t, u)$

$I_S B(t) = \int_0^t e^{(t-s)A} f(s) dX(s)''$ when $\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$ with $\alpha + \eta > 1$

How to find a mild solution? Fixed point for $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ given by

$$\Gamma(f)(t) = e^{tA} y_0 + I_S(\sigma(f))(t), \quad t \in [0, 1], \quad \text{what is } \mathcal{B}???$$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1}I_S B)(s, t) - B(s, t)| \leq C(\beta, B)|t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v)$, P partition of $[0, t]$

If $B(s, t) = e^{(t-s)A} f(s) (\delta_1 X)(s, t)$, then $(\delta_{S,2} B)(s, t, u) = -(\delta_{S,1} f)(s, t) (\delta_1 X)(t, u)$

$I_S B(t) = \int_0^t e^{(t-s)A} f(s) dX(s)$ when $\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$ with $\alpha + \eta > 1$

How to find a mild solution? Fixed point for $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ given by

$$\Gamma(f)(t) = e^{tA} y_0 + I_S(\sigma(f))(t), \quad t \in [0, 1], \quad \text{what is } \mathcal{B}???$$

$\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$, $\sigma \in C_b^1(E; E)$, then

$$(\delta_{S,1}(\sigma(f)))(s, t) = \sigma(f(t)) - \sigma(e^{(t-s)A} f(s)) + (e^{(t-s)A} - Id)(\sigma(f))(s) + \dots \in C^\eta?$$



$$C_{S,2}^{\alpha,\beta}([0,1]; E) := \{B \in C^\alpha([0,1]_{<}^2; E) : \delta_{S,2}B \in C^\beta([0,1]_{<}^3; E)\}$$

Mild Sewing Lemma (Gubinelli-Tindel '10): If $0 < \alpha \leq 1 < \beta$, then there exists a unique continuous linear map $I_S : C_{S,2}^{\alpha,\beta}([0,1]; E) \rightarrow C^\alpha([0,1]; E)$ such that $I_S B(0) = 0$ and

- $|(\delta_{S,1}I_S B)(s, t) - B(s, t)| \leq C(\beta, B)|t - s|^\beta$
- $I_S B(t) = \lim_{|P| \rightarrow 0} \sum_{[u,v] \in P} e^{(t-v)A} B(u, v), P \text{ partition of } [0, t]$

If $B(s, t) = e^{(t-s)A} f(s) (\delta_1 X)(s, t)$, then $(\delta_{S,2} B)(s, t, u) = -(\delta_{S,1} f)(s, t) (\delta_1 X)(t, u)$

$I_S B(t) = \int_0^t e^{(t-s)A} f(s) dX(s)''$ when $\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$ with $\alpha + \eta > 1$

How to find a mild solution? Fixed point for $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ given by

$$\Gamma(f)(t) = e^{tA} y_0 + I_S(\sigma(f))(t), \quad t \in [0, 1], \quad \text{what is } \mathcal{B}???$$

$\delta_{S,1} f \in C^\eta([0,1]_{<}^2; E)$, $\sigma \in C_b^1(E; E)$, then

$$(\delta_{S,1}(\sigma(f)))(s, t) = \sigma(f(t)) - \sigma(e^{(t-s)A} f(s)) + (e^{(t-s)A} - Id)(\sigma(f))(s) + \dots \in C^\eta?$$

If (e^{tA}) is analytic, then $\|e^{rA} - Id\|_{\mathcal{L}(D_A(\eta, \infty), E)} \leq Cr^\eta, \eta \in [0, 1]$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq C t^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq C t^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq C t^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq C t^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1)$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq C t^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq C t^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1]$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$

Theorem (A-Lorenzi-Tessitore '22): Let $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta \in (1 - \alpha, \alpha)$, $y_0 \in E_\eta$ and σ smooth.



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq C t^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq C t^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1]$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$

Theorem (A-Lorenzi-Tessitore '22): Let $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta \in (1 - \alpha, \alpha)$, $y_0 \in E_\eta$ and σ smooth. Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq Ct^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq Ct^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1]$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$

Theorem (A-Lorenzi-Tessitore '22): Let $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta \in (1 - \alpha, \alpha)$, $y_0 \in E_\eta$ and σ smooth. Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

Further, $Y(t) \in D(A)$ for every $t \in (0, 1]$, it satisfies the integral representation

$$Y(t) = y_0 + \int_0^t AY(s) ds + \int_0^t \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq Ct^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq Ct^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1]$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$

Theorem (A-Lorenzi-Tessitore '22): Let $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta \in (1 - \alpha, \alpha)$, $y_0 \in E_\eta$ and σ smooth. Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

Further, $Y(t) \in D(A)$ for every $t \in (0, 1]$, it satisfies the integral representation

$$Y(t) = y_0 + \int_0^t AY(s) ds + \int_0^t \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

and $F(Y(t)) - F(Y(s)) = \int_s^t \langle DF(Y(r)), AY(s) \rangle dr + \int_s^t \langle DF(Y(r)), \sigma(Y(r)) \rangle dX(r)$



We assume that exist $(E_\eta)_{\eta \geq 0}$, $E_\eta \subseteq E_\theta$ if $\eta \geq \theta$, $E_0 = E$, $E_1 = D(A)$ such that

$$\|e^{tA}\|_{\mathcal{L}(E_\theta, E_\eta)} \leq C t^{\theta-\eta}, \quad \|e^{tA} - Id\|_{\mathcal{L}(E_\eta, E_\theta)} \leq C t^{\eta-\theta}, \quad t \in [0, 1], \quad \eta \geq \theta$$

Mild Sewing Lemma (2) (Gubinelli-Tindel '10, A-Lorenzi-Tessitore '22): If $\alpha \leq 1 < \beta$, then the map $I_S : C_{S,2}^{\alpha,\beta}([0, 1]; E_\eta) \rightarrow C^\alpha([0, 1]; E_\eta)$ satisfies, for every $\varepsilon \in [0, 1]$,

$$R \in C^{\beta-\varepsilon}([0, 1]_{<}^2; E_{\eta+\varepsilon}), \quad R(s, t) := (\delta_{S,1} I_S B)(s, t) - B(s, t)$$

Theorem (A-Lorenzi-Tessitore '22): Let $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta \in (1 - \alpha, \alpha)$, $y_0 \in E_\eta$ and σ smooth. Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

Further, $Y(t) \in D(A)$ for every $t \in (0, 1]$, it satisfies the integral representation

$$Y(t) = y_0 + \int_0^t AY(s) ds + \int_0^t \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

$$\text{and } F(Y(t)) - F(Y(s)) = \int_s^t \langle DF(Y(r)), AY(s) \rangle dr + \int_s^t \langle DF(Y(r)), \sigma(Y(r)) \rangle dX(r)$$

$$\mathcal{B} := \{f \in C([0, 1]; E_\eta) : \delta_{S,1} f \in C^\eta([0, 1]_{<}^2; E_\eta)\} \Rightarrow y_0 \in E_\eta$$



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$

Mild Singular Sewing Lemma (A-Lorenzi-Tessitore '24, Neamtu-Hocquet '24): Let $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $\gamma < \alpha$ and $B \in C_{-\gamma}^\alpha((0, 1]_{<}^2; E_\eta)$ s. t. $\delta_{S,2}B \in C_{-\gamma}^{\alpha+\eta}((0, 1]_{<}^3; E)$.

Then, $I_S B$, defined on $(0, 1]$, extends to $[0, 1]$ with values in E_η and

$$\|\delta_{S,1} I_S B(s, t)\|_{E_{\eta+\varepsilon}} \leq C(\varepsilon, B) |t - s|^{\alpha - \gamma - \varepsilon}, \quad \varepsilon \in [0, \alpha - \gamma]$$



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$

Mild Singular Sewing Lemma (A-Lorenzi-Tessitore '24, Neamtu-Hocquet '24): Let $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $\gamma < \alpha$ and $B \in C_{-\gamma}^\alpha((0, 1]_{<}^2; E_\eta)$ s. t. $\delta_{S,2}B \in C_{-\gamma}^{\alpha+\eta}((0, 1]_{<}^3; E)$.

Then, $I_S B$, defined on $(0, 1]$, extends to $[0, 1]$ with values in E_η and

$$\|\delta_{S,1} I_S B(s, t)\|_{E_{\eta+\varepsilon}} \leq C(\varepsilon, B) |t - s|^{\alpha - \gamma - \varepsilon}, \quad \varepsilon \in [0, \alpha - \gamma]$$

Theorem (A-Lorenzi-Tessitore '24): if $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $y_0 \in E_\theta$ with $\theta \in [0, \eta]$ and $2\eta - \theta < \alpha$, σ smooth.



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$

Mild Singular Sewing Lemma (A-Lorenzi-Tessitore '24, Neamtu-Hocquet '24): Let $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $\gamma < \alpha$ and $B \in C_{-\gamma}^\alpha((0, 1]_{<}^2; E_\eta)$ s. t. $\delta_{S,2}B \in C_{-\gamma}^{\alpha+\eta}((0, 1]_{<}^3; E)$.

Then, $I_S B$, defined on $(0, 1]$, extends to $[0, 1]$ with values in E_η and

$$\|\delta_{S,1} I_S B(s, t)\|_{E_{\eta+\varepsilon}} \leq C(\varepsilon, B) |t - s|^{\alpha - \gamma - \varepsilon}, \quad \varepsilon \in [0, \alpha - \gamma]$$

Theorem (A-Lorenzi-Tessitore '24): if $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $y_0 \in E_\theta$ with $\theta \in [0, \eta]$ and $2\eta - \theta < \alpha$, σ smooth.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$

Mild Singular Sewing Lemma (A-Lorenzi-Tessitore '24, Neamtu-Hocquet '24): Let $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $\gamma < \alpha$ and $B \in C_{-\gamma}^\alpha((0, 1]_{<}^2; E_\eta)$ s. t. $\delta_{S,2}B \in C_{-\gamma}^{\alpha+\eta}((0, 1]_{<}^3; E)$.

Then, $I_S B$, defined on $(0, 1]$, extends to $[0, 1]$ with values in E_η and

$$\|\delta_{S,1} I_S B(s, t)\|_{E_{\eta+\varepsilon}} \leq C(\varepsilon, B) |t - s|^{\alpha - \gamma - \varepsilon}, \quad \varepsilon \in [0, \alpha - \gamma]$$

Theorem (A-Lorenzi-Tessitore '24): if $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $y_0 \in E_\theta$ with $\theta \in [0, \eta]$ and $2\eta - \theta < \alpha$, σ smooth.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

Moreover, $Y(t) \in D(A)$ for every $t \in (0, 1]$.



How can we weaken the condition $y_0 \in E_\eta$? Sewing Lemma with **singularities**: $\gamma \geq 0$

$$C_{-\gamma}^\alpha((0, 1]_{<}^N; E_\eta) := \left\{ B : (0, 1]_{<}^N \rightarrow E_\eta : \sup_{0 \leq t_1 < \dots < t_N \leq 1} (t_1)^\gamma \frac{\|B(t_1, \dots, t_N)\|_\eta}{|t_N - t_1|^\alpha} < \infty \right\}$$

Mild Singular Sewing Lemma (A-Lorenzi-Tessitore '24, Neamtu-Hocquet '24): Let $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $\gamma < \alpha$ and $B \in C_{-\gamma}^\alpha((0, 1]_{<}^2; E_\eta)$ s. t. $\delta_{S,2}B \in C_{-\gamma}^{\alpha+\eta}((0, 1]_{<}^3; E)$.

Then, $I_S B$, defined on $(0, 1]$, extends to $[0, 1]$ with values in E_η and

$$\|\delta_{S,1} I_S B(s, t)\|_{E_{\eta+\varepsilon}} \leq C(\varepsilon, B) |t - s|^{\alpha - \gamma - \varepsilon}, \quad \varepsilon \in [0, \alpha - \gamma]$$

Theorem (A-Lorenzi-Tessitore '24): if $X \in C^\alpha$, $\alpha \in (\frac{1}{2}, 1)$, $\eta > 1 - \alpha$, $y_0 \in E_\theta$ with $\theta \in [0, \eta]$ and $2\eta - \theta < \alpha$, σ smooth.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$, defined as

$$\Gamma(Y)(t) = e^{tA} y_0 + \int_0^t e^{(t-s)A} \sigma(Y(s)) dX(s), \quad t \in [0, 1]$$

Moreover, $Y(t) \in D(A)$ for every $t \in (0, 1]$.

$$\mathcal{B} := \{f \in C_{\theta-\eta}((0, 1]; E_\eta) \cap C_b((0, 1]; E) : \delta_{S,1} f \in C^\eta([0, 1]_{<}^2; E_\eta)\}$$



Mild Integration: the Rough Case

Assume that $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\alpha$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$.



Mild Integration: the Rough Case

Assume that $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\alpha$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. We have

$$\int_s^t e^{(t-s)A} f(r) dX(r) = e^{(t-s)A} f(s) (\delta_1 X)(s, t) + \int_s^t e^{(t-r)A} (f(r) - e^{(r-s)A} f(s)) dX(r)$$



Mild Integration: the Rough Case

Assume that $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\alpha$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. We have

$$\int_s^t e^{(t-s)A} f(r) dX(r) = e^{(t-s)A} f(s) (\delta_1 X)(s, t) + \int_s^t e^{(t-r)A} (f(r) - e^{(r-s)A} f(s)) dX(r)$$

When $A = 0$ and f is controlled by X , we replace $f(r) - e^{(r-s)A} f(s)$ with

$$f(r) - f(s) = f'(s) (\delta_1 X)(s, r) + R(s, r), \quad R(s, r) = O(|r - s|^{2\alpha})$$



Mild Integration: the Rough Case

Assume that $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\alpha$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. We have

$$\int_s^t e^{(t-s)A} f(r) dX(r) = e^{(t-s)A} f(s) (\delta_1 X)(s, t) + \int_s^t e^{(t-r)A} (f(r) - e^{(r-s)A} f(s)) dX(r)$$

When $A = 0$ and f is controlled by X , we replace $f(r) - e^{(r-s)A} f(s)$ with

$$f(r) - f(s) = f'(s) (\delta_1 X)(s, r) + R(s, r), \quad R(s, r) = O(|r - s|^{2\alpha})$$

Gubinelli-Tindel '10, Gerasimovics-Hairer '19: We say that $f : [0, 1] \rightarrow E$ is S -controlled by X if there exists $f' : [0, 1] \rightarrow E$ such that (f' S -Gubinelli derivative of f)

$$R_S^f(s, t) := (\delta_{s,1} f)(s, t) - e^{(t-s)A} f'(s) (\delta_1 X)(s, t), \quad R_S^f \in C^\rho([0, 1]_{<}^2; E), \quad \rho > \alpha$$



Mild Integration: the Rough Case

Assume that $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\alpha$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. We have

$$\int_s^t e^{(t-s)A} f(r) dX(r) = e^{(t-s)A} f(s) (\delta_1 X)(s, t) + \int_s^t e^{(t-r)A} (f(r) - e^{(r-s)A} f(s)) dX(r)$$

When $A = 0$ and f is controlled by X , we replace $f(r) - e^{(r-s)A} f(s)$ with

$$f(r) - f(s) = f'(s) (\delta_1 X)(s, r) + R(s, r), \quad R(s, r) = O(|r - s|^{2\alpha})$$

Gubinelli-Tindel '10, Gerasimovics-Hairer '19: We say that $f : [0, 1] \rightarrow E$ is S -controlled by X if there exists $f' : [0, 1] \rightarrow E$ such that (f' S -Gubinelli derivative of f)

$$R_S^f(s, t) := (\delta_{S,1} f)(s, t) - e^{(t-s)A} f'(s) (\delta_1 X)(s, t), \quad R_S^f \in C^\rho([0, 1]_{<}^2; E), \quad \rho > \alpha$$

For f S -controlled by X , we get

$$\int_s^t e^{(t-r)A} f(r) dX(r) = e^{(t-s)A} f(s) (\delta_1 X)(s, t) + e^{(t-s)A} f'(s) \mathbb{X}(s, t) + O(|t - s|^{\rho+\alpha})$$



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is S -controlled by X , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is S -controlled by X , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$

If $\delta_{S,1}f' \in C^\zeta([0, 1]_{<}^2; E)$, $\zeta + 2\alpha$, $\rho + \alpha > 1$, then $\delta_{S,2}B \in C^\mu([0, 1]_{<}^3; E)$, $\mu > 1$



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is **S-controlled by X** , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$

If $\delta_{S,1}f' \in C^\zeta([0, 1]_{<}^2; E)$, $\zeta + 2\alpha$, $\rho + \alpha > 1$, then $\delta_{S,2}B \in C^\mu([0, 1]_{<}^3; E)$, $\mu > 1$

Mild Sewing Lemma $\Rightarrow I_S B(t) = " \int_0^t e^{(t-r)A}f(r)dX(r) "$ exists

Theorem (A-Lorenzi-Tessitore '25): Let $X \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $\eta \in (1 - 2\alpha, \alpha)$, $y_0 \in E_{\eta+\alpha}$, σ linear and bounded.



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is **S-controlled by X** , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$

If $\delta_{S,1}f' \in C^\zeta([0, 1]_{<}^2; E)$, $\zeta + 2\alpha$, $\rho + \alpha > 1$, then $\delta_{S,2}B \in C^\mu([0, 1]_{<}^3; E)$, $\mu > 1$

Mild Sewing Lemma $\Rightarrow I_S B(t) = " \int_0^t e^{(t-r)A}f(r)d\mathbf{X}(r) "$ exists

Theorem (A-Lorenzi-Tessitore '25): Let $\mathbf{X} \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $\eta \in (1 - 2\alpha, \alpha)$, $y_0 \in E_{\eta+\alpha}$, σ linear and bounded.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ defined as

$$\Gamma(Y)(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))d\mathbf{X}(s), \quad t \in [0, 1]$$



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is **S-controlled by X** , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$

If $\delta_{S,1}f' \in C^\zeta([0, 1]_{<}^2; E)$, $\zeta + 2\alpha$, $\rho + \alpha > 1$, then $\delta_{S,2}B \in C^\mu([0, 1]_{<}^3; E)$, $\mu > 1$

Mild Sewing Lemma $\Rightarrow I_S B(t) = " \int_0^t e^{(t-r)A}f(r)d\mathbf{X}(r) "$ exists

Theorem (A-Lorenzi-Tessitore '25): Let $\mathbf{X} \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $\eta \in (1 - 2\alpha, \alpha)$, $y_0 \in E_{\eta+\alpha}$, σ linear and bounded.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ defined as

$$\Gamma(Y)(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))d\mathbf{X}(s), \quad t \in [0, 1]$$

Moreover, $Y(t) \in D(A)$ for every $t \in (0, 1]$, it satisfies the integral representation

$$Y(t) = y_0 + \int_0^t A Y(s)ds + \int_0^t \sigma(Y(s))d\mathbf{X}(s), \quad t \in [0, 1]$$



Set $B(s, t) = e^{(t-s)A}f(s)(\delta_1 X)(s, t) + e^{(t-s)A}f'(s)\mathbb{X}(s, t)$. If f is **S-controlled by X** , then

$$(\delta_{S,2}B)(s, t, u) = -R_S^f(s, t)(\delta_1 X)(t, u) - (\delta_{S,1}f')(s, t)\mathbb{X}(t, u)$$

If $\delta_{S,1}f' \in C^\zeta([0, 1]_{<}^2; E)$, $\zeta + 2\alpha$, $\rho + \alpha > 1$, then $\delta_{S,2}B \in C^\mu([0, 1]_{<}^3; E)$, $\mu > 1$

Mild Sewing Lemma $\Rightarrow I_S B(t) = \int_0^t e^{(t-r)A}f(r)d\mathbf{X}(r)$ exists

Theorem (A-Lorenzi-Tessitore '25): Let $\mathbf{X} \in \mathcal{C}^\alpha([0, 1])$, $\alpha \in (\frac{1}{3}, \frac{1}{2}]$, $\eta \in (1 - 2\alpha, \alpha)$, $y_0 \in E_{\eta+\alpha}$, σ linear and bounded.

Then, there exists a unique fixed point Y for the operator $\Gamma : \mathcal{B} \rightarrow \mathcal{B}$ defined as

$$\Gamma(Y)(t) = e^{tA}y_0 + \int_0^t e^{(t-s)A}\sigma(Y(s))d\mathbf{X}(s), \quad t \in [0, 1]$$

Moreover, $Y(t) \in D(A)$ for every $t \in (0, 1]$, it satisfies the integral representation

$$Y(t) = y_0 + \int_0^t A Y(s)ds + \int_0^t \sigma(Y(s))d\mathbf{X}(s), \quad t \in [0, 1]$$

$$\text{and } F(Y(t)) - F(Y(s)) = \int_s^t \langle DF(Y(r)), AY(r) \rangle dr + \int_s^t \langle DF(Y(r)), \sigma(Y(r)) \rangle d\mathbf{X}(r) (+?)$$

$$\mathcal{B} = \{f \in \dots : R_S^f \in C^{\alpha+\eta}([0, 1]_{<}^2; E_\eta), f' \in C^\eta([0, 1]_{<}^2; E_\eta)\}$$



THANK YOU FOR YOUR ATTENTION!!

