

# Pinning and Wetting Transition for (1+1)-Dimensional Fields with Laplacian Interaction

Francesco Caravenna

`caravenna@math.unipd.it`

Università degli Studi di Padova

Seminar on Stochastic Processes

Zürich, April 30<sup>th</sup>, 2008

# References

- ▶ [CD1] F. Caravenna and J.-D. Deuschel  
*Pinning and wetting transition for  $(1+1)$ -dimensional fields with Laplacian interaction*, Ann. Probab. (to appear)
- ▶ [CD2] F. Caravenna and J.-D. Deuschel  
*Scaling limits of  $(1+1)$ -dimensional pinning models with Laplacian interaction*, preprint (2008).

# Outline

## 1. The Models

Introduction

Wetting and pinning models

## 2. Free Energy Results

The free energy

The phase transition

The disordered case

## 3. Path Results

Path results

Refined critical scaling limit

## 4. Sketch of the Proofs

Integrated random walk

Markov renewal theory

# Outline

## 1. The Models

Introduction

Wetting and pinning models

## 2. Free Energy Results

The free energy

The phase transition

The disordered case

## 3. Path Results

Path results

Refined critical scaling limit


## 4. Sketch of the Proofs

Integrated random walk

Markov renewal theory


# Some motivations

DNA denaturation transition at high temperature




# Some motivations

DNA denaturation transition at high temperature




(1+1)-dimensional model: field above an impenetrable wall




# Some motivations

DNA denaturation transition at high temperature



(1+1)-dimensional model: field above an impenetrable wall



# The general wetting model

The field  $\varphi = \{\varphi_i\}_{1 \leq i \leq N}$  in the **free case**:

$$\mathbb{P}_{0,N}^w(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{0,N}^w} \prod_{i=1}^N d\varphi_i^+$$

- ▶  $d\varphi_i^+$  is the Lebesgue measure on  $[0, \infty)$
- ▶  $\mathcal{H}_N(\varphi)$  describes the structure of the chain (**to be specified**)
- ▶  $\mathcal{Z}_{0,N}^w$  is the normalization constant (**partition function**)



# The general wetting model

The field  $\varphi = \{\varphi_i\}_{1 \leq i \leq N}$  in the **interacting case**:

$$\mathbb{P}_{\varepsilon, N}^w(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{\varepsilon, N}^w} \prod_{i=1}^N (d\varphi_i^+ + \varepsilon \cdot \delta_0(d\varphi_i))$$

- ▶  $d\varphi_i^+$  is the Lebesgue measure on  $[0, \infty)$
- ▶  $\mathcal{H}_N(\varphi)$  describes the structure of the chain (to be specified)
- ▶  $\mathcal{Z}_{\varepsilon, N}^w$  is the normalization constant (partition function)
- ▶  $\delta_0(\cdot)$  is the Dirac mass at zero
- ▶  $\varepsilon \geq 0$  is the strength of the pinning interaction

# The general pinning model

Analogous to the wetting case but **without repulsion**:  $d\varphi_i^+ \rightarrow d\varphi_i$


# The general pinning model

Analogous to the wetting case but **without repulsion**:  $d\varphi_i^+ \rightarrow d\varphi_i$



$$\mathbb{P}_{\varepsilon, N}^p(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{\varepsilon, N}^p} \prod_{i=1}^N (d\varphi_i + \varepsilon \cdot \delta_0(d\varphi_i))$$

- ▶  $d\varphi_i$  is the Lebesgue measure on  $\mathbb{R}$
- ▶  $\mathcal{H}_N(\varphi)$  is the same Hamiltonian (to be specified)
- ▶  $\mathcal{Z}_{\varepsilon, N}^p$  is the normalization constant (partition function)
- ▶  $\delta_0(\cdot)$  is the Dirac mass at zero
- ▶  $\varepsilon \geq 0$  is the strength of the pinning interaction

# Pinning VS Wetting (+ boundary conditions)



# Pinning VS Wetting (+ boundary conditions)



# Some questions

Once  $\mathcal{H}_N(\varphi)$  is chosen and  $\varepsilon \geq 0$  is fixed:

- ▶ What are the properties of  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  for large  $N$ ?

# Some questions

Once  $\mathcal{H}_N(\varphi)$  is chosen and  $\varepsilon \geq 0$  is fixed:

- ▶ What are the properties of  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  for large  $N$ ?
- ▶ Is the field **localized** at  $\varphi = 0$  or **delocalized** in  $\varphi \neq 0$ ?  
(How to define localization and delocalization?)

# Some questions

Once  $\mathcal{H}_N(\varphi)$  is chosen and  $\varepsilon \geq 0$  is fixed:

- ▶ What are the properties of  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  for large  $N$ ?
- ▶ Is the field **localized** at  $\varphi = 0$  or **delocalized** in  $\varphi \neq 0$ ?  
(How to define localization and delocalization?)
- ▶ Does the answer depend on  $\varepsilon \geq 0$  (**phase transition**) and/or on the choice of  $\mathcal{H}_N(\varphi)$ ?



# Some questions

Once  $\mathcal{H}_N(\varphi)$  is chosen and  $\varepsilon \geq 0$  is fixed:

- ▶ What are the properties of  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  for large  $N$ ?
- ▶ Is the field **localized** at  $\varphi = 0$  or **delocalized** in  $\varphi \neq 0$ ?  
(How to define localization and delocalization?)
- ▶ Does the answer depend on  $\varepsilon \geq 0$  (**phase transition**) and/or on the choice of  $\mathcal{H}_N(\varphi)$ ?

How to choose  $\mathcal{H}_N(\varphi)$ ?

# The choice of $\mathcal{H}_N(\varphi)$

The simplest choice is the **gradient case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=1}^N V(\nabla \varphi_i), \quad \nabla \varphi_i := \varphi_i - \varphi_{i-1},$$

# The choice of $\mathcal{H}_N(\varphi)$

The simplest choice is the **gradient case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=1}^N V(\nabla \varphi_i), \quad \nabla \varphi_i := \varphi_i - \varphi_{i-1},$$

$$V(\cdot) : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\} : \quad \int_{\mathbb{R}} e^{-V(x)} dx < +\infty \quad (= 1).$$

+ regularity (see later). **Gaussian case:**  $V(x) \propto x^2$

# The choice of $\mathcal{H}_N(\varphi)$

The simplest choice is the **gradient case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=1}^N V(\nabla \varphi_i), \quad \nabla \varphi_i := \varphi_i - \varphi_{i-1},$$

$$V(\cdot) : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\} : \quad \int_{\mathbb{R}} e^{-V(x)} dx < +\infty \quad (= 1).$$

+ regularity (see later). **Gaussian case:**  $V(x) \propto x^2$

- ▶  $\mathbb{P}_{0,N}^p$  is (the bridge of) a **random walk** of step law  $e^{-V(x)}$
- ▶  $\mathbb{P}_{0,N}^w$  is (the bridge of) a random walk **conditioned to stay  $\geq 0$**

# The choice of $\mathcal{H}_N(\varphi)$

The simplest choice is the **gradient case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=1}^N V(\nabla \varphi_i), \quad \nabla \varphi_i := \varphi_i - \varphi_{i-1},$$

$$V(\cdot) : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\} : \quad \int_{\mathbb{R}} e^{-V(x)} dx < +\infty \quad (= 1).$$

+ regularity (see later). **Gaussian case:**  $V(x) \propto x^2$

- ▶  $\mathbb{P}_{0,N}^p$  is (the bridge of) a **random walk** of step law  $e^{-V(x)}$
- ▶  $\mathbb{P}_{0,N}^w$  is (the bridge of) a random walk **conditioned to stay  $\geq 0$**

[Isozaki, Yoshida SPA 01] [Deuschel, Giacomin, Zambotti PTRF 05]

[Caravenna, Giacomin, Zambotti EJP 06]

# The choice of $\mathcal{H}_N(\varphi)$

We rather consider the **Laplacian case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=0}^N V(\Delta\varphi_i)$$

$$\Delta\varphi_i := \nabla\varphi_{i+1} - \nabla\varphi_i = \varphi_{i+1} + \varphi_{i-1} - 2\varphi_i$$

# The choice of $\mathcal{H}_N(\varphi)$

We rather consider the **Laplacian case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=0}^N V(\Delta\varphi_i)$$

$$\Delta\varphi_i := \nabla\varphi_{i+1} - \nabla\varphi_i = \varphi_{i+1} + \varphi_{i-1} - 2\varphi_i$$

- ▶ Simplest non-nearest neighbor interaction
- ▶ **Semi-flexible polymers**:  $\Delta$  favors affine configurations

# The choice of $\mathcal{H}_N(\varphi)$

We rather consider the **Laplacian case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=0}^N V(\Delta\varphi_i) + \tilde{V}(\nabla\varphi_i)$$

$$\Delta\varphi_i := \nabla\varphi_{i+1} - \nabla\varphi_i = \varphi_{i+1} + \varphi_{i-1} - 2\varphi_i$$

- ▶ Simplest non-nearest neighbor interaction
- ▶ **Semi-flexible polymers**:  $\Delta$  favors affine configurations
- ▶  $\nabla$  and  $\Delta$  together?



# The choice of $\mathcal{H}_N(\varphi)$

We rather consider the **Laplacian case**:

$$\mathcal{H}_N(\varphi) := \sum_{i=0}^N V(\Delta\varphi_i)$$

$$\Delta\varphi_i := \nabla\varphi_{i+1} - \nabla\varphi_i = \varphi_{i+1} + \varphi_{i-1} - 2\varphi_i$$

- ▶ Simplest non-nearest neighbor interaction
- ▶ **Semi-flexible polymers**:  $\Delta$  favors affine configurations
- ▶  $\nabla$  and  $\Delta$  together?

Interpretation of the free case  $\varepsilon = 0$ :

- ▶  $\mathbb{P}_{0,N}^p$  is (the bridge of) the integral of a **random walk**
- ▶  $\mathbb{P}_{0,N}^w$  is further **conditioned to stay  $\geq 0$**

# Laplacian interaction in $(d + 1)$ -dimension

Fields  $\varphi : \{1, \dots, N\}^d \rightarrow \mathbb{R}$  with Laplacian interaction for  $d \geq 2$  are models for **semiflexible membranes**

# Laplacian interaction in $(d + 1)$ -dimension

Fields  $\varphi : \{1, \dots, N\}^d \rightarrow \mathbb{R}$  with Laplacian interaction for  $d \geq 2$  are models for [semiflexible membranes](#)

The problem of [entropic repulsion](#) (probability for the field to stay non-negative) has been studied in the [Gaussian case](#):

[Sakagawa JMP 04] [Kurt SPA 07] [Kurt preprint 07]

# Laplacian interaction in $(d + 1)$ -dimension

Fields  $\varphi : \{1, \dots, N\}^d \rightarrow \mathbb{R}$  with Laplacian interaction for  $d \geq 2$  are models for [semiflexible membranes](#)

The problem of [entropic repulsion](#) (probability for the field to stay non-negative) has been studied in the [Gaussian case](#):

[Sakagawa JMP 04] [Kurt SPA 07] [Kurt preprint 07]

---

Henceforth we study  $\mathbb{P}_{\varepsilon, N}^{\text{P}}$  and  $\mathbb{P}_{\varepsilon, N}^{\text{W}}$  with [Laplacian interaction](#) and boundary conditions  $\varphi_{-1} = \varphi_0 = \varphi_N = \varphi_{N+1} = 0$

# Laplacian interaction in $(d + 1)$ -dimension

Fields  $\varphi : \{1, \dots, N\}^d \rightarrow \mathbb{R}$  with Laplacian interaction for  $d \geq 2$  are models for [semiflexible membranes](#)

The problem of [entropic repulsion](#) (probability for the field to stay non-negative) has been studied in the [Gaussian case](#):

[Sakagawa JMP 04] [Kurt SPA 07] [Kurt preprint 07]

Henceforth we study  $\mathbb{P}_{\varepsilon, N}^{\text{P}}$  and  $\mathbb{P}_{\varepsilon, N}^{\text{W}}$  with [Laplacian interaction](#) and boundary conditions  $\varphi_{-1} = \varphi_0 = \varphi_N = \varphi_{N+1} = 0$

Assumptions on  $V$ :

$$\int_{\mathbb{R}} e^{-V(x)} dx = 1, \quad \int_{\mathbb{R}} x e^{-V(x)} dx = 0, \quad \int_{\mathbb{R}} x^2 e^{-V(x)} dx = 1$$

+ regularity:  $x \mapsto e^{-V(x)}$  continuous and  $V(0) < +\infty$ .

# Outline

## 1. The Models

Introduction

Wetting and pinning models

## 2. Free Energy Results

The free energy

The phase transition

The disordered case

## 3. Path Results

Path results

Refined critical scaling limit

## 4. Sketch of the Proofs

Integrated random walk

Markov renewal theory

# How to define localization and delocalization?

Recall the **partition function**: (zero boundary conditions)

$$\mathcal{Z}_{\varepsilon, N}^a = \int_{\Omega_N^a} e^{-\mathcal{H}_N(\varphi)} \prod_{i=1}^{N-1} (d\varphi_i + \varepsilon \delta_0(d\varphi_i))$$

where  $a \in \{p, w\}$  and  $\Omega_N^p = \mathbb{R}^{N-1}$  while  $\Omega_N^w = [0, \infty)^{N-1}$

# How to define localization and delocalization?

Recall the **partition function**: (zero boundary conditions)

$$\mathcal{Z}_{\varepsilon, N}^a = \int_{\Omega_N^a} e^{-\mathcal{H}_N(\varphi)} \prod_{i=1}^{N-1} (d\varphi_i + \varepsilon \delta_0(d\varphi_i))$$

where  $a \in \{p, w\}$  and  $\Omega_N^p = \mathbb{R}^{N-1}$  while  $\Omega_N^w = [0, \infty)^{N-1}$

## Free Energy

$$F^a(\varepsilon) := \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z}_{\varepsilon, N}^a \quad (\text{super-additivity})$$



# How to define localization and delocalization?

Recall the **partition function**: (zero boundary conditions)

$$\mathcal{Z}_{\varepsilon, N}^a = \int_{\Omega_N^a} e^{-\mathcal{H}_N(\varphi)} \prod_{i=1}^{N-1} (d\varphi_i + \varepsilon \delta_0(d\varphi_i))$$

where  $a \in \{p, w\}$  and  $\Omega_N^p = \mathbb{R}^{N-1}$  while  $\Omega_N^w = [0, \infty)^{N-1}$

## Free Energy

$$F^a(\varepsilon) := \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z}_{\varepsilon, N}^a \quad (\text{super-additivity})$$

**Basic observation:**  $F^a(\varepsilon) \geq F^a(0) = 0$  for all  $\varepsilon \geq 0$  and  $a \in \{p, w\}$

$$\mathcal{Z}_{\varepsilon, N}^a \geq \mathcal{Z}_{0, N}^a \approx N^{-c} \quad (c > 0)$$

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

Setting  $\ell_N := \#\{i \leq N : \varphi_i = 0\}$  we have:

► if  $\varepsilon > \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow D^a(\varepsilon) > 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

Setting  $\ell_N := \#\{i \leq N : \varphi_i = 0\}$  we have:

- ▶ if  $\varepsilon > \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow D^a(\varepsilon) > 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability
- ▶ if  $\varepsilon < \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability (**delocalization**)

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

Setting  $\ell_N := \#\{i \leq N : \varphi_i = 0\}$  we have:

- ▶ if  $\varepsilon > \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow D^a(\varepsilon) > 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability
- ▶ if  $\varepsilon < \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability (**delocalization**)
- ▶ if  $\varepsilon = \varepsilon_c^a$ ? Depends on the model:

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

Setting  $\ell_N := \#\{i \leq N : \varphi_i = 0\}$  we have:

- ▶ if  $\varepsilon > \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow D^a(\varepsilon) > 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability
- ▶ if  $\varepsilon < \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability (**delocalization**)
- ▶ if  $\varepsilon = \varepsilon_c^a$ ? Depends on the model:
  - ▶ if  $F^a(\varepsilon_c^a + h) = o(h)$  [**> 1<sup>st</sup> order trans.**]  $\varepsilon = \varepsilon_c^a$  is **delocalized**

# Localization and delocalization

## Definition

The  $a$ -model at  $\varepsilon \geq 0$  is said to be **localized** if  $F^a(\varepsilon) > 0$ .

$$\text{localized} \iff \varepsilon > \varepsilon_c^a := \sup\{\varepsilon \geq 0 : F^a(\varepsilon) = 0\} \in [0, \infty]$$

Setting  $\ell_N := \#\{i \leq N : \varphi_i = 0\}$  we have:

- ▶ if  $\varepsilon > \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow D^a(\varepsilon) > 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability
- ▶ if  $\varepsilon < \varepsilon_c^a$  then  $\frac{\ell_N}{N} \rightarrow 0$  in  $\mathbb{P}_{\varepsilon, N}^a$ -probability (**delocalization**)
- ▶ if  $\varepsilon = \varepsilon_c^a$ ? Depends on the model:
  - ▶ if  $F^a(\varepsilon_c^a + h) = o(h)$  [**> 1<sup>st</sup> order trans.**]  $\varepsilon = \varepsilon_c^a$  is **delocalized**
  - ▶ if  $F^a(\varepsilon_c^a + h) \geq C h$  [**1<sup>st</sup> order trans.**]  $\varepsilon = \varepsilon_c^a$  may be **localized** (phase coexistence, dependence of boundary conditions)



# The phase transition

## Theorem ([CD1])

Both  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  undergo a non-trivial phase transition:

$$0 < \varepsilon_c^p < \varepsilon_c^w < \infty$$

and  $F^a(\varepsilon)$  is analytic on  $[0, \varepsilon_c^a) \cup (\varepsilon_c^a, \infty)$ . (*variational formula*)

# The phase transition

## Theorem ([CD1])

Both  $\mathbb{P}_{\varepsilon,N}^p$  and  $\mathbb{P}_{\varepsilon,N}^w$  undergo a non-trivial phase transition:

$$0 < \varepsilon_c^p < \varepsilon_c^w < \infty$$

and  $F^a(\varepsilon)$  is analytic on  $[0, \varepsilon_c^a) \cup (\varepsilon_c^a, \infty)$ . (*variational formula*)

► In the *pinning model* the transition is exactly of  $2^{nd}$  order:

$$C_1 \frac{h}{\log \frac{1}{h}} \leq F^p(\varepsilon_c^p + h) \leq o(h)$$

# The phase transition

## Theorem ([CD1])

Both  $\mathbb{P}_{\varepsilon, N}^p$  and  $\mathbb{P}_{\varepsilon, N}^w$  undergo a non-trivial phase transition:

$$0 < \varepsilon_c^p < \varepsilon_c^w < \infty$$

and  $F^a(\varepsilon)$  is analytic on  $[0, \varepsilon_c^a) \cup (\varepsilon_c^a, \infty)$ . (*variational formula*)

► In the *pinning model* the transition is exactly of *2<sup>nd</sup> order*:

$$C_1 \frac{h}{\log \frac{1}{h}} \leq F^p(\varepsilon_c^p + h) \leq o(h)$$

► In the *wetting model* the transition is of *1<sup>st</sup> order*:

$$F^w(\varepsilon_c^w + h) \sim C_2 h, \quad \ell_N \sim D N \quad (D > 0)$$

# The gradient case

## Differences in the gradient case

- ▶ the transition is non-trivial only in the **wetting model**:

$$\varepsilon_c^{p,\nabla} = 0, \quad 0 < \varepsilon_c^{w,\nabla} < \infty$$

# The gradient case

## Differences in the gradient case

- ▶ the transition is non-trivial only in the **wetting model**:

$$\varepsilon_c^{p,\nabla} = 0, \quad 0 < \varepsilon_c^{w,\nabla} < \infty$$

- ▶ the transition is of **2<sup>nd</sup> order**:

$$F^p(\varepsilon_c^{p,\nabla} + h) \sim C_p h^2, \quad F^w(\varepsilon_c^{w,\nabla} + h) \sim C_w h^2$$

# The gradient case



## Differences in the gradient case

- ▶ the transition is non-trivial only in the **wetting model**:

$$\varepsilon_c^{p,\nabla} = 0, \quad 0 < \varepsilon_c^{w,\nabla} < \infty$$

- ▶ the transition is of **2<sup>nd</sup> order**:

$$F^p(\varepsilon_c^{p,\nabla} + h) \sim C_p h^2, \quad F^w(\varepsilon_c^{w,\nabla} + h) \sim C_w h^2$$



# A look at the disordered case

Disordered version of our model: ( $d\varphi_i^p = d\varphi_i$  and  $d\varphi_i^w = d\varphi_i^+$ )

$$\mathbb{P}_{\varepsilon, \beta, \omega, N}^a(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{\varepsilon, \beta, \omega, N}^a} \prod_{i=1}^N (d\varphi_i^a + \varepsilon e^{\beta \omega_i} \delta_0(d\varphi_i))$$

where  $\beta \geq 0$  and  $\{\omega_i\}_{i \in \mathbb{N}}$  are IID  $\mathcal{N}(0, 1)$  (law  $P$  indep.  $\mathbb{P}^a$ ).

# A look at the disordered case

**Disordered version** of our model: ( $d\varphi_i^p = d\varphi_i$  and  $d\varphi_i^w = d\varphi_i^+$ )

$$\mathbb{P}_{\varepsilon, \beta, \omega, N}^a(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{\varepsilon, \beta, \omega, N}^a} \prod_{i=1}^N (d\varphi_i^a + \varepsilon e^{\beta \omega_i} \delta_0(d\varphi_i))$$

where  $\beta \geq 0$  and  $\{\omega_i\}_{i \in \mathbb{N}}$  are IID  $\mathcal{N}(0, 1)$  (law  $P$  indep.  $\mathbb{P}^a$ ).

## Quenched free energy

$$F^a(\varepsilon, \beta) := \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z}_{\varepsilon, \beta, \omega, N}^a \geq 0$$

exists  $P(d\omega)$ -a.s. and **does not depend on  $\omega$**  (self-averaging)



# A look at the disordered case

**Disordered version** of our model: ( $d\varphi_i^p = d\varphi_i$  and  $d\varphi_i^w = d\varphi_i^+$ )

$$\mathbb{P}_{\varepsilon, \beta, \omega, N}^a(d\varphi_1, \dots, d\varphi_N) := \frac{e^{-\mathcal{H}_N(\varphi)}}{\mathcal{Z}_{\varepsilon, \beta, \omega, N}^a} \prod_{i=1}^N (d\varphi_i^a + \varepsilon e^{\beta \omega_i} \delta_0(d\varphi_i))$$

where  $\beta \geq 0$  and  $\{\omega_i\}_{i \in \mathbb{N}}$  are IID  $\mathcal{N}(0, 1)$  (law  $P$  indep.  $\mathbb{P}^a$ ).

## Quenched free energy

$$F^a(\varepsilon, \beta) := \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathcal{Z}_{\varepsilon, \beta, \omega, N}^a \geq 0$$

exists  $P(d\omega)$ -a.s. and **does not depend on  $\omega$**  (self-averaging)

**Localization:**  $F^a(\varepsilon, \beta) > 0 \iff \varepsilon > \varepsilon_c^a(\beta)$  (**critical line**)

# Smoothing effect of disorder

What is the behavior of  $\varepsilon_c^a(\beta)$  for small  $\beta$  ? ( $\varepsilon_c^a = \varepsilon_c^a(0)$ )

What is the **regularity of the transition** in the disordered case?

# Smoothing effect of disorder

What is the behavior of  $\varepsilon_c^a(\beta)$  for small  $\beta$  ? ( $\varepsilon_c^a = \varepsilon_c^a(0)$ )

What is the **regularity of the transition** in the disordered case?

**Theorem ([Giacomin and Toninelli, CMP 06])**

*Both in the  $\nabla$  and  $\Delta$  case, both for  $a = p$  and for  $a = w$ :  
for every  $\beta > 0$  there exists  $C_\beta > 0$  such that*

$$F^a(\varepsilon_c^a(\beta) + h) \leq C_\beta h^2$$

When disorder is present the transition is **at least of 2<sup>nd</sup> order**

**Sharp contrast** with homogeneous case ► cf.

# Smoothing effect of disorder

What is the behavior of  $\varepsilon_c^a(\beta)$  for small  $\beta$  ? ( $\varepsilon_c^a = \varepsilon_c^a(0)$ )

What is the **regularity of the transition** in the disordered case?

**Theorem ([Giacomin and Toninelli, CMP 06])**

*Both in the  $\nabla$  and  $\Delta$  case, both for  $a = p$  and for  $a = w$ :  
for every  $\beta > 0$  there exists  $C_\beta > 0$  such that*

$$F^a(\varepsilon_c^a(\beta) + h) \leq C_\beta h^2$$

When disorder is present the transition is **at least of 2<sup>nd</sup> order**

**Sharp contrast** with homogeneous case ► cf.

**Very general proof:** rare-stretches in  $\omega$  (Large Deviations)

# Outline

## 1. The Models

Introduction

Wetting and pinning models

## 2. Free Energy Results

The free energy

The phase transition

The disordered case

## 3. Path Results

Path results

Refined critical scaling limit

## 4. Sketch of the Proofs

Integrated random walk

Markov renewal theory

# Some deeper questions

We have established the existence of a **phase transition**:

$$\ell_N = o(N) \text{ if } \varepsilon < \varepsilon_c^a \quad \text{VS} \quad \ell_N \sim D \cdot N \text{ if } \varepsilon > \varepsilon_c^a$$

Can we say something more precise? Typical paths of  $\mathbb{P}_{\varepsilon, N}^a$ ?

# Some deeper questions

We have established the existence of a **phase transition**:

$$\ell_N = o(N) \text{ if } \varepsilon < \varepsilon_c^a \quad \text{VS} \quad \ell_N \sim D \cdot N \text{ if } \varepsilon > \varepsilon_c^a$$

Can we say something more precise? Typical paths of  $\mathbb{P}_{\varepsilon, N}^a$ ?

**Yes** in the **pinning case** and under additional assumptions on  $V(\cdot)$ :

- ▶ **symmetry**:  $V(x) = V(-x)$  for every  $x \in \mathbb{R}$
- ▶ **uniform strict convexity**:  $\exists \gamma > 0$  s. t.  $V(x) - \gamma \frac{x^2}{2}$  is convex
- ▶ **regularity**:  $x \mapsto e^{-V(x)}$  is continuous and  $V(0) < \infty$

$$\int_{\mathbb{R}} e^{-V(x)} dx = 1 \qquad \int_{\mathbb{R}} x^2 e^{-V(x)} dx = 1$$

# Some deeper questions

We have established the existence of a **phase transition**:

$$\ell_N = o(N) \text{ if } \varepsilon < \varepsilon_c^a \quad \text{VS} \quad \ell_N \sim D \cdot N \text{ if } \varepsilon > \varepsilon_c^a$$

Can we say something more precise? Typical paths of  $\mathbb{P}_{\varepsilon, N}^a$ ?

Yes in the **pinning case** and under additional assumptions on  $V(\cdot)$ :

- ▶ **symmetry**:  $V(x) = V(-x)$  for every  $x \in \mathbb{R}$
- ▶ **uniform strict convexity**:  $\exists \gamma > 0$  s. t.  $V(x) - \gamma \frac{x^2}{2}$  is convex
- ▶ **regularity**:  $x \mapsto e^{-V(x)}$  is continuous and  $V(0) < \infty$

$$\int_{\mathbb{R}} e^{-V(x)} dx = 1 \qquad \int_{\mathbb{R}} x^2 e^{-V(x)} dx = 1$$

Let us set  $M_N := \max_{1 \leq i \leq N} |\varphi_i|$



# A closer look at the typical paths [CD2]

► delocalized regime  $\varepsilon < \varepsilon_c^p$ :  $\ell_N = O(1)$  and  $M_N \approx N^{3/2}$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p(\varphi_i \neq 0 \text{ for } i \in \{K, N - K\}) = 1$$

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p\left(\frac{1}{K} N^{3/2} \leq M_N \leq K N^{3/2}\right) = 1$$

# A closer look at the typical paths [CD2]

- delocalized regime  $\varepsilon < \varepsilon_c^p$ :  $\ell_N = O(1)$  and  $M_N \approx N^{3/2}$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p(\varphi_i \neq 0 \text{ for } i \in \{K, N - K\}) = 1$$

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p\left(\frac{1}{K} N^{3/2} \leq M_N \leq K N^{3/2}\right) = 1$$

- localized regime  $\varepsilon > \varepsilon_c^p$ :  $M_N = O((\log N)^2)$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p(M_N \leq K (\log N)^2) = 1$$

# A closer look at the typical paths [CD2]

- delocalized regime  $\varepsilon < \varepsilon_c^p$ :  $\ell_N = O(1)$  and  $M_N \approx N^{3/2}$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p(\varphi_i \neq 0 \text{ for } i \in \{K, N - K\}) = 1$$

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p\left(\frac{1}{K} N^{3/2} \leq M_N \leq K N^{3/2}\right) = 1$$

- localized regime  $\varepsilon > \varepsilon_c^p$ :  $M_N = O((\log N)^2)$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon, N}^p(M_N \leq K (\log N)^2) = 1$$

- critical regime  $\varepsilon = \varepsilon_c^p$ :  $M_N \approx \frac{N^{3/2}}{(\log N)^c}$  and  $\ell_N \approx \frac{N}{\log N}$ :

$$\lim_{K \rightarrow \infty} \liminf_{N \rightarrow \infty} \mathbb{P}_{\varepsilon_c^p, N}^p\left(\frac{1}{K} \frac{N^{3/2}}{(\log N)^{3/2}} \leq M_N \leq K \frac{N^{3/2}}{\log N}\right) = 1$$

# Scaling Limits

We rescale and interpolate linearly the field: for  $t \in [0, 1]$

$$\widehat{\varphi}_N(t) := \frac{\varphi_{\lfloor Nt \rfloor}}{N^{3/2}} + (Nt - \lfloor Nt \rfloor) \frac{\varphi_{\lfloor Nt \rfloor + 1} - \varphi_{\lfloor Nt \rfloor}}{N^{3/2}}$$

# Scaling Limits

We rescale and interpolate linearly the field: for  $t \in [0, 1]$

$$\widehat{\varphi}_N(t) := \frac{\varphi_{\lfloor Nt \rfloor}}{N^{3/2}} + (Nt - \lfloor Nt \rfloor) \frac{\varphi_{\lfloor Nt \rfloor + 1} - \varphi_{\lfloor Nt \rfloor}}{N^{3/2}}$$

Let  $\{B_t\}_{t \in [0,1]}$  **standard BM**,  $I_t := \int_0^t B_s \, ds$  **integrated BM**

$$\{(\widehat{B}_t, \widehat{I}_t)\}_{t \in [0,1]} = \{(B_t, I_t)\}_{t \in [0,1]} \quad \text{cond. on} \quad (B_1, I_1) = (0, 0)$$

# Scaling Limits

We rescale and interpolate linearly the field: for  $t \in [0, 1]$

$$\widehat{\varphi}_N(t) := \frac{\varphi_{\lfloor Nt \rfloor}}{N^{3/2}} + (Nt - \lfloor Nt \rfloor) \frac{\varphi_{\lfloor Nt \rfloor + 1} - \varphi_{\lfloor Nt \rfloor}}{N^{3/2}}$$

Let  $\{B_t\}_{t \in [0,1]}$  **standard BM**,  $I_t := \int_0^t B_s \, ds$  **integrated BM**

$$\{(\widehat{B}_t, \widehat{I}_t)\}_{t \in [0,1]} = \{(B_t, I_t)\}_{t \in [0,1]} \quad \text{cond. on} \quad (B_1, I_1) = (0, 0)$$

## Theorem (Scaling Limits [CD2])

*The rescaled field  $\{\widehat{\varphi}_N(t)\}_{t \in [0,1]}$  under  $\mathbb{P}_{\varepsilon, N}^p$  converges in distribution on  $C([0, 1])$  as  $N \rightarrow \infty$ , for every  $\varepsilon \geq 0$ . The limit is*

- ▶ *If  $\varepsilon < \varepsilon_c^p$ , the law of  $\{\widehat{I}_t\}_{t \in [0,1]}$*
- ▶ *If  $\varepsilon = \varepsilon_c^p$  or  $\varepsilon > \varepsilon_c^p$ , the law concentrated on  $f(t) \equiv 0$*

# The critical regime

For  $\varepsilon = \varepsilon_c^p$  the field has **very large fluctuations** ( $\approx \frac{N^{3/2}}{(\log N)^c}$ ).

Can we extract a **non-trivial scaling limit**?

# The critical regime

For  $\varepsilon = \varepsilon_c^p$  the field has **very large fluctuations** ( $\approx \frac{N^{3/2}}{(\log N)^c}$ ).

Can we extract a **non-trivial scaling limit**?

**Not** in  $C([0, 1])$  or  $D([0, 1])$ :

$$\frac{1}{N} \{i \in \{1, \dots, N\} : \varphi_i = 0\} \quad \text{becomes dense in } [0, 1]$$




# The critical regime

For  $\varepsilon = \varepsilon_c^p$  the field has **very large fluctuations** ( $\approx \frac{N^{3/2}}{(\log N)^c}$ ).

Can we extract a **non-trivial scaling limit**?

**Not** in  $C([0, 1])$  or  $D([0, 1])$ :

$$\frac{1}{N} \{i \in \{1, \dots, N\} : \varphi_i = 0\} \quad \text{becomes dense in } [0, 1]$$



# The critical regime

**Alternative idea:** look at the field in a **distributional** sense

$$\mu_N(dt) := \frac{(\log N)^{5/2}}{N^{3/2}} \varphi_{\lfloor Nt \rfloor} dt = \tilde{\varphi}_N(t) dt$$

$\mu_N(\cdot)$  is a (random) **finite signed measure** on  $[0, 1]$

# The critical regime

**Alternative idea:** look at the field in a **distributional sense**

$$\mu_N(dt) := \frac{(\log N)^{5/2}}{N^{3/2}} \varphi_{\lfloor Nt \rfloor} dt = \tilde{\varphi}_N(t) dt$$

$\mu_N(\cdot)$  is a (random) **finite signed measure** on  $[0, 1]$

Let  $\{L_t\}_{t \in [0,1]}$  be the **stable symmetric Lévy process** of index  $\frac{2}{5}$

0 drift      0 Brownian component       $\Pi(dx) = \frac{c}{|x|^{7/5}} dx$

# The critical regime

**Alternative idea:** look at the field in a **distributional sense**

$$\mu_N(dt) := \frac{(\log N)^{5/2}}{N^{3/2}} \varphi_{\lfloor Nt \rfloor} dt = \tilde{\varphi}_N(t) dt$$

$\mu_N(\cdot)$  is a (random) **finite signed measure** on  $[0, 1]$

Let  $\{L_t\}_{t \in [0,1]}$  be the **stable symmetric Lévy process** of index  $\frac{2}{5}$

0 drift      0 Brownian component       $\Pi(dx) = \frac{c}{|x|^{7/5}} dx$

The paths of  $L$  are a.s. of bounded variation, hence we set

$$dL((a, b)) := L_b - L_a$$

# The critical regime

**Alternative idea:** look at the field in a **distributional sense**

$$\mu_N(dt) := \frac{(\log N)^{5/2}}{N^{3/2}} \varphi_{[Nt]} dt = \tilde{\varphi}_N(t) dt$$

$\mu_N(\cdot)$  is a (random) **finite signed measure** on  $[0, 1]$

Let  $\{L_t\}_{t \in [0,1]}$  be the **stable symmetric Lévy process** of index  $\frac{2}{5}$

0 drift      0 Brownian component       $\Pi(dx) = \frac{c}{|x|^{7/5}} dx$


The paths of  $L$  are a.s. of bounded variation, hence we set

$$dL((a, b)) := L_b - L_a$$

## Theorem ([CD2])

*The random signed measure  $\mu_N$  under  $\mathbb{P}_{\varepsilon_c^p, N}^p$  converges in distribution as  $N \rightarrow \infty$  toward  $dL$ .*

# The critical regime



# Outline

## 1. The Models

Introduction

Wetting and pinning models

## 2. Free Energy Results

The free energy

The phase transition

The disordered case

## 3. Path Results

Path results

Refined critical scaling limit

## 4. Sketch of the Proofs

Integrated random walk

Markov renewal theory

# A random walk viewpoint ( $\varepsilon = 0$ )

Let  $\{X_i\}_{i \in \mathbb{N}}$  be IID random variables with law

$$\mathbf{P}(X_i \in dx) := e^{-V(x)} dx \quad \mathbf{E}(X_i) = 0 \quad \mathbf{Var}(X_i) = 1$$



# A random walk viewpoint ( $\varepsilon = 0$ )

Let  $\{X_i\}_{i \in \mathbb{N}}$  be IID random variables with law

$$\mathbf{P}(X_i \in dx) := e^{-V(x)} dx \quad \mathbf{E}(X_i) = 0 \quad \mathbf{Var}(X_i) = 1$$

Random walk:  $Y_n := X_1 + \dots + X_n$

Integrated random walk:

$$Z_n := Y_1 + \dots + Y_n = nX_1 + (n-1)X_2 + \dots + X_n$$

# A random walk viewpoint ( $\varepsilon = 0$ )

Let  $\{X_i\}_{i \in \mathbb{N}}$  be IID random variables with law

$$\mathbf{P}(X_i \in dx) := e^{-V(x)} dx \quad \mathbf{E}(X_i) = 0 \quad \mathbf{Var}(X_i) = 1$$

Random walk:  $Y_n := X_1 + \dots + X_n$

Integrated random walk:

$$Z_n := Y_1 + \dots + Y_n = nX_1 + (n-1)X_2 + \dots + X_n$$

The free case  $\varepsilon = 0$

- ▶ The field  $\{\varphi_i\}_{1 \leq i \leq N}$  under  $\mathbb{P}_{0,N}^{\mathbb{P}}$  is distributed like  $\{Z_i\}_{1 \leq i \leq N}$  conditionally on  $(Y_N, Z_N) = (0, 0)$ . ( $Z_{N/2} \approx N^{3/2}$ )

# A random walk viewpoint ( $\varepsilon = 0$ )

Let  $\{X_i\}_{i \in \mathbb{N}}$  be IID random variables with law

$$\mathbf{P}(X_i \in dx) := e^{-V(x)} dx \quad \mathbf{E}(X_i) = 0 \quad \mathbf{Var}(X_i) = 1$$

Random walk:  $Y_n := X_1 + \dots + X_n$

Integrated random walk:

$$Z_n := Y_1 + \dots + Y_n = nX_1 + (n-1)X_2 + \dots + X_n$$

## The free case $\varepsilon = 0$

- ▶ The field  $\{\varphi_i\}_{1 \leq i \leq N}$  under  $\mathbb{P}_{0,N}^{\mathbf{P}}$  is distributed like  $\{Z_i\}_{1 \leq i \leq N}$  conditionally on  $(Y_N, Z_N) = (0, 0)$ . ( $Z_{N/2} \approx N^{3/2}$ )
- ▶ Under  $\mathbb{P}_{0,N}^{\mathbf{W}}$  the same, under the further conditioning  $\{Z_1 \geq 0, \dots, Z_N \geq 0\}$

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

**Auxiliary chain:**  $J_i := \varphi_{\tau_i - 1}$  for  $i \in \mathbb{N}$

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

**Auxiliary chain:**  $J_i := \varphi_{\tau_i - 1}$  for  $i \in \mathbb{N}$

Conditionally on  $\tau$  and  $J$ , the excursions  $\{e_i(\cdot)\}_{i \in \mathbb{N}}$  under  $\mathbb{P}_{\varepsilon, N}^a$  are independent and distributed:



# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

**Auxiliary chain:**  $J_i := \varphi_{\tau_i - 1}$  for  $i \in \mathbb{N}$

Conditionally on  $\tau$  and  $J$ , the excursions  $\{e_i(\cdot)\}_{i \in \mathbb{N}}$  under  $\mathbb{P}_{\varepsilon, N}^a$  are independent and distributed:

- ▶ for  $a = p$  like bridges of the process  $\{Z_i\}_i$

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

**Auxiliary chain:**  $J_i := \varphi_{\tau_{i-1}}$  for  $i \in \mathbb{N}$

Conditionally on  $\tau$  and  $J$ , the excursions  $\{e_i(\cdot)\}_{i \in \mathbb{N}}$  under  $\mathbb{P}_{\varepsilon, N}^a$  are independent and distributed:

- ▶ for  $a = p$  like bridges of the process  $\{Z_i\}_i$
- ▶ for  $a = w$  like bridges of the process  $\{Z_i\}_i$  conditioned to stay non-negative

# Excursions and contact set

What happens for  $\varepsilon > 0$ ? The same holds for the excursions.

**Contact set:**  $\tau := \{i \in \mathbb{N} : \varphi_i = 0\} = \{\tau_i\}_{i \geq 0}$

**Excursions:**  $\{e_i(k)\}_k := \{\varphi_{\tau_{i-1}+k}\}_{0 \leq k \leq \tau_i - \tau_{i-1}}$  for  $i \in \mathbb{N}$

**Auxiliary chain:**  $J_i := \varphi_{\tau_i - 1}$  for  $i \in \mathbb{N}$

Conditionally on  $\tau$  and  $J$ , the excursions  $\{e_i(\cdot)\}_{i \in \mathbb{N}}$  under  $\mathbb{P}_{\varepsilon, N}^a$  are independent and distributed:

- ▶ for  $a = p$  like bridges of the process  $\{Z_i\}_i$
- ▶ for  $a = w$  like bridges of the process  $\{Z_i\}_i$  conditioned to stay non-negative

Once we know  $\tau, J$ , the **whole field**  $\{\varphi_i\}_i$  is **reconstructed** by pasting independent excursions from  $\{Z_i\}_i$  (cond. to stay  $\geq 0$ )

# The law of the excursions

Pinning case: good control (Donsker's inv. pr. + LLT)

$$\left\{ \frac{Z_{\langle Nt \rangle}}{N^{3/2}} \right\}_{t \in [0,1]} \text{ condit. on } (Y_N, Z_N) = (0, 0) \implies \left\{ \widehat{l}_t \right\}_{t \in [0,1]}$$

# The law of the excursions

**Pinning case:** good control (Donsker's inv. pr. + LLT)

$$\left\{ \frac{Z_{\langle Nt \rangle}}{N^{3/2}} \right\}_{t \in [0,1]} \text{ condit. on } (Y_N, Z_N) = (0, 0) \implies \{\widehat{l}_t\}_{t \in [0,1]}$$

**Wetting case:** several open issues

- **Integrated BM conditioned to stay non-negative** is studied, but not its bridge

# The law of the excursions

**Pinning case:** good control (Donsker's inv. pr. + LLT)

$$\left\{ \frac{Z_{\langle Nt \rangle}}{N^{3/2}} \right\}_{t \in [0,1]} \text{ condit. on } (Y_N, Z_N) = (0, 0) \implies \left\{ \widehat{l}_t \right\}_{t \in [0,1]}$$

**Wetting case:** several open issues

- ▶ **Integrated BM conditioned to stay non-negative** is studied, but not its bridge
- ▶ Invariance principle seems in any case very difficult

# The law of the excursions

**Pinning case:** good control (Donsker's inv. pr. + LLT)

$$\left\{ \frac{Z_{\lfloor Nt \rfloor}}{N^{3/2}} \right\}_{t \in [0,1]} \text{ condit. on } (Y_N, Z_N) = (0,0) \implies \{\widehat{l}_t\}_{t \in [0,1]}$$

**Wetting case:** several open issues

- ▶ **Integrated BM conditioned to stay non-negative** is studied, but not its bridge
- ▶ Invariance principle seems in any case very difficult
- ▶ **Entropic repulsion:**

$$\mathbf{P}(Z_1 \geq 0, \dots, Z_N \geq 0) \approx ?$$

$$\mathbf{P}(Z_1 \geq 0, \dots, Z_N \geq 0 \mid Y_N = 0, Z_N = 0) \approx ?$$

# The law of the excursions

**Pinning case:** good control (Donsker's inv. pr. + LLT)

$$\left\{ \frac{Z_{\lfloor Nt \rfloor}}{N^{3/2}} \right\}_{t \in [0,1]} \text{ condit. on } (Y_N, Z_N) = (0,0) \implies \{\widehat{l}_t\}_{t \in [0,1]}$$

**Wetting case:** several open issues

- ▶ **Integrated BM conditioned to stay non-negative** is studied, but not its bridge
- ▶ Invariance principle seems in any case very difficult
- ▶ **Entropic repulsion:**

$$\mathbf{P}(Z_1 \geq 0, \dots, Z_N \geq 0) \approx N^{-1/4} \quad [\text{Sinai (SRW)}]$$

$$\mathbf{P}(Z_1 \geq 0, \dots, Z_N \geq 0 \mid Y_N = 0, Z_N = 0) \approx N^{-1/2} \quad [\text{conj.}]$$



# Markov renewal processes

Given a (sub-)probability kernel  $K_{x,dy}(n)$ :

$$\int_{y \in \mathbb{R}} \sum_{n \in \mathbb{N}} K_{x,dy}(n) = c \leq 1, \quad \forall x \in \mathbb{R}$$

we build the Markov renewal process  $\tau$  with modulating chain  $J$ :

$$\mathcal{P}\left(\tau_{i+1} - \tau_i = n, J_{i+1} \in dy \mid J_i = x\right) := K_{x,dy}(n)$$

# Markov renewal processes

Given a (sub-)probability kernel  $K_{x,dy}(n)$ :

$$\int_{y \in \mathbb{R}} \sum_{n \in \mathbb{N}} K_{x,dy}(n) = c \leq 1, \quad \forall x \in \mathbb{R}$$

we build the Markov renewal process  $\tau$  with modulating chain  $J$ :

$$\mathcal{P}(\tau_{i+1} - \tau_i = n, J_{i+1} \in dy \mid J_i = x) := K_{x,dy}(n)$$

The law of  $(\tau, J)$  conditionally on  $\{N, N+1\} \subseteq \tau$  is

$$\mathcal{P}(\tau_i = t_i, J_i \in dy_i \mid \{N, N+1\} \subseteq \tau) = \frac{1}{C_N} \prod_i K_{y_{i-1}, dy_i}(t_i - t_{i-1})$$

with  $C_N = \mathcal{P}(\{N, N+1\} \subseteq \tau)$ .

# The law of the contact set

Consider the following kernels: for  $n \in \mathbb{N}$  and  $x, y \in \mathbb{R}$

$$G_{x,dy}^p(n) := \varepsilon \frac{\mathbf{P}_x(Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

$$G_{x,dy}^w(n) := \varepsilon \frac{\mathbf{P}_x(Z_i \geq 0, i \leq n, Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

# The law of the contact set

Consider the following kernels: for  $n \in \mathbb{N}$  and  $x, y \in \mathbb{R}$

$$G_{x,dy}^p(n) := \varepsilon \frac{\mathbf{P}_x(Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

$$G_{x,dy}^w(n) := \varepsilon \frac{\mathbf{P}_x(Z_i \geq 0, i \leq n, Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

The law of  $(\tau, J)$  is given by

$$\mathbb{P}_{\varepsilon, N}^a(\tau_i = t_i, J_i \in dy_i, i \leq k) = \frac{1}{Z_{\varepsilon, N}^a} \prod_{i=1}^k G_{y_{i-1}, dy_i}^a(t_i - t_{i-1})$$

# The law of the contact set

Consider the following kernels: for  $n \in \mathbb{N}$  and  $x, y \in \mathbb{R}$

$$G_{x,dy}^p(n) := \varepsilon \frac{\mathbf{P}_x(Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

$$G_{x,dy}^w(n) := \varepsilon \frac{\mathbf{P}_x(Z_i \geq 0, i \leq n, Z_{n-1} \in dy, Z_n \in dz)}{dz} \Big|_{z=0}$$

The law of  $(\tau, J)$  is given by

$$\mathbb{P}_{\varepsilon, N}^a(\tau_i = t_i, J_i \in dy_i, i \leq k) = \frac{1}{Z_{\varepsilon, N}^a} \prod_{i=1}^k G_{y_{i-1}, dy_i}^a(t_i - t_{i-1})$$

Reminds of [Markov renewal theory](#)

# Markov renewal processes

We exploit the **invariance properties**: for every  $F$ ,  $v(y)$

$$\begin{aligned} & \mathbb{P}_{\varepsilon, N}^a (\tau_i = t_i, J_i \in dy_i, i \leq k) \\ &= \frac{e^{FN}}{\mathcal{Z}_{\varepsilon, N}^a} \prod_{i=1}^k G_{y_{i-1}, dy_i}^a(t_i - t_{i-1}) e^{-F(t_i - t_{i-1})} \frac{v(y_i)}{v(y_{i-1})} \end{aligned}$$

# Markov renewal processes

We exploit the **invariance properties**: for every  $F$ ,  $v(y)$

$$\begin{aligned} \mathbb{P}_{\varepsilon, N}^a (\tau_i = t_i, J_i \in dy_i, i \leq k) \\ = \frac{e^{FN}}{\mathcal{Z}_{\varepsilon, N}^a} \prod_{i=1}^k G_{y_{i-1}, dy_i}^a(t_i - t_{i-1}) e^{-F(t_i - t_{i-1})} \frac{v(y_i)}{v(y_{i-1})} \end{aligned}$$

If we determine  $F$ ,  $v(\cdot)$  such that

$$K_{x, dy}(n) := G_{x, dy}^a(n) e^{-F \cdot n} \frac{v(y)}{v(x)}$$

is a **probability kernel**, we have the crucial relation

$$\mathbb{P}_{\varepsilon, N}^a (\tau_i = t_i, J_i \in dy_i) = \mathcal{P}(\tau_i = t_i, J_i \in dy_i \mid \{N, N+1\} \subseteq \tau)$$

# A Perron-Frobenius problem

It turns out that:

- $F$  is the solution of the equation

$$\text{spectral radius of } \left( \sum_{n \in \mathbb{N}} G_{x,dy}^a(n) e^{-F \cdot n} \right)_{x,y} = 1$$

when such a solution exists and  $F = 0$  otherwise.



# A Perron-Frobenius problem

It turns out that:

- ▶  $F$  is the solution of the equation

$$\text{spectral radius of } \left( \sum_{n \in \mathbb{N}} G_{x,dy}^a(n) e^{-F \cdot n} \right)_{x,y} = 1$$

when such a solution exists and  $F = 0$  otherwise.

- ▶ In fact  $F = F^a(\varepsilon)$  is the **free energy**

# A Perron-Frobenius problem

It turns out that:

- $F$  is the solution of the equation

$$\text{spectral radius of } \left( \sum_{n \in \mathbb{N}} G_{x,dy}^a(n) e^{-F \cdot n} \right)_{x,y} = 1$$

when such a solution exists and  $F = 0$  otherwise.

- In fact  $F = F^a(\varepsilon)$  is the **free energy**
- $v(\cdot)$  is the **principal eigenfunction**:

$$\int_{y \in \mathbb{R}} \left( \sum_{n \in \mathbb{N}} G_{x,dy}^a(n) e^{-F \cdot n} \right)_{x,y} v(y) = v(x)$$