

# Assignment 1

Submission Deadline: **30 September, 2025** at 23:59

Course Website: <https://ti.inf.ethz.ch/ew/courses/LA25/index.html>

## Exercises

You can get feedback from your TA for Exercise 2 by handing in your solution as pdf via Moodle before the deadline.

### 1. Lines in $\mathbb{R}^m$ (in-class) (★★☆)

- a) Let  $\mathbf{0} \in \mathbb{R}^m$  denote the vector whose entries are all zero. We say that a set  $L$  is a line through the origin in  $\mathbb{R}^m$  if and only if there exists  $\mathbf{w} \in \mathbb{R}^m$  with  $\mathbf{w} \neq \mathbf{0}$  such that  $L = \{\lambda \mathbf{w} : \lambda \in \mathbb{R}\}$ . Let now  $L$  be a line through the origin in  $\mathbb{R}^m$  and let  $\mathbf{u}$  be an arbitrary non-zero element of  $L$ . Prove that  $L = \{\lambda \mathbf{u} : \lambda \in \mathbb{R}\}$ .
- b) For two lines through the origin  $L_1$  and  $L_2$  in  $\mathbb{R}^m$ , prove that we have either  $L_1 \cap L_2 = \{\mathbf{0}\}$  or  $L_1 \cap L_2 = L_1 = L_2$ .
- c) Consider an arbitrary line through the origin  $L$  in  $\mathbb{R}^2$ . Prove that  $L$  is a hyperplane, i.e. find a vector  $\mathbf{d} \neq \mathbf{0}$  such that

$$L = \{\mathbf{v} \in \mathbb{R}^2 : \mathbf{v} \cdot \mathbf{d} = 0\}.$$

### 2. Orthogonality and Linear independence (hand-in) (★☆☆)

- a) For which number  $s \in \mathbb{R}$  are the two following vectors orthogonal?

$$\mathbf{v} = \begin{pmatrix} s \\ 3 \\ 2 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ s \end{pmatrix}$$

- b) For which number  $t \in \mathbb{R}$  are the three following vectors linearly dependent?

$$\mathbf{u} = \begin{pmatrix} t \\ t \\ 0 \end{pmatrix}, \mathbf{v} = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \mathbf{w} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

- c) Show that if two nonzero vectors  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  are orthogonal, then they are linearly independent. Prove that the converse is not necessarily true, i.e. provide an example of two linearly independent vectors that are not orthogonal.

### 3. Cauchy-Schwarz inequality (★★☆)

Consider an arbitrary vector  $\mathbf{v} \in \mathbb{R}^m$ .

- a) Prove the inequality

$$\sum_{i=1}^m v_i \leq \sqrt{m} \|\mathbf{v}\|.$$

b) Prove the inequality

$$\sum_{i=1}^m \sqrt{i} v_i \leq m \| \mathbf{v} \|.$$

#### 4. Linear independence (★★☆)

Let  $\mathbf{e}_1, \dots, \mathbf{e}_m \in \mathbb{R}^m$  be the standard unit vectors in  $\mathbb{R}^m$ . Consider the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_m \in \mathbb{R}^m$  with  $\mathbf{v}_i := \mathbf{e}_i + \mathbf{e}_{i+1}$  for all  $i \in \{1, 2, \dots, m-1\}$  and  $\mathbf{v}_m := \mathbf{e}_m + \mathbf{e}_1$ .

For example, we get

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

in the case  $m = 3$ , and

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \mathbf{v}_4 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

in the case  $m = 4$ .

a) Prove that  $\mathbf{v}_1, \dots, \mathbf{v}_m$  are linearly dependent if  $m$  is even.

b) Prove that  $\mathbf{v}_1, \dots, \mathbf{v}_m$  are linearly independent if  $m$  is odd.

#### 5. Angle between two vectors (★★★)

Consider two non-zero vectors  $\mathbf{v} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$  and  $\mathbf{w} = \begin{pmatrix} z \\ x \\ y \end{pmatrix}$  in  $\mathbb{R}^3$  with  $x + y + z = 0$ . Determine

the value of  $\cos(\alpha)$  where  $\alpha$  denotes the angle between the two vectors  $\mathbf{v}$  and  $\mathbf{w}$ . You are not required to compute (or look up)  $\alpha$ , but you are of course welcome to do so.

#### 6. Challenge 1.6 (★★★)

This exercise asks you to solve Challenge 1.6 from the lecture notes (check it out for some guiding questions).

Let  $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$  and  $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}$  be arbitrary vectors in  $\mathbb{R}^2$  and assume that  $\mathbf{v} \neq \mathbf{0}$  and that  $\mathbf{w} \neq \lambda \mathbf{v}$  for all  $\lambda \in \mathbb{R}$ . Let  $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \in \mathbb{R}^2$  be arbitrary. Prove that  $\mathbf{u}$  can be written as a linear combination of  $\mathbf{v}$  and  $\mathbf{w}$ .