

# Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

Flávio Codeço Coelho and Luiz Max Carvalho

Applied Mathematics School, Fundação Getúlio Vargas

May 1st, 2015

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Summary

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

## Building blocks

- Variable Force of Infection
- Vector dynamics

## Modeling Dengue

- Single-strain model
- Variable Force of Infection

## Parameter estimation

- Estimating  $S_0$
- Attack Ratio

Motivation

Building blocks

- Variable Force of Infection
- Vector dynamics

Modeling Dengue

- Single-strain model
- Variable Force of Infection

Parameter estimation

- Estimating  $S_0$
- Attack Ratio

# Dengue Dynamics

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

- Dengue is a Multi-Strain vector-borne disease

## Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict
- ▶ Vector population dynamics plays a major role in the modulation of incidence

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict
- ▶ Vector population dynamics plays a major role in the modulation of incidence
- ▶ Immunological structure of the population is also a key factor, but is mostly unknown.

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio



# 4 epidemics

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

Building blocks


- Variable Force of Infection
- Vector dynamics

Modeling Dengue


- Single-strain model
- Variable Force of Infection

Parameter estimation


- Estimating  $S_0$
- Attack Ratio




(a) 2010



(b) 2011



(c) 2012



(d) 2013

# Effective Reproductive number( $R_t$ )

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

**Variable Force of Infection**  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

The effective reproductive number can be easily estimated from the incidence time-series,  $Y_t$ :

$$R_t = \left( \frac{Y_{t+1}}{Y_t} \right)^{1/n} \quad (1)$$

Where  $n$  is the ratio between the length of reporting interval and the mean generation time of the disease.

Nishiura et. al. (2010)


But what about the uncertainty about  $R_t$ <sup>1</sup>?

We explore the approach of Ederer and Mantel[4], whose objective is to obtain confidence intervals for the ratio of two Poisson counts. Let  $Y_t \sim \text{Poisson}(\lambda_t)$  and  $Y_{t+1} \sim \text{Poisson}(\lambda_{t+1})$  and define  $S = Y_t + Y_{t+1}$ . The authors note that by conditioning on the sum  $S$

$$Y_{t+1}|S \sim \text{Binomial}(S, \theta_t) \quad (2)$$

$$\theta_t = \frac{\lambda_{t+1}}{\lambda_t + \lambda_{t+1}} \quad (3)$$

---

<sup>1</sup>Coelho, FC and Carvalho, LM (Submitted) 

# $R_t$ 's Uncertainty

Let  $c_\alpha(\theta_t) = \{\theta_t^{(L)}, \theta_t^{(U)}\}$  be such that  $Pr(\theta_t^{(L)} < \theta_t < \theta_t^{(U)}) = \alpha$ . Analogously, define  $c_\alpha(R_t) = \{R_t^{(L)}, R_t^{(U)}\}$  such that  $Pr(R_t^{(L)} < R_t < R_t^{(U)}) = \alpha$ . Ederer and Mantel (1974) [4] show that one can construct a  $100\alpha\%$  confidence interval for  $R_t$  by noting that

$$R_t^{(L)} = \frac{\theta_t^{(L)}}{(1 - \theta_t^{(L)})} \quad \text{and} \quad R_t^{(U)} = \frac{\theta_t^{(U)}}{(1 - \theta_t^{(U)})} \quad (4)$$

# $R_t$ 's Uncertainty

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Taking a Bayesian conjugate distribution approach, If we choose a Beta conjugate prior with parameters  $a_0$  and  $b_0$  for the Binomial likelihood in (2), the posterior distribution for  $\theta_t$  is

$$p(\theta_t | Y_{t+1}, S) \sim \text{Beta}(Y_{t+1} + a_0, Y_t + b_0) \quad (5)$$

Combining equations (4) and (5) tells us that the induced posterior distribution of  $R_t$  is a Beta prime (or inverted Beta) with parameters  $a_1 = Y_{t+1} + a_0$  and  $b_1 = Y_t + b_0$  [?]. The density of the induced distribution is then

$$f_P(R_t | a_1, b_1) = \frac{\Gamma(a_1 + b_1)}{\Gamma(a_1)\Gamma(b_1)} R_t^{a_1-1} (1 + R_t)^{-(a_1+b_1)} \quad (6)$$

Thus, the expectation of  $R_t$  is  $a_1/(b_1 - 1)$  and its variance is  $a_1(a_1 + b_1 - 1)/((b_1 - 2)(b_1 - 1)^2)$ .

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection


Parameter estimation

Estimating  $S_0$   
Attack Ratio

# $R_t$ 's Uncertainty

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho



Motivation

Building blocks

- Variable Force of Infection
- Vector dynamics

Modeling Dengue

- Single-strain model
- Variable Force of Infection

Parameter estimation

- Estimating  $S_0$
- Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio



# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection

Vector dynamics

Modeling Dengue

Single-strain model

Variable Force of Infection

Parameter estimation

Estimating  $S_0$

Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs
- ▶ Effects on mosquito reproduction are non-linear

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs
- ▶ Effects on mosquito reproduction are non-linear
- ▶ Delayed influence

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# $R_t$ vs. Temperature

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks




Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio



# Single Strain SIR

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

Why not multi-strain? No Multi-strain data!!

$$\begin{aligned}\frac{dS}{dt} &= -\beta(t)SI \\ \frac{dI}{dt} &= \beta(t)SI - \tau I \\ \frac{dR}{dt} &= \tau I\end{aligned}\tag{7}$$

where  $S(t) + I(t) + R(t) = 1 \forall t$ .

# Variable Force of Infection

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

From  $R_t$ , we can define a force of infection which varies with time:

$$\beta(t) = \frac{R_t \cdot \tau}{S} \quad (8)$$

But how do we get the value of  $S$ ? we need to estimate  $S_0$ .

Bayesian framework:

- Define priors for  $S_0$  in the range  $(0,1)$

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (9)$$

Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (9)$$



Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model
- ▶ calculate Likelihood of data given current parameterization

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (9)$$


Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model
- ▶ calculate Likelihood of data given current parameterization
- ▶ Determine posterior probability of parameterization

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (9)$$

# Models vs Data

fitting the model to data (Rio de janeiro) to estimate  $S_0^2$ .



Posterior distribution for Susceptible (S) and infectious (I) individuals. Blue dots are data.

<sup>2</sup>Coelho FC et al., 2011

Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

Flávio Codeço Coelho and Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

Once we have  $S_0$ , we can calculate the attack ratio:

$$A_j = \frac{\sum Y_j}{S_{0j}} \quad (10)$$

# Attack ratio

**Table :** Median attack ratio and 95% credibility intervals calculated according to (10). Values are presented as percentage of total population.  $\dagger$ : Year corresponds to the start of the epidemic, however the peak of cases may occur in the following year.  $\ddagger$ : Susceptible fraction. These results show considerable variation in AR between epidemics, consistent with the acquiring and loss of serotype-specific immunity.

Year $\dagger$	median Attack Ratio	$S_0^{\ddagger}$
1996	0.39 (0.17-0.54)	0.00171(0.0012-0.0038)
1997	0.87 (0.74-0.87)	0.00273(0.0027-0.0032)
1998	0.5 (0.49-0.5)	0.00142(0.0014-0.0014)
1999	0.11 (0.037-0.2)	0.00345(0.0018-0.01)
2000	0.25 (0.24-0.27)	0.0155(0.015-0.016)
2001	0.48 (0.47-0.49)	0.0495(0.048-0.051)
2005	0.15 (0.1-0.21)	0.0147(0.01-0.021)
2006	0.11 (0.08-0.14)	0.0281(0.022-0.037)
2007	0.15 (0.15-0.15)	0.135(0.13-0.14)
2008	0.14 (0.031-0.31)	0.00672(0.003-0.024)
2010	0.18 (0.17-0.19)	0.0454(0.043-0.048)
2011	0.086 (0.082-0.094)	0.215(0.2-0.23)
2012	0.14 (0.13-0.15)	0.0621(0.058-0.068)

# Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

## Motivation

### Variable Force of Infection

### Vector dynamics

Single-strain model  
Variable Force of Infection

### Estimating $S_0$

#### Attack Ratio

- A set of small navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and other slide controls.

# Collaboration possibilities

- ▶ Relaxing other simplifying assumptions in classical epidemiological models.
- ▶ Looking into modeling spatio-temporal dynamics taking into account Environmental stock of eggs.
- ▶ Separating noise from relevant variability in observational data, and finding new ways of integrating it into models.

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
**Attack Ratio**

# Thank you!