

# Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

Flávio Codeço Coelho and Luiz Max Carvalho

Escola de Matemática Aplicada, Fundação Getúlio Vargas

11-02-2015

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Summary

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

## Building blocks

- Variable Force of Infection
- Vector dynamics

## Modeling Dengue

- Single-strain model
- Variable Force of Infection

## Parameter estimation

- Estimating  $S_0$
- Attack Ratio

Motivation

Building blocks

- Variable Force of Infection
- Vector dynamics

Modeling Dengue

- Single-strain model
- Variable Force of Infection

Parameter estimation

- Estimating  $S_0$
- Attack Ratio

# Dengue Dynamics

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

- Dengue is a Multi-Strain vector-borne disease

## Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict
- ▶ Vector population dynamics plays a major role in the modulation of incidence

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Dengue Dynamics

- ▶ Dengue is a Multi-Strain vector-borne disease
- ▶ 4 major viral strains in circulation in Brazil
- ▶ Case-notification data is aggregated, i.e., does not discriminate serotype except for a handful of cases.
- ▶ It's a Seasonal disease, but recurrence pattern is hard to predict
- ▶ Vector population dynamics plays a major role in the modulation of incidence
- ▶ Immunological structure of the population is also a key factor, but is mostly unknown.

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks

Variable Force of Infection  
Vector dynamics

### Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

Estimating  $S_0$   
Attack Ratio



# 4 epidemics

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

## Motivation

### Building blocks


- Variable Force of Infection
- Vector dynamics

### Modeling Dengue


- Single-strain model
- Variable Force of Infection

### Parameter estimation


- Estimating  $S_0$
- Attack Ratio




(a) 2010



(b) 2011



(c) 2012



(d) 2013

# Effective Reproductive number( $R_t$ )

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

**Variable Force of Infection**  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

The effective reproductive number can be easily estimated from the incidence time-series,  $Y_t$ :

$$R_t = \left( \frac{Y_{t+1}}{Y_t} \right)^{1/n} \quad (1)$$


Where  $n$  is the ration between the length of reporting interval and the mean generation time of the disease.

Nishiura et. al. (2010)

# $R_t$ 's uncertainty

But what about the uncertainty about  $R_t$  ?

If we assume that the counts  $Y_t$  are Poisson distributed for all  $t$ , we can derive the probability distribution of  $R_t$ <sup>1</sup>



Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

<sup>1</sup>Coelho, FC and Carvalho, LM (Submitted)

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection

Vector dynamics

Modeling Dengue

Single-strain model

Variable Force of Infection

Parameter estimation

Estimating  $S_0$

Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs
- ▶ Effects on mosquito reproduction are non-linear

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Environmental determinants

- ▶ A. Aegypti population dynamics display marked seasonality
- ▶ Temperature, Humidity and rainfall are important factors
- ▶ Environmental stock of eggs
- ▶ Effects on mosquito reproduction are non-linear
- ▶ Delayed influence

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio



# $R_t$ vs. Temperature

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks




Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio



# Single Strain SIR

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

Why not multi-strain? No Multi-strain data!!

$$\begin{aligned}\frac{dS}{dt} &= -\beta(t)SI \\ \frac{dI}{dt} &= \beta(t)SI - \tau I \\ \frac{dR}{dt} &= \tau I\end{aligned}\tag{2}$$

where  $S(t) + I(t) + R(t) = 1 \forall t$ .

# Variable Force of Infection

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

From  $R_t$ , we can define a force of infection which varies with time:

$$\beta(t) = \frac{R_t \cdot \tau}{S} \quad (3)$$

But how do we get the value of  $S$ ? we need to estimate  $S_0$ .

Bayesian framework:

- Define priors for  $S_0$  in the range  $(0,1)$

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (4)$$

Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (4)$$

# Estimating $S_0$

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model
- ▶ calculate Likelihood of data given current parameterization

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (4)$$

# Estimating $S_0$

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio


Bayesian framework:

- ▶ Define priors for  $S_0$  in the range  $(0,1)$
- ▶ Samples from prior, calculate  $\beta(t)$  and run the model
- ▶ calculate Likelihood of data given current parameterization
- ▶ Determine posterior probability of parameterization

$$p(S_{0j}|\mathbf{Y}_j) \propto L(\mathbf{Y}_j|S_{0j}, R_t, m, \tau)\pi(S_{0j}) \quad (4)$$

# Models vs Data

fitting the model to data (Rio de janeiro) to estimate  $S_0^2$ .



Posterior distribution for Susceptible (S) and infectious (I) individuals. Blue dots are data.

<sup>2</sup>Coelho FC et al., 2011

Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

Flávio Codeço Coelho and Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio



Once we have  $S_0$ , we can calculate the attack ratio:

$$A_j = \frac{\sum Y_j}{S_{0j}} \quad (5)$$

# Attack ratio

**Table :** Median attack ratio and 95% credibility intervals calculated according to (5). Values are presented as percentage of total population. <sup>†</sup>: Year corresponds to the start of the epidemic, however the peak of cases may occur in the following year. <sup>‡</sup>: Susceptible fraction. These results show considerable variation in AR between epidemics, consistent with the acquiring and loss of serotype-specific immunity.

Year <sup>†</sup>	median Attack Ratio	$S_0^{\dagger}$
1996	0.39 (0.17-0.54)	0.00171(0.0012-0.0038)
1997	0.87 (0.74-0.87)	0.00273(0.0027-0.0032)
1998	0.5 (0.49-0.5)	0.00142(0.0014-0.0014)
1999	0.11 (0.037-0.2)	0.00345(0.0018-0.01)
2000	0.25 (0.24-0.27)	0.0155(0.015-0.016)
2001	0.48 (0.47-0.49)	0.0495(0.048-0.051)
2005	0.15 (0.1-0.21)	0.0147(0.01-0.021)
2006	0.11 (0.08-0.14)	0.0281(0.022-0.037)
2007	0.15 (0.15-0.15)	0.135(0.13-0.14)
2008	0.14 (0.031-0.31)	0.00672(0.003-0.024)
2010	0.18 (0.17-0.19)	0.0454(0.043-0.048)
2011	0.086 (0.082-0.094)	0.215(0.2-0.23)
2012	0.14 (0.13-0.15)	0.0621(0.058-0.068)

Estimating the Attack Ratio  
of Dengue Epidemics under  
Time-varying Force of  
Infection using Aggregated  
Notification Data

Flávio Codeço Coelho and  
Luiz Max Carvalho

Motivation

Building blocks

Variable Force of Infection  
Vector dynamics

Modeling Dengue

Single-strain model  
Variable Force of Infection

Parameter estimation

Estimating  $S_0$   
Attack Ratio

# Estimating the Attack Ratio of Dengue Epidemics under Time-varying Force of Infection using Aggregated Notification Data

## Motivation

## Building blocks

### Variable Force of Infection

#### Vector dynamics

## Modeling Dengue

Single-strain model  
Variable Force of Infection

### Parameter estimation

### Estimating $S_0$

- ◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ≡ ≡ ↺ 🔍 ↻