

MINE 2020 OF THE YEAR

Mortality and Length of Stay in MIMIC III

Dr. Fani Deligianni,

fani.deligianni@glasgow.ac.uk

Lecturer (Assistant Professor)

nttps://www.gla.ac.uk/schools/computing/staff/fanideligianni Lead of the Computing Technologies for Healthcare Theme

WORLD CHANGING GLASGOW




Mortality


- Most commonly used outcome in prediction studies
- Deceptively simple outcome to define
- Susceptible to selection biases
- Interpretation varies according to the time window



MIMIC-III: ICU Mortality




MIMIC-III: ICU Mortality






MIMIC-III: Hospital Mortality




MIMIC-III: Hospital Mortality






MIMIC-III: Mortality

(PERCENT OF UNIT STAYS)



Total Mortality: 8.5% (4565 patients)


(PERCENT OF UNIT STAYS)




Total Mortality: 11.5% (5748 patients)



MIMIC-III: Hospitalisation Length



MIMIC-III: Hospitalisation Length





Summary

- Mortality and length of stay are common outcomes in prediction models
- They seem easy to define but small differences in their estimation can result in different conclusions
- MIMIC-III provides a useful substrate to understand how to form these queries and extract data

References

- Johnson et al. 'MIMIC-III, a freely accessible critical care database', Scientific Data, 2016.
- modeling and persistence: A comparative review', Journal Gamal et al. 'Standardized electronic health record data of Biomedical Informatics, 2020.