

Multidimensional Random Subset Sum Problem



Francesco d'Amore
COATI team

Based on joint work with L. Becchetti, A. Carvalho Walraven da Cunha, A. Clementi, H. Lesfari, E. Natale, and L. Trevisan

Aalto University
21 September 2022

The Subset Sum problem (SSP)

Input:

- Sequence of n integers x_1, \dots, x_n
- Target value z

The Subset Sum problem (SSP)

Input:

- Sequence of n integers x_1, \dots, x_n
- Target value z

Question:

- Is there a subset $S \subseteq [n]$ such that $\sum_{i \in S} x_i = z$?

The Subset Sum problem (SSP)

Input:

- Sequence of n integers x_1, \dots, x_n
- Target value z

Question:

- Is there a subset $S \subseteq [n]$ such that $\sum_{i \in S} x_i = z$?

Example:

- Sequence: $-9, -7, -1, -1, 0, +3, +4, +5, +9, +11$
- Target value: 2

The Subset Sum problem (SSP)

Input:

- Sequence of n integers x_1, \dots, x_n
- Target value z

Question:

- Is there a subset $S \subseteq [n]$ such that $\sum_{i \in S} x_i = z$?

Example:

- Sequence: $-9, -7, -1, -1, 0, +3, +4, +5, +9, +11$
- Target value: 2
- $-9 + 11 = 2$
- $-7 + 9 = 2$
- $-1 + 3 = 2$
- $-1 - 1 + 4 = 2$
- $-7 + 4 + 5 = 2$
- etc.

Properties of the SSP

- **NP-complete:** reduction from 3-SAT

Properties of the SSP

- **NP-complete:** reduction from 3-SAT
- “**Equivalent**” to the **Number Partition problem** (NPP)
 - Solving the SSP for x_1, \dots, x_n and target $z = \frac{1}{2} \sum_{i \in [n]} x_i$ solves the NPP
 - Solving the NPP for $x_1, \dots, x_n, 2z, \sum_{i \in [n]} x_i$ solves the SSP with tgt z

Properties of the SSP

- **NP-complete:** reduction from 3-SAT
- “Equivalent” to the **Number Partition problem** (NPP)
 - Solving the SSP for x_1, \dots, x_n and target $z = \frac{1}{2} \sum_{i \in [n]} x_i$ solves the NPP
 - Solving the NPP for $x_1, \dots, x_n, 2z, \sum_{i \in [n]} x_i$ solves the SSP with tgt z
- **Common base** for NP-completeness proofs
 - The NPP is among the **six basic NP-complete problems** in the Garey and Johnson's book

Properties of the SSP

- **NP-complete:** reduction from 3-SAT
- “Equivalent” to the **Number Partition problem** (NPP)
 - Solving the SSP for x_1, \dots, x_n and target $z = \frac{1}{2} \sum_{i \in [n]} x_i$ solves the NPP
 - Solving the NPP for $x_1, \dots, x_n, 2z, \sum_{i \in [n]} x_i$ solves the SSP with tgt z
- **Common base** for NP-completeness proofs
 - The NPP is among the **six basic NP-complete problems** in the Garey and Johnson’s book
- **Applications:**
 - combinatorial number theory [Zhi-Wei, 2003]
 - cryptography [Gemmel et Johnston, 2001; Kate et Goldberg, 2011]

The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:


- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z

The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z




The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z



Question:


- How large n for a **subset** $S \subseteq [n]$ to exist, with $z = \sum_{i \in S} X_i$, with h. p.?

The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z
- **Error** parameter $\varepsilon > 0$



Question:

- How large n for a **subset** $S \subseteq [n]$ to exist, with $z = \sum_{i \in S} X_i$, with h. p.?


$$|z - \sum_{i \in S} X_i| \leq 2\varepsilon$$

The Random SSP (RSPP)

- Interested in the **average case**, or in **high probabilities** (h. p.)
- The input values are **random variables**

Input:

- **Sequence** of n independent **random variables** X_1, \dots, X_n
- **Target** value z
- **Error** parameter $\varepsilon > 0$



Question:

- How large n for a **subset** $S \subseteq [n]$ to exist, with $z = \sum_{i \in S} X_i$, with h. p.?

$$|z - \sum_{i \in S} X_i| \leq 2\varepsilon$$

We say that S **2ε -approximates** z

Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP


- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$

Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [[Lueker, 1982](#); [Lueker, 1998](#)]
- Proved in a simpler way in [[Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022](#)]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$




Related literature

- Mainly, results from Lueker [Lueker, 1982; Lueker, 1998]
- Proved in a simpler way in [Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$



Stronger question:


- How large n such that, with h. p., for any $z \in [-1, 1]$ a **subset** $S_z \subseteq [n]$ exists, with $|z - \sum_{i \in S_z} X_i| \leq 2\varepsilon$?
 - **all values** in $[-1, 1]$ must be **2ε -approximated** by **some subset**

Related literature

- Mainly, results from Lueker [Lueker, 1982; Lueker, 1998]
- Proved in a simpler way in [Carvalho, d'Amore, Giroire, Lesfari, Natale, and Viennot, 2022]

Specific instance of RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- **Error** parameter $\varepsilon > 0$



Stronger question:

- How large n such that, with h. p., for any $z \in [-1, 1]$ a **subset** $S_z \subseteq [n]$ exists, with $|z - \sum_{i \in S_z} X_i| \leq 2\varepsilon$?
 - **all values** in $[-1, 1]$ must be **2ε -approximated** by **some subset**

Result for the RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- Small **error** parameter $\varepsilon > 0$

Result for the RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- Small **error** parameter $\varepsilon > 0$

Theorem: [Lueker, 1998]

- There exist two constant $\kappa, C > 0$, such that, if $n \geq C \log \frac{1}{\varepsilon}$, the probability that, for any $z \in [-1, 1]$, a subset $S_z \subseteq [n]$ exists, with $|z - \sum_{i \in S_z} X_i| \leq 2\varepsilon$, is

$$1 - 2 \exp \left[\frac{1}{\kappa n} \left(n - C \log \frac{1}{\varepsilon} \right)^2 \right]$$

Result for the RSSP

- X_1, \dots, X_n **uniform** random variables over $[-1, 1]$
- Small **error** parameter $\varepsilon > 0$

Theorem: [Lueker, 1998]

- There exist two constant $\kappa, C > 0$, such that, if $n \geq C \log \frac{1}{\varepsilon}$, the probability that, for any $z \in [-1, 1]$, a subset $S_z \subseteq [n]$ exists, with $|z - \sum_{i \in S_z} X_i| \leq 2\varepsilon$, is $1 - 2 \exp \left[\frac{1}{\kappa n} \left(n - C \log \frac{1}{\varepsilon} \right)^2 \right]$

Corollaries:

- the result applies to a **wider class of distributions**: any density $f \geq b > 0$ for $x \in [-a, a]$
- **upper bound** on the expectation of
 - the $[a, b]$ -**Subset Sum gap**: minimum value of 2ε such that any real in $[a, b]$ can be 2ε -approximated by some subset S of n **variables**
 - the $[a, b]$ -**Number Partition gap**: minimum value of 2ε such that any real in $[a, b]$ can be 2ε -approximated by using coefficients $\{-1, +1\}$ with n **variables**

Applications of the RSSP

Machine learning:

- Proof of the **Strong Lottery Ticket Hypothesis** [Pensia et al., NeurIPS 2020]
 - *any target network of width d and depth ℓ can be approximated by pruning a random network that is a factor $\mathcal{O}(\log d\ell)$ wider and twice as deep*
 - **feed-forward, fully connected, ReLU activation**
- Related results [Carvalho et al., ICLR 2022; Burkholz et al., ICLR 2022]
- Pruning in Federated Learning [Wang et al., EMNLP 2021]

Proof sketch of the RSSP result


- $z \in \mathbb{R}$, $0 < \varepsilon < 1/3$
- Imagine **revealing** the r.v.s **one by one**: at time t , we have revealed X_1, \dots, X_t

Proof sketch of the RSSP result

- $z \in \mathbb{R}$, $0 < \varepsilon < 1/3$
- Imagine **revealing** the r.v.s **one by one**: at time t , we have revealed X_1, \dots, X_t
- $f_t(z) = 1$ if z is **ε -apx** with a subset of X_1, \dots, X_t **and** $z \in [1, 1]$, 0 otherwise



Proof sketch of the RSSP result

- $z \in \mathbb{R}$, $0 < \varepsilon < 1/3$
- Imagine **revealing** the r.v.s **one by one**: at time t , we have revealed X_1, \dots, X_t
- $f_t(z) = 1$ if z is **ε -apx** with a subset of X_1, \dots, X_t **and** $z \in [1, 1]$, 0 otherwise
- $v_t = \frac{1}{2} \int_{-1}^1 f_t(z) dz$ **ratio** of $[-1, 1]$ that is approximated



Proof sketch of the RSSP result



- $z \in \mathbb{R}$, $0 < \varepsilon < 1/3$
- Imagine **revealing** the r.v.s **one by one**: at time t , we have revealed X_1, \dots, X_t
- $f_t(z) = 1$ if z is **ε -apx** with a subset of X_1, \dots, X_t **and** $z \in [1, 1]$, 0 otherwise
- $v_t = \frac{1}{2} \int_{-1}^1 f_t(z) dz$ **ratio** of $[-1, 1]$ that is approximated



- In **expectation**: v_t **increases exponentially** each round until $1/2$; $w_t = 1 - v_t$ **decreases exponentially** each round until ε

Proof sketch of the RSSP result

- $z \in \mathbb{R}$, $0 < \varepsilon < 1/3$
- Imagine **revealing** the r.v.s **one by one**: at time t , we have revealed X_1, \dots, X_t
- $f_t(z) = 1$ if z is **ε -apx** with a subset of X_1, \dots, X_t **and** $z \in [1, 1]$, 0 otherwise
- $v_t = \frac{1}{2} \int_{-1}^1 f_t(z) dz$ **ratio** of $[-1, 1]$ that is approximated



- In **expectation**: v_t **increases exponentially** each round until $1/2$; $w_t = 1 - v_t$ **decreases exponentially** each round until ε
- A lot of math: convert into **probabilities** and **concentrate**

Multidimensional RSSP (MRSSP)

- Natural generalization

Multidimensional RSSP (MRSSP)

- Natural generalization

Input:


- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$

Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$




Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$



Question:


- How large n for a **subset** $S \subseteq [n]$ to exist, with $\|\mathbf{z} - \sum_{i \in S} X_i\|_\infty \leq 2\varepsilon$, with h. p.?

Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$



Question:

- How large n for a **subset** $S \subseteq [n]$ to exist, with $\|\mathbf{z} - \sum_{i \in S} X_i\|_\infty \leq 2\varepsilon$, with h. p.?

Observations:


- If $X_i \sim \text{Unif}([-1, 1]^d)$, same proof as before leads to $n \geq \exp(d^{\Omega(1)}) \log \frac{1}{\varepsilon}$ variables

Multidimensional RSSP (MRSSP)

- Natural generalization

Input:

- **Sequence** of n independent **random vectors** $X_1, \dots, X_n \in [-1, +1]^d$
- **Target** vector $\mathbf{z} \in [-1, +1]^d$
- **Error** parameter $\varepsilon > 0$



Question:

- How large n for a **subset** $S \subseteq [n]$ to exist, with $\|\mathbf{z} - \sum_{i \in S} X_i\|_\infty \leq 2\varepsilon$, with h. p.?

Observations:

- If $X_i \sim \text{Unif}([-1, 1]^d)$, same proof as before leads to $n \geq \exp(d^{\Omega(1)}) \log \frac{1}{\varepsilon}$ variables
- No success with **method of average bounded differences**, or **Janson's variant of Chernoff bound**

MRSSP: definition of the problem

Input:

- **Sequence** of n independent **standard normal random vectors** $X_1, \dots, X_n \sim \mathcal{N}(\mathbf{0}, I_d)$
 - to better deal with sums
- **Error** parameter $\varepsilon > 0$

MRSSP: definition of the problem

Input:

- **Sequence** of n independent **standard normal random vectors** $X_1, \dots, X_n \sim \mathcal{N}(\mathbf{0}, I_d)$
 - to better deal with sums
- **Error** parameter $\varepsilon > 0$

Question:

- How large n such that, with h. p., for **any** $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}} \subseteq [n]$ exists with $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$?


MRSSP: definition of the problem

Input:

- **Sequence** of n independent **standard normal random vectors** $X_1, \dots, X_n \sim \mathcal{N}(\mathbf{0}, I_d)$
 - to better deal with sums
- **Error** parameter $\varepsilon > 0$

Question:

- How large n such that, with h. p., for **any** $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}} \subseteq [n]$ exists with $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$?



MRSSP: definition of the problem

Input:


- **Sequence** of n independent **standard normal random vectors** $X_1, \dots, X_n \sim \mathcal{N}(\mathbf{0}, I_d)$
 - to better deal with sums
- **Error** parameter $\varepsilon > 0$

Question:

- How large n such that, with h. p., for **any** $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}} \subseteq [n]$ exists with $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$?

Observations:

- $\Theta\left(\frac{1}{\varepsilon^d}\right) = 2^{\Theta(d \log \frac{1}{\varepsilon})}$ **∞ -norm balls** of radius ε are **necessary to cover** the hypercube
 - at least $\Omega\left(d \log \frac{1}{\varepsilon}\right)$ vectors



MRSSP: upper bound in expectation


Negletting constants

- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .

MRSSP: upper bound in expectation

Negletting constants


- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .
- **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes



MRSSP: upper bound in expectation

Negletting constants


- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .
 - **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes
- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets



MRSSP: upper bound in expectation

Negletting constants


- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .
 - **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes
- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets
 - **sum** distributed as $\mathcal{N}(\mathbf{0}, \frac{n}{2}I_d)$



MRSSP: upper bound in expectation

Negletting constants


- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .
 - **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes
- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets
 - **sum** distributed as $\mathcal{N}(\mathbf{0}, \frac{n}{2} I_d)$
 - each **component** has prob. $\frac{\varepsilon}{\sqrt{\frac{n}{2}}}$ to be in $B_\varepsilon(\mathbf{z})$



MRSSP: upper bound in expectation

Negletting constants

- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .
 - **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes
- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets
 - **sum** distributed as $\mathcal{N}(\mathbf{0}, \frac{n}{2} I_d)$
 - each **component** has prob. $\frac{\varepsilon}{\sqrt{\frac{n}{2}}}$ to be in $B_\varepsilon(\mathbf{z})$
 - joint **probability** $\varepsilon^d \left(\frac{n}{2}\right)^{-\frac{d}{2}} = 2^{-d(\log \frac{1}{\varepsilon} + \frac{1}{2} \log \frac{n}{2})}$ to be in $B_\varepsilon(\mathbf{z})$



MRSSP: upper bound in expectation

Negletting constants

- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .

- **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes

- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets


- **sum** distributed as $\mathcal{N}(\mathbf{0}, \frac{n}{2} I_d)$

- each **component** has prob. $\frac{\varepsilon}{\sqrt{\frac{n}{2}}}$ to be in $B_\varepsilon(\mathbf{z})$

- joint **probability** $\varepsilon^d \left(\frac{n}{2}\right)^{-\frac{d}{2}} = 2^{-d(\log \frac{1}{\varepsilon} + \frac{1}{2} \log \frac{n}{2})}$ to be in $B_\varepsilon(\mathbf{z})$

- **Expected number of covering subsets**: $2^{n-o(n)} \cdot 2^{-d(\log \frac{1}{\varepsilon} + \frac{1}{2} \log \frac{n}{2})} > 1$ if

$$n \geq Cd \log \frac{1}{\varepsilon}$$



MRSSP: upper bound in expectation

Negletting constants

- Fix $\mathbf{z} \in [-1, 1]^d$ and a 2ε -side **hypercube** $B_\varepsilon(\mathbf{z})$ around \mathbf{z} — the **∞ -norm ball** of **radius** ε .


- **reminder**: $2^{\Theta(d \log \frac{1}{\varepsilon})}$ such cubes

- Consider **subsets** of **size** $n/2$: $\binom{n}{n/2} \approx 2^{n-o(n)}$ such subsets

- **sum** distributed as $\mathcal{N}(\mathbf{0}, \frac{n}{2} I_d)$

- each **component** has prob. $\frac{\varepsilon}{\sqrt{\frac{n}{2}}}$ to be in $B_\varepsilon(\mathbf{z})$


- joint **probability** $\varepsilon^d \left(\frac{n}{2}\right)^{-\frac{d}{2}} = 2^{-d(\log \frac{1}{\varepsilon} + \frac{1}{2} \log \frac{n}{2})}$ to be in $B_\varepsilon(\mathbf{z})$



- Expected number of covering subsets**: $2^{n-o(n)} \cdot 2^{-d(\log \frac{1}{\varepsilon} + \frac{1}{2} \log \frac{n}{2})} > 1$ if

$$n \geq Cd \log \frac{1}{\varepsilon}$$

- even using $2^{n-o(n)-\Theta(d \log \frac{1}{\varepsilon})}$ subsets **yields** the **same bound**



Our result

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Our result

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$

Our result

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with **exponentially small prob.** in n , for any $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Our result

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with **exponentially small prob.** in n , for any $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Generalizations:

- The **domain** $[-1, 1]^d$ can be **widened** to $[-\sqrt{\frac{n}{d\sqrt{d}}}, \sqrt{\frac{n}{d\sqrt{d}}}]^d$

Our result

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with **exponentially small prob.** in n , for any $\mathbf{z} \in [-1, 1]^d$, a **subset** $S_{\mathbf{z}}$ **exists** such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Generalizations:

- The **domain** $[-1, 1]^d$ can be **widened** to $[-\sqrt{\frac{n}{d\sqrt{d}}}, \sqrt{\frac{n}{d\sqrt{d}}}]^d$
- The **distribution class** is **larger**: every distribution that **contains** a gaussian

Proof sketch


- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$

Proof sketch

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$

Proof sketch


- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$



Proof sketch


- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$

- Fix any **subset** $S \subseteq [n]$ of **size** αn
 - $Y_S = 1$ if $\sum_{i \in S} X_i \sim \mathcal{N}(\mathbf{0}, \alpha n I_d)$ **lies** in $B_\varepsilon(\mathbf{z})$
 - **probability** $Y_S = 1$ is roughly $\frac{(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
- **Gaussian density:** $\frac{1}{\sqrt{(2\pi\sigma)^d}} \exp\left[-\frac{\|x\|_2^2}{2\sigma^d}\right]$



Proof sketch


- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$



- Fix any **subset** $S \subseteq [n]$ of **size** αn
 - $Y_S = 1$ if $\sum_{i \in S} X_i \sim \mathcal{N}(\mathbf{0}, \alpha n I_d)$ **lies** in $B_\varepsilon(\mathbf{z})$
 - **probability** $Y_S = 1$ is roughly $\frac{(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
- **Gaussian density:** $\frac{1}{\sqrt{(2\pi\sigma)^d}} \exp\left[-\frac{\|x\|_2^2}{2\sigma^d}\right]$
- \mathcal{C} set of **subset** $S \subseteq [n]$ of some **fixed size** αn
 - **cardinality** $\binom{n}{\alpha n} \in [\alpha^{\alpha n}, (\alpha e)^{\alpha n}]$

Proof sketch

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$



- Fix any **subset** $S \subseteq [n]$ of **size** αn
- $Y_S = 1$ if $\sum_{i \in S} X_i \sim \mathcal{N}(\mathbf{0}, \alpha n I_d)$ **lies** in $B_\varepsilon(\mathbf{z})$

- **probability** $Y_S = 1$ is roughly $\frac{(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$

- **Gaussian density:** $\frac{1}{\sqrt{(2\pi\sigma)^d}} \exp\left[-\frac{\|x\|_2^2}{2\sigma^d}\right]$


- \mathcal{C} set of **subset** $S \subseteq [n]$ of some **fixed size** αn
- **cardinality** $\binom{n}{\alpha n} \in [\alpha^{\alpha n}, (\alpha e)^{\alpha n}]$

- $Y = \sum_{S \in \mathcal{C}} Y_S$: **number of subsets hitting** $B_\varepsilon(\mathbf{z})$

- $\mathbb{E}[Y] = \sum_{S \in \mathcal{C}} \mathbb{E}[Y_S] = |\mathcal{C}| \cdot \mathbb{P}(Y_S = 1)$ roughly $\frac{|\mathcal{C}|(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$

Proof sketch

- X_1, \dots, X_n **standard normal** random vectors $\sim \mathcal{N}(\mathbf{0}, I_d)$
- **Error** parameter $0 < \varepsilon < 1$
- Fix $\mathbf{z} \in [-1, 1]^d$, and $B_\varepsilon(\mathbf{z})$
- Fix any **subset** $S \subseteq [n]$ of **size** αn
 - $Y_S = 1$ if $\sum_{i \in S} X_i \sim \mathcal{N}(\mathbf{0}, \alpha n I_d)$ **lies** in $B_\varepsilon(\mathbf{z})$
 - **probability** $Y_S = 1$ is roughly $\frac{(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
- **Gaussian density:** $\frac{1}{\sqrt{(2\pi\sigma)^d}} \exp\left[-\frac{\|x\|_2^2}{2\sigma^d}\right]$
- \mathcal{C} set of **subset** $S \subseteq [n]$ of some **fixed size** αn
 - **cardinality** $\binom{n}{\alpha n} \in [\alpha^{\alpha n}, (\alpha e)^{\alpha n}]$
- $Y = \sum_{S \in \mathcal{C}} Y_S$: **number of subsets hitting** $B_\varepsilon(\mathbf{z})$
 - $\mathbb{E}[Y] = \sum_{S \in \mathcal{C}} \mathbb{E}[Y_S] = |\mathcal{C}| \cdot \mathbb{P}(Y_S = 1)$ roughly $\frac{|\mathcal{C}|(2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
 - $n \sim \frac{d}{\alpha \log \alpha} \log \frac{1}{\varepsilon}$ in **expectation**



Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P} \left(Y \geq \frac{\mathbb{E}[Y]}{2} \right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2} \right)^2}$

Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P} \left(Y \geq \frac{\mathbb{E}[Y]}{2} \right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2} \right)^2}$

- $\text{V}[Y] = \text{V} \left[\sum_{S \in \mathcal{C}} Y_S \right] = \sum_{S, T \in \mathcal{C}} \text{Cov} [Y_S, Y_T]$

Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$

- $\text{V}[Y] = \text{V}\left[\sum_{S \in \mathcal{C}} Y_S\right] = \sum_{S, T \in \mathcal{C}} \text{Cov}[Y_S, Y_T]$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{E}[Y_S Y_T] - \mathbb{E}[Y_S] \mathbb{E}[Y_T]$$

Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$

- $\text{V}[Y] = \text{V}\left[\sum_{S \in \mathcal{C}} Y_S\right] = \sum_{S, T \in \mathcal{C}} \text{Cov}[Y_S, Y_T]$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{E}[Y_S Y_T] - \mathbb{E}[Y_S] \mathbb{E}[Y_T]$$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{P}(Y_S = 1, Y_T = 1) - \mathbb{P}(Y_S = 1) \mathbb{P}(Y_T = 1)$$

Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\mathbb{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$

- $\mathbb{V}[Y] = \mathbb{V}\left[\sum_{S \in \mathcal{C}} Y_S\right] = \sum_{S, T \in \mathcal{C}} \text{Cov}[Y_S, Y_T]$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{E}[Y_S Y_T] - \mathbb{E}[Y_S] \mathbb{E}[Y_T]$$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{P}(Y_S = 1, Y_T = 1) - \mathbb{P}(Y_S = 1) \mathbb{P}(Y_T = 1) = (\mathbb{P}(Y_S = 1) [1 - \mathbb{P}(Y_S = 1)])$$

Strategy: problem

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- **Second moment method** to convert in probabilities

- Chebyshev implies $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$


- $\text{V}[Y] = \text{V}\left[\sum_{S \in \mathcal{C}} Y_S\right] = \sum_{S, T \in \mathcal{C}} \text{Cov}[Y_S, Y_T]$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{E}[Y_S Y_T] - \mathbb{E}[Y_S] \mathbb{E}[Y_T]$$

$$= \sum_{S, T \in \mathcal{C}} \mathbb{P}(Y_S = 1, Y_T = 1) - \mathbb{P}(Y_S = 1) \mathbb{P}(Y_T = 1)$$


Main difficulty: counting and dealing with dependencies among subsets

- joint probability of Y_S and Y_T




Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**




Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$




Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$
 - **restrict** \mathcal{C} to sets whose **pairwise intersection size** at most $2\alpha^2 n$



Strategy: solution


- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$
 - **restrict** \mathcal{C} to sets whose **pairwise intersection size** at most $2\alpha^2 n$
 - such a set has **cardinality** **at least** $2^{\frac{\alpha^2 n}{6}}$



Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$
 - **restrict** \mathcal{C} to sets whose **pairwise intersection size** at most $2\alpha^2 n$
 - such a set has **cardinality** **at least** $2^{\frac{\alpha^2 n}{6}}$

- $Y = \sum_{S \in \mathcal{C}} Y_S$: **number of subsets hitting** $B_\varepsilon(\mathbf{z})$
 - $\mathbb{E}[Y]$ roughly **greater** than $\frac{2^{\frac{\alpha^2 n}{6}} (2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$




Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$
 - **restrict** \mathcal{C} to sets whose **pairwise intersection size** at most $2\alpha^2 n$
 - such a set has **cardinality** **at least** $2^{\frac{\alpha^2 n}{6}}$

- $Y = \sum_{S \in \mathcal{C}} Y_S$: **number of subsets hitting** $B_\varepsilon(\mathbf{z})$

- $\mathbb{E}[Y]$ roughly **greater** than $\frac{2^{\frac{\alpha^2 n}{6}} (2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
- **already** $n \geq C \frac{d}{\alpha^2} \log \frac{1}{\varepsilon}$ in **expectation**




Strategy: solution

- \mathcal{C} set of **subset** $S \subseteq [n]$ of **size** αn
 - pick two **elements** from \mathcal{C} **uniformly at random**
 - **expected intersection** $\alpha^2 n$
 - **restrict** \mathcal{C} to sets whose **pairwise intersection size** at most $2\alpha^2 n$
 - such a set has **cardinality** **at least** $2^{\frac{\alpha^2 n}{6}}$

- $Y = \sum_{S \in \mathcal{C}} Y_S$: **number of subsets hitting** $B_\varepsilon(\mathbf{z})$

- $\mathbb{E}[Y]$ roughly **greater** than $\frac{2^{\frac{\alpha^2 n}{6}} (2\varepsilon)^d}{(2\pi\alpha n)^{\frac{d}{2}}}$
- **already** $n \geq C \frac{d}{\alpha^2} \log \frac{1}{\varepsilon}$ in **expectation**



- $V[Y] = \sum_{S, T \in \mathcal{C}} \mathbb{P}(Y_S = 1, Y_T = 1) - \mathbb{P}(Y_S = 1) \mathbb{P}(Y_T = 1)$
 - a lot of math to get **tight bounds**

Chebyshev

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- Chebyshev: $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$
- $\mathbb{P}(Y \geq 1) \geq \frac{1}{3}$ if $\alpha \leq \frac{1}{6\sqrt{d}}$ and $n \geq C \frac{d}{\alpha^2} (\log \frac{1}{\varepsilon} + \log d)$

Chebyshev

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$

- Chebyshev: $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$
- $\mathbb{P}(Y \geq 1) \geq \frac{1}{3}$ if $\alpha \leq \frac{1}{6\sqrt{d}}$ and $n \geq C \frac{d}{\alpha^2} (\log \frac{1}{\varepsilon} + \log d)$
- subset size $\alpha n = \Theta\left(\frac{n}{\sqrt{d}}\right)$

Chebyshev

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$


- Chebyshev: $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$
- $\mathbb{P}(Y \geq 1) \geq \frac{1}{3}$ if $\alpha \leq \frac{1}{6\sqrt{d}}$ and $n \geq C \frac{d}{\alpha^2} (\log \frac{1}{\varepsilon} + \log d)$
- subset size $\alpha n = \Theta\left(\frac{n}{\sqrt{d}}\right)$
- intersection at most $2\alpha^2 n = \Theta\left(\frac{n}{d}\right)$
- bound on n given from estimation of $|\mathcal{C}| \geq 2^{\frac{\alpha^2 n}{6}}$

Chebyshev

$Y = \sum_{S \in \mathcal{C}} Y_S$: number of subsets hitting $B_\varepsilon(\mathbf{z})$


- Chebyshev: $\mathbb{P}\left(Y \geq \frac{\mathbb{E}[Y]}{2}\right) \geq 1 - \frac{\text{V}[Y]}{\left(\frac{\mathbb{E}[Y]}{2}\right)^2}$
- $\mathbb{P}(Y \geq 1) \geq \frac{1}{3}$ if $\alpha \leq \frac{1}{6\sqrt{d}}$ and $n \geq C \frac{d}{\alpha^2} (\log \frac{1}{\varepsilon} + \log d)$
- subset size $\alpha n = \Theta\left(\frac{n}{\sqrt{d}}\right)$
- intersection at most $2\alpha^2 n = \Theta\left(\frac{n}{d}\right)$
- bound on n given from estimation of $|\mathcal{C}| \geq 2^{\frac{\alpha^2 n}{6}}$

- for all hypercubes?



Amplification argument: union bound


- **Amplification**
- $\frac{1}{\epsilon^d}$ smaller **hypercubes**



Amplification argument: union bound

- **Amplification**


- $\frac{1}{\varepsilon^d}$ smaller **hypercubes**
- each has prob. $\frac{1}{3}$ of being **ε -apx** by $n \sim Cd^2(\log \frac{1}{\varepsilon} + \log d)$ **random vectors**



Amplification argument: union bound

- **Amplification**


- $\frac{1}{\epsilon^d}$ smaller **hypercubes**
- each has prob. $\frac{1}{3}$ of being **ϵ -apx** by $n \sim Cd^2(\log \frac{1}{\epsilon} + \log d)$ **random vectors**
- **repeat** k times with **independent** samples




Amplification argument: union bound

- **Amplification**


- $\frac{1}{\varepsilon^d}$ smaller **hypercubes**
- each has prob. $\frac{1}{3}$ of being **ε -apx** by $n \sim Cd^2(\log \frac{1}{\varepsilon} + \log d)$ **random vectors**
- **repeat** k times with **independent** samples




$n_1 \sim n$




$n_2 \sim n$



$n_3 \sim n$



$n_k \sim n$




...


Amplification argument: union bound

- **Amplification**


- $\frac{1}{\epsilon^d}$ smaller **hypercubes**
- each has prob. $\frac{1}{3}$ of being **ϵ -apx** by $n \sim Cd^2(\log \frac{1}{\epsilon} + \log d)$ **random vectors**
- **repeat** k times with **independent** samples




$n_1 \sim n$




$n_2 \sim n$



$n_3 \sim n$



$n_k \sim n$



- **Prob. to fail** $\leq \frac{1}{\epsilon^d} \cdot \left(\frac{2}{3}\right)^k$

- $k \sim d \log \frac{1}{\epsilon}$ to have **exponentially small prob.** to fail

Recap of results

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with exponentially small prob. in n , for any $\mathbf{z} \in [-1, 1]^d$, a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Recap of results

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with exponentially small prob. in n , for any $\mathbf{z} \in [-1, 1]^d$, a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Generalizations:

- The domain $[-1, 1]^d$ can be widened to $[-\sqrt{\frac{n}{d\sqrt{d}}}, \sqrt{\frac{n}{d\sqrt{d}}}]^d$

Recap of results

Theorems:

- If $\mathbf{z} \in [-1, 1]^d$ and $n \geq Cd^2(\log \frac{1}{\varepsilon} + \log d)$ a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq \varepsilon$ w.p. $\frac{1}{3}$
- If $n \geq Cd^3 \log \frac{1}{\varepsilon} \cdot (\log \frac{1}{\varepsilon} + \log d)$, with exponentially small prob. in n , for any $\mathbf{z} \in [-1, 1]^d$, a subset $S_{\mathbf{z}}$ exists such that $\|\mathbf{z} - \sum_{i \in S_{\mathbf{z}}} X_i\|_{\infty} \leq 2\varepsilon$

Generalizations:

- The domain $[-1, 1]^d$ can be widened to $[-\sqrt{\frac{n}{d\sqrt{d}}}, \sqrt{\frac{n}{d\sqrt{d}}}]^d$
- The distribution class is larger: every distribution that contains a gaussian

Wider distribution class

$\phi(x)$ pdf of a standard Gaussian, $p \in (0, 1]$

Wider distribution class


$\phi(x)$ pdf of a standard Gaussian, $p \in (0, 1]$

- Every pdf $f : \mathbb{R} \rightarrow [0, +\infty)$
 - $f(x) = p\phi(x) + (1 - p)g(x)$

Wider distribution class

$\phi(x)$ pdf of a standard Gaussian, $p \in (0, 1]$

- Every pdf $f : \mathbb{R} \rightarrow [0, +\infty)$
 - $f(x) = p\phi(x) + (1 - p)g(x)$




- **Green:** standard Gaussian
- **Red:** resized standard Gaussian
- **Black:** example of $f(x)$

Wider distribution class

$\phi(x)$ pdf of a standard Gaussian, $p \in (0, 1]$

- Every pdf $f : \mathbb{R} \rightarrow [0, +\infty)$
 - $f(x) = p\phi(x) + (1 - p)g(x)$




- **Green:** standard Gaussian
- **Red:** resized standard Gaussian
- **Black:** example of $f(x)$

- **Equivalently:** with prob. p it's a standard Gaussian
 - just amplify more the number of variables
 - only a constant worse

Wider distribution class

$\phi(x)$ pdf of a standard Gaussian, $p \in (0, 1]$

- Every pdf $f : \mathbb{R} \rightarrow [0, +\infty)$
 - $f(x) = p\phi(x) + (1 - p)g(x)$



- **Green:** standard Gaussian
- **Red:** resized standard Gaussian
- **Black:** example of $f(x)$

- **Equivalently:** with prob. p it's a standard Gaussian
 - just amplify more the number of variables
 - only a constant worse
- Idea from [\[Lueker, 1998\]](#)

Work in progress . . .

- **Improve the bound**
 - **lower bound** $n \sim d \log \frac{1}{\varepsilon}$
 - make use of **more subsets**

Work in progress . . .

- **Improve the bound**
 - **lower bound** $n \sim d \log \frac{1}{\varepsilon}$
 - make use of **more subsets**
- **Bad news:** [Borst et al., 2022] has a **huge overlap**
 - application to the **integrality gap** of the **binary integer problem**

Work in progress . . .

- **Improve the bound**
 - **lower bound** $n \sim d \log \frac{1}{\varepsilon}$
 - make use of **more subsets**
- **Bad news:** [Borst et al., 2022] has a **huge overlap**
 - application to the **integrality gap** of the **binary integer problem**
 - similar **subset sum** result
 - **error** must be **exponentially small** in the dimension
 - n always **asymptotically greater** than $d^{2.25}$

Work in progress . . .

- **Improve the bound**
 - **lower bound** $n \sim d \log \frac{1}{\varepsilon}$
 - make use of **more subsets**
- **Bad news:** [Borst et al., 2022] has a **huge overlap**
 - application to the **integrality gap** of the **binary integer problem**
 - similar **subset sum** result
 - **error** must be **exponentially small** in the dimension
 - n always **asymptotically greater** than $d^{2.25}$
- **Even wider class of distributions**
 - if **convergence** to Gaussian is **fast enough**

Work in progress . . .

- **Improve the bound**
 - **lower bound** $n \sim d \log \frac{1}{\varepsilon}$
 - make use of **more subsets**
- **Bad news:** [Borst et al., 2022] has a **huge overlap**
 - application to the **integrality gap** of the **binary integer problem**
 - similar **subset sum** result
 - **error** must be **exponentially small** in the dimension
 - n always **asymptotically greater** than $d^{2.25}$
- **Even wider class of distributions**
 - if **convergence** to Gaussian is **fast enough**
- **Neural network** applications: investigate further **structured pruning**

The end



Thank
You!