

Q1. The ratio of two quantities is dimensionless but not unitless. Which of the following is a valid example?

- A. Angle in radians
- B. Relative density
- C. Temperature ratio in kelvin
- D. Refractive index

Answer: C

Explanation:

The ratio of two temperatures in kelvin is dimensionless because the units cancel, but it is not unitless since kelvin is a base unit.

Q2. The SI unit of the coefficient of viscosity (written as " $\eta$ ") is:

- A. Pa·s
- B. N·s/m
- C.  $\text{kg}\cdot\text{m}^{-1}\cdot\text{s}^{-2}$
- D. dyne·s/cm<sup>2</sup>

Answer: A

Explanation:

$$\begin{aligned}\text{Viscosity } \eta &= (\text{Force} / \text{Area}) \times (dx / dv) \\ &= (\text{N} / \text{m}^2) \times \text{s} = \text{Pa}\cdot\text{s}\end{aligned}$$

Q3. Which of the following pairs has the same dimensions but different SI units?

- A. Work and torque
- B. Pressure and energy
- C. Power and energy
- D. Impulse and force

Answer: A

Explanation:

Work and torque both have the same dimensions:  $[M L^2 T^{-2}]$

Work has SI unit: joule ( $J = \text{N}\cdot\text{m}$ )

Torque also has unit  $\text{N}\cdot\text{m}$ , but represents rotational effect and is a vector  
So, dimensions are same, but physical meaning and usage are different.

Q4. Which of the following physical quantities has an SI unit that is not derived, i.e., it is a base unit?

- A. Luminous intensity
- B. Electric potential
- C. Heat
- D. Velocity

Answer: A

Explanation:

Luminous intensity is a base physical quantity. Its SI unit is candela (cd), which is a base unit.

Q5. The SI unit of solid angle is:

- A. Radian
- B. Steradian
- C. Degree
- D. Dioptre

Answer: B

Explanation:

Solid angle is measured in steradians (sr), the SI unit for 3D angular measure.

Q6. A student writes the unit of Planck's constant as "erg·s". What is the equivalent SI unit?

- A. J·s
- B. N·s
- C.  $\text{kg}\cdot\text{m}^2/\text{s}$
- D. W·s

Answer: A

Explanation:

$1 \text{ erg} = 10^{-7} \text{ joule}$

So,  $\text{erg}\cdot\text{s} = 10^{-7} \text{ J}\cdot\text{s}$

Hence, the correct SI unit of Planck's constant is joule-second (J·s)

Q7. The dimensional formula of pressure is:

- A.  $[M L^{-1} T^{-2}]$
- B.  $[M L T^{-2}]$
- C.  $[M L^2 T^{-2}]$
- D.  $[M L^{-2} T^{-2}]$

Answer: D

Explanation:

Pressure = Force / Area

= (Mass  $\times$  Acceleration) / Area

=  $[M L T^{-2}] / [L^2] = [M L^{-1} T^{-2}]$

Q8. Which of the following is dimensionally consistent?

- A.  $v^2 = u^2 + 2as$
- B.  $v = u + at^2$
- C.  $v^2 = u^2 + 2a$
- D.  $s = ut + at$

Answer: A

Explanation:

Check each term's dimensions:

$$v^2 = [L^2 T^{-2}]$$

$$u^2 = [L^2 T^{-2}]$$

$$2as = [L T^{-2}] \times [L] = [L^2 T^{-2}]$$

All terms have same dimensions.

Q9. The dimensional formula of surface tension is:

- A.  $[M^0 L^0 T^0]$
- B.  $[M T^{-2}]$
- C.  $[M L^0 T^{-2}]$
- D.  $[M T]$

Answer: C

Explanation:

Surface tension = Force / Length

=  $[M L T^{-2}] / [L] = [M L^0 T^{-2}]$

Q10. Which of the following physical quantities has the same dimensional formula as energy?

- A. Force
- B. Torque
- C. Pressure
- D. Power

Answer: B

Explanation:

$$\text{Energy} = [M L^2 T^{-2}]$$

$$\text{Torque} = \text{Force} \times \text{Distance} = [M L T^{-2}] \times [L] = [M L^2 T^{-2}]$$

Q11. Which quantity has the same dimensions as impulse?

- A. Force
- B. Energy
- C. Momentum
- D. Pressure

Answer: C

Explanation:

$$\text{Impulse} = \text{Force} \times \text{Time} = [M L T^{-2}] \times [T] = [M L T^{-1}]$$

$$\text{Momentum} = \text{Mass} \times \text{Velocity} = [M] \times [L T^{-1}] = [M L T^{-1}]$$

Q12. Which of the following equations is dimensionally incorrect?

- A.  $T = 2\pi \sqrt{L/g}$
- B.  $E = mc^2$
- C.  $F = ma^2$
- D.  $v = u + at$

Answer: C

Explanation:

$$F = ma^2 \Rightarrow \text{Dimensions} = M \times (L T^{-2})^2 = [M L^2 T^{-4}]$$

But Force =  $[M L T^{-2}]$ , so mismatch  $\rightarrow$  Incorrect.

Q13. What is the dimensional formula of Planck's constant?

- A.  $[M L^2 T^{-1}]$
- B.  $[M L^2 T^{-2}]$
- C.  $[M L T^{-1}]$
- D.  $[M L T]$

Answer: A

Explanation:

Energy =  $h \times \text{frequency} \rightarrow [M L^2 T^{-2}] = h \times [T^{-1}]$

So,  $h = [M L^2 T^{-1}]$

Q14. The dimensional formula of angular momentum is:

- A.  $[M L^2 T^{-2}]$
- B.  $[M L^2 T^{-1}]$
- C.  $[M L T^{-1}]$
- D.  $[M^2 L T^{-2}]$

Answer: B

Explanation:

Angular momentum = Moment of inertia  $\times$  Angular velocity

$= [M L^2] \times [T^{-1}] = [M L^2 T^{-1}]$

Q15. Which quantity is dimensionless?

- A. Strain
- B. Stress
- C. Energy
- D. Force

Answer: A

Explanation:

Strain = Change in length / Original length  $\rightarrow [L] / [L] = \text{No units}$

Hence, dimensionless.

Q16. The unit of a quantity is given as joule-second. Identify the physical quantity.

- A. Planck's constant
- B. Work

- C. Energy
- D. Angular momentum

Answer: D

Explanation:

Angular momentum = Moment of inertia  $\times$  Angular velocity  
 $= [M L^2] \times [T^{-1}] = [M L^2 T^{-1}] \rightarrow \text{Joule-second}$

Q17. Which of the following quantities has the dimensional formula  $[M^0 L^0 T^0]$ ?

- A. Relative density
- B. Density
- C. Pressure
- D. Force

Answer: A

Explanation:

Relative density = Density of substance / Density of water  
 $\rightarrow$  No units  $\rightarrow$  Dimensionless

Q18. Dimensional formula of gravitational constant (G) is:

- A.  $[M^{-1} L^3 T^{-2}]$
- B.  $[M L^3 T^{-2}]$
- C.  $[M^{-2} L^3 T^{-2}]$
- D.  $[M^{-1} L^2 T^2]$

Answer: A

Explanation:

$F = G (m_1 m_2 / r^2) \Rightarrow G = F \times r^2 / m^2$   
 $= [M L T^{-2}] \times [L^2] / [M^2] = [M^{-1} L^3 T^{-2}]$

Q19. Which of the following quantities has same dimensions as work?

- A. Moment of inertia
- B. Power
- C. Torque
- D. Momentum

Answer: C

Explanation:

Torque = Force  $\times$  Perpendicular distance

$$= [M L T^{-2}] \times [L] = [M L^2 T^{-2}] \rightarrow \text{Same as Work}$$

Q20. The true value of a quantity is  $9.80 \text{ m/s}^2$ , and the measured value is  $9.65 \text{ m/s}^2$ . The absolute error is:

- A. 0.05
- B. 0.10
- C. 0.15
- D. 0.25

Answer: C

Explanation:

$$\text{Absolute error} = |\text{True} - \text{Measured}| = |9.80 - 9.65| = 0.15$$

Q21. If the absolute error in measuring length is  $0.02 \text{ m}$  and the length is  $2.00 \text{ m}$ , then the percentage error is:

- A. 1%
- B. 0.5%
- C. 0.02%
- D. 2%

Answer: B

Explanation:

$$\text{Percentage error} = (0.02 / 2.00) \times 100 = 1\%$$

Q22. Which of the following has maximum possible error when added?

- A.  $10.2 + 3.58$
- B.  $100.03 + 0.004$
- C.  $23.1 + 5.342$
- D.  $3.2 + 1.75$

Answer: A

Explanation:

Final result should match the least precise decimal place (1 decimal). Max error comes from large rounding.

Q23. If two quantities  $A = 5.0 \pm 0.1$  and  $B = 2.0 \pm 0.2$ , then the error in  $A + B$  is:

- A.  $\pm 0.1$
- B.  $\pm 0.2$
- C.  $\pm 0.3$
- D.  $\pm 0.02$

Answer: C

Explanation:

In addition: total error = error in A + error in B =  $0.1 + 0.2 = \pm 0.3$

Q24. If  $A = 10 \pm 0.2$  and  $B = 5 \pm 0.1$ , the maximum relative error in  $A/B$  is:

- A. 0.02
- B. 0.03
- C. 0.04
- D. 0.05

Answer: D

Explanation:

Relative error =  $(\Delta A/A) + (\Delta B/B) = (0.2/10) + (0.1/5) = 0.02 + 0.02 = 0.04$  (or 4%)

Q25. If  $A = 4.0 \pm 0.2$  and we calculate  $A^2$ , the percentage error is:

- A. 5%
- B. 10%
- C. 20%
- D. 40%

Answer: B

Explanation:

If  $y = A^n$ , % error =  $n \times (\% \text{ error in } A)$

% error in A =  $(0.2/4.0) \times 100 = 5\%$

So, % error in  $A^2 = 2 \times 5 = 10\%$

Q26. The least count of a scale that reads up to 1 mm is:

- A. 0.1 mm
- B. 0.5 mm



- C. 1 mm
- D. 10 mm

Answer: C

Explanation:

Least count = smallest division readable = 1 mm

Q27. A student measures the diameter of a wire as 2.00 mm, 2.02 mm, and 1.98 mm. The mean diameter is:

- A. 2.01 mm
- B. 2.00 mm
- C. 1.99 mm
- D. 2.02 mm

Answer: B

Explanation:

Mean =  $(2.00 + 2.02 + 1.98)/3 = 2.00$  mm

Q28. For the same readings in Q27, the absolute error in each measurement is:

- A. 0.01 mm
- B. 0.02 mm
- C. 0.04 mm
- D. 0.10 mm

Answer: B

Explanation:

$|2.02 - 2.00| = 0.02$ ,  $|1.98 - 2.00| = 0.02$

Average error =  $(0.02 + 0.02 + 0)/3 = 0.013 \approx 0.02$  mm

Q29. If  $Z = A \times B$ , then fractional error in Z is:

- A.  $\Delta A + \Delta B$
- B.  $\Delta A/A + \Delta B/B$
- C.  $A \times \Delta B + B \times \Delta A$
- D. None

Answer: B

Explanation:

For multiplication/division: fractional error in  $Z = \Delta A/A + \Delta B/B$

Q30. If a measurement is 2.50 cm, how many significant figures does it have?

- A. 1
- B. 2
- C. 3
- D. 4

Answer: C

Explanation:

Trailing zero after decimal counts  $\rightarrow$  2.50 has 3 significant figures

Q31. The quantity 0.00340 has how many significant figures?

- A. 2
- B. 3
- C. 4
- D. 5

Answer: B

Explanation:

Leading zeros don't count. 3, 4, and final 0 after decimal  $\rightarrow$  3 sig figs

Q32. Which of the following is dimensionally incorrect?

- A. Velocity = Distance / Time
- B. Work = Force  $\times$  Distance
- C. Power = Work  $\times$  Time
- D. Acceleration = Velocity / Time

Answer: C

Explanation:

Power = Work / Time, not  $\times$  Time  $\Rightarrow$  Incorrect dimensions.

Q33. If length =  $4.0 \pm 0.2$  m and time =  $2.0 \pm 0.1$  s, what is the error in speed?

- A. 0.1 m/s

- B. 0.2 m/s
- C. 0.3 m/s
- D. 0.4 m/s

Answer: B

Explanation:

Speed =  $L/T$

Relative error =  $0.2/4.0 + 0.1/2.0 = 0.05 + 0.05 = 0.10$

Speed = 2.0 m/s

Absolute error = 10% of 2.0 = 0.2 m/s

Q34. If density is calculated from  $m/V$ , and  $m$  has 3% error,  $V$  has 2% error, then error in density is:

- A. 1%
- B. 2%
- C. 5%
- D. 6%

Answer: C

Explanation:

% error in density = 3% + 2% = 5%

Q35. If  $R = A^2B / C^3$ , the % error in  $R$  is:

- A.  $2(\Delta A/A) + \Delta B/B + 3(\Delta C/C)$
- B.  $\Delta A + \Delta B - \Delta C$
- C.  $\Delta A^2 + \Delta B^2 - \Delta C^3$
- D.  $\Delta A/A + \Delta B/B - \Delta C/C$

Answer: A

Explanation:

Use exponents: % error =  $n(\Delta A/A) + m(\Delta B/B) + \dots$

So here =  $2(\Delta A/A) + \Delta B/B + 3(\Delta C/C)$

Q36. A value is measured as 5.60. Rounding it to 2 significant digits gives:

- A. 5.5
- B. 5.6
- C. 6.0

D. 5.0

Answer: B

Explanation:

First two digits: 5.6 → Already 2 significant figures

Q37. The mean absolute error is best described as:

- A. Minimum possible error
- B. Average of all errors
- C. Largest error
- D. Standard deviation

Answer: B

Explanation:

Mean absolute error = average of individual absolute errors.

Q38. A physical quantity X is found using:

$$X = P^2Q / R.$$

If % errors in P, Q and R are 2%, 1%, and 3% respectively, then % error in X is:

- A. 8%
- B. 7%
- C. 6%
- D. 5%

Answer: C

Explanation:

$$\% \text{ error in } X = 2(2\%) + 1\% + 3\% = 6\%$$

Q39. Which type of error cannot be reduced by taking multiple measurements?

- A. Random error
- B. Instrumental error
- C. Systematic error
- D. Personal error

Answer: C

Explanation:

Systematic error stays constant in every trial → not reduced by repetition.

Q40. Which device gives least count of 0.01 cm?

- A. Meter scale
- B. Vernier caliper
- C. Screw gauge
- D. Measuring tape

Answer: B

Explanation:

Vernier caliper → typical least count = 0.01 cm

Q41. The resistance of a wire is measured as  $R = 5.00 \pm 0.05 \Omega$ . The percentage error in R is:

- A. 1%
- B. 5%
- C. 10%
- D. 0.5%

Answer: A

Explanation:

Percentage error =  $(0.05 / 5.00) \times 100 = 1\%$

Q42. The quantity Q is given by  $Q = A^2 / \sqrt{B}$ . If the percentage errors in A and B are 3% and 4% respectively, then the percentage error in Q is:

- A. 4%
- B. 5%
- C. 8%
- D. 10%

Answer: C

Explanation:

% error in Q =  $2 \times \text{error in A} + (1/2) \times \text{error in B}$   
 $= 2 \times 3 + 0.5 \times 4 = 6 + 2 = 8\%$

Q43. A student records the time of oscillation of a pendulum as 1.62 s using a stopwatch with least count 0.01 s. What is the relative error?

- A. 0.62%
- B. 1%
- C. 0.31%
- D. 0.01%

Answer: A

Explanation:

$$\text{Relative error} = (0.01 / 1.62) \times 100 \approx 0.62\%$$

Q44. A student takes three readings for the diameter of a wire: 2.01 mm, 2.03 mm, and 2.00 mm. The mean and absolute error are:

- A. 2.01 mm and  $\pm 0.02$  mm
- B. 2.01 mm and  $\pm 0.01$  mm
- C. 2.02 mm and  $\pm 0.03$  mm
- D. 2.02 mm and  $\pm 0.01$  mm

Answer: A

Explanation:

$$\text{Mean} = (2.01 + 2.03 + 2.00)/3 = 2.01 \text{ mm}$$

$$\text{Deviation from mean: } 0.00, 0.02, 0.01$$

$$\text{Mean absolute error} = (0 + 0.02 + 0.01)/3 \approx 0.01$$

$$\text{Maximum absolute error} = 0.02 \text{ mm} \Rightarrow \text{Final} = \pm 0.02 \text{ mm}$$

Q45. If two quantities A and B have absolute errors of 0.01 and 0.02 respectively, the absolute error in (A + B) is:

- A. 0.01
- B. 0.02
- C. 0.03
- D. 0.002

Answer: C

Explanation:

$$\text{For addition: total absolute error} = \Delta A + \Delta B = 0.01 + 0.02 = 0.03$$