

¹ **Cross-scale interaction of host tree size and climate governs bark
2 beetle-induced tree mortality**

³ Michael J. Koontz^{1,2,3*}, Andrew M. Latimer^{1,2}, Leif A. Mortenson⁴, Christopher J. Fettig⁵, Malcolm P.
⁴ North^{1,2,6}

⁵ ¹Graduate Group in Ecology, University of California, Davis, CA, USA

⁶ ²Department of Plant Sciences, University of California, Davis, CA, USA

⁷ ³Earth Lab, University of Colorado-Boulder; Boulder, CO, USA

⁸ ⁴USDA Forest Service, Pacific Southwest Research Station, Placerville, CA, USA

⁹ ⁵USDA Forest Service, Pacific Southwest Research Station, Davis, CA, USA

¹⁰ ⁶USDA Forest Service, Pacific Southwest Research Station, Mammoth Lakes, CA, USA

¹¹ *Correspondence: michael.koontz@colorado.edu

¹² *Keywords:* *Dendroctonus brevicomis*, disturbance, drones, *Pinus ponderosa*, Sierra Nevada, structure from motion, forest structure, climate change-type drought, macroecology

¹⁴ *Abstract word count:* 310

¹⁵ *Overall .docx word count:* 11499

¹⁶ *Main text word count:* 4034 (Intro: 1264 Results: 368 (41+110+217); Discussion: 2402)

¹⁷ *Methods word count:* 3093 (661+642+1463+327)

¹⁸ *Text boxes word count:* 0

¹⁹ Date report generated: April 26, 2020

²⁰ **Abstract**

²¹ The Californian hot drought of 2012 to 2015 created favorable conditions for unprecedented ponderosa pine
²² (*Pinus ponderosa*) mortality in the Sierra Nevada mountain range, largely attributable to the western pine
²³ beetle (*Dendroctonus brevicomis*; WPB). Climate conditions and forest density may interact to affect tree
²⁴ mortality, but density is a coarse gauge of forest structure that can affect WPB behavior in a number of
²⁵ ways. Measuring broad-scale climate conditions simultaneously with local forest composition and structure–
²⁶ the spatial distribution and size of trees– will refine our understanding of how these variables interact, but
²⁷ is generally expensive and/or labor-intensive. We use drone surveys over 32 distinct sites along a 350-km
²⁸ latitudinal and 1000-m elevational gradient in western slope Sierra Nevada ponderosa pine/mixed-conifer
²⁹ forests and structure from motion (SfM) processing to segment and classify more than 450,000 trees over

30 9 km² of forest with WPB-induced tree mortality. We validated the segmentation and classification with
31 data from 160 coincident field plots (each 0.041 ha in area) throughout the 32 sites, with the caveat that
32 dead trees were all classified as ponderosa pine. We modeled the probability of ponderosa pine mortality as a
33 function of forest structure and composition and their interaction with site-level climatic water deficit (CWD),
34 accounting for spatial covariance using exact Gaussian processes. A greater local proportion of host trees
35 strongly increased the probability of host mortality, with greater host density amplifying this effect. Further,
36 we found a strong interaction between host size and CWD such that larger trees increased the probability of
37 host mortality at hot/dry sites, but smaller trees tended to drive mortality in cool/wet sites. Our results
38 demonstrate a variable response of WPB to local forest structure and composition across an environmental
39 gradient, which may help reconcile differences between observed ecosystem-wide tree mortality patterns and
40 predictions from models based on coarser-scale forest structure. Climate change adaptation strategies should
41 consider that future disturbance outcomes may depend on interactions between local forest structure and
42 broad-scale environmental gradients, with the potential for cross-scale interactions that challenge our current
43 understanding of forest insect dynamics.

44 Introduction

45 Bark beetles dealt the final blow to many of the nearly 150 million trees killed in the California hot drought
46 of 2012 to 2015 and its aftermath (USDAFS 2019). A harbinger of climate change effects to come, record
47 high temperatures exacerbated the drought (Griffin and Anchukaitis 2014), which increased water stress in
48 trees (Asner et al. 2016), making them more susceptible to colonization by bark beetles (Fettig 2012, Kolb
49 et al. 2016). Further, a century of fire suppression policy has enabled forests to grow into dense stands,
50 which can also makes them more vulnerable to bark beetles (Fettig 2012). This combination of environmental
51 conditions and forest structural characteristics led to tree mortality events of unprecedented size in the driest,
52 densest forests across the state (Young et al. 2017).

53 The mechanisms underlying the link between tree susceptibility to colonization by insects and hot, dry
54 conditions are often directly attributed to tree physiology (Bentz et al. 2010, Kolb et al. 2016), while the link
55 to forest density is multifaceted (Fettig 2012). Because forest density is a coarse metric of the forest features
56 to which bark beetles respond (Raffa et al. 2008), our understanding of the connection between forest density
57 and insect disturbance severity could be enhanced with more finely-resolved measures of forest structure as
58 well as explicit consideration of tree species composition (Stephenson et al. 2019, Fettig et al. 2019). Finally,
59 the challenge of simultaneously measuring the effects of both local-scale forest features (such as structure and
60 composition) and broad-scale environmental conditions (such as climatic water deicit; CWD) on forest insect

61 disturbance leaves their interaction effect relatively underexplored (Seidl et al. 2016, Stephenson et al. 2019,
62 Fettig et al. 2019).

63 The ponderosa pine/mixed-conifer forests in California's Sierra Nevada region are characterized by regular
64 bark beetle disturbances, primarily by the influence of western pine beetle (*Dendroctonus brevicomis*; WPB)
65 on its host ponderosa pine (*Pinus ponderosa*) (Fettig 2016). WPB is a primary bark beetle— its reproductive
66 success is contingent upon host tree mortality, which itself requires enough beetles to mass attack the host tree
67 and overwhelm its defenses (Raffa and Berryman 1983). This Allee effect creates a strong coupling between
68 beetle selection behavior of host trees and host tree susceptibility to colonization (Raffa and Berryman 1983,
69 Logan et al. 1998, Wallin and Raffa 2004). A key defense mechanism of conifers to bark beetle attack is to
70 flood beetle bore holes with resin, which physically expels colonizing beetles, can be toxic to the colonizers
71 and their fungi, and may interrupt beetle communication (Franceschi et al. 2005, Raffa et al. 2015). Under
72 normal conditions, weakened trees with compromised defenses are the most susceptible to colonization and
73 will be the main targets of primary bark beetles like WPB (Bentz et al. 2010, Boone et al. 2011, Raffa et al.
74 2015). Under severe water stress, many trees no longer have the resources available to mount a defense (Kolb
75 et al. 2016) and thus prolonged drought can often trigger increased bark beetle-induced tree mortality as
76 average tree vigor declines (Bentz et al. 2010). As the local population density of beetles increases due to
77 successful reproduction within spatially-aggregated weakened trees, as might occur during drought, mass
78 attacks grow in size and become capable of overwhelming formidable tree defenses such that even healthy
79 trees may be susceptible to colonization and mortality (Bentz et al. 2010, Boone et al. 2011, Raffa et al.
80 2015). Thus, water stress can be a key determinant of whether individual trees are susceptible to bark beetles
81 under many conditions, and this environmental condition may interact with beetle population dynamics to
82 drive tree susceptibility under extreme conditions (Bentz et al. 2010, Stephenson et al. 2019).

83 WPB activity is strongly influenced by forest structure— the spatial distribution and size of trees— and tree
84 species composition. Taking forest structure alone, high-density forests are more prone to bark beetle-induced
85 tree mortality (Fettig 2012) which may arise as greater competition for water resources amongst crowded trees
86 and thus average tree resistance is lower (Hayes et al. 2009), or because smaller gaps between trees protect
87 pheromone plumes from dissipation by the wind and thus enhance intraspecific beetle communication (Thistle
88 et al. 2004). Tree size is another aspect of forest structure that affects bark beetle host selection behavior with
89 smaller trees tending to have lower capacity for resisting attack, and larger trees being more desirable targets
90 on account of their thicker phloem providing greater nutritional content (Chubaty et al. 2009, Boone et al.
91 2011, Graf et al. 2012). Taking forest composition alone, WPB activity in the Sierra Nevada mountain range
92 of California is necessarily tied to the regional distribution of its exclusive host, ponderosa pine (Fettig 2016).

93 Colonization by primary bark beetles can also depend on the relative frequencies of tree species in a more
94 local area, akin to reduced oligophagous insect herbivory in forests comprising taxonomically-distinct tree
95 species compared to monocultures (Jactel and Brockerhoff 2007). The interaction between forest structure
96 and composition also drives WPB activity. For instance, high-density forests with high host availability
97 may experience greater beetle-induced tree mortality because dispersal distances between potential host
98 trees are shorter reducing predation of adults searching for hosts and facilitating higher rates of colonization
99 (Miller and Keen 1960, Berryman 1982, Fettig et al. 2007) or because high host availability reduces the
100 chance of individual beetles wasting their limited resources flying to and landing on a non-host tree (Moeck
101 et al. 1981, Evenden et al. 2014). Stand-scale measures of forest structure and composition thus paint a
102 fundamentally limited picture of the mechanisms by which these forest characteristics affect bark beetle
103 disturbance, but finer-grain information explicitly recognizing tree size, tree species, and local tree density
104 should more appropriately capture the ecological processes underlying insect-induced tree mortality (Kaiser
105 et al. 2013). Additionally, considering the effects of local forest structure and composition with the effects of
106 environmental conditions may help refine our understanding of tree mortality patterns in widespread events
107 such as during the recent California hot drought.

108 The vast spatial extent of tree mortality in the 2012 to 2015 California hot drought challenges our ability to
109 simultaneously consider how broad-scale environmental conditions may interact with local forest structure
110 and composition to affect the dynamic between bark beetle selection and colonization of host trees, and host
111 tree susceptibility to attack (Anderegg et al. 2015, Stephenson et al. 2019). Measuring local forest structure
112 generally requires expensive instrumentation (Kane et al. 2014, Asner et al. 2016) or labor-intensive field
113 surveys (Larson and Churchill 2012, Stephenson et al. 2019, Fettig et al. 2019), which constrains survey
114 extent and frequency. Small, unhumanned aerial systems (sUAS) enable relatively fast and cheap remote
115 imaging over dozens of hectares of forest, which can be used to measure complex forest structure at the
116 individual tree scale (Morris et al. 2017, Shiklomanov et al. 2019). Distributing such surveys across an
117 environmental gradient can overcome the data acquisition challenge inherent in investigating phenomena
118 with both a strong local- and a strong broad-scale component.

119 We used ultra-high resolution, sUAS-derived remote sensing data over a network of 32 sites in Sierra Nevada
120 ponderosa pine/mixed-conifer forests spanning 1000 m of elevation and 350 km of latitude (see Fettig et
121 al. 2019) and covering a total of 9 km² to ask how broad-scale environmental conditions interacted with
122 local forest structure and composition to affect the probability of tree mortality during the cumulative tree
123 mortality event of 2012 to 2018. We asked:

- 124 1. How does the proportion of host trees in a local area and average host tree size affect WPB-induced

- 125 tree mortality?
- 126 2. How does the density of all tree species (hereafter “overall density”) affect WPB-induced tree mortality?
- 127 3. How does environmentally-driven tree moisture stress affect WPB-induced tree mortality?
- 128 4. Do the effects of forest structure, forest composition, and environmental condition interact to influence
- 129 WPB-induced tree mortality?

130 **Methods**

131 **Study system**

132 We built our study coincident with 160 vegetation/forest insect monitoring plots at 32 sites established
133 between 2016 and 2017 by Fettig et al. (2019) (Figure 1). The study sites were chosen to reflect typical
134 west-side Sierra Nevada yellow pine/mixed-conifer forests and were dominated by ponderosa pine (Fettig
135 et al. 2019). Plots were located in WPB-attacked, yellow pine/mixed-conifer forests across the Eldorado,
136 Stanislaus, Sierra and Sequoia National Forests and were stratified by elevation (914-1219 m, 1219-1524
137 m, 1524-1829 m above sea level). In the Sequoia National Forest, the southernmost National Forest in our
138 study, plots were stratified with the lowest elevation band of 1219-1524 m and extended to an upper elevation
139 band of 1829-2134 m to capture a more similar forest community composition as at the more northern
140 National Forests. The sites have variable forest structure and plot locations were selected in areas with >35%
141 ponderosa pine basal area and >10% ponderosa pine mortality. At each site, five 0.041 ha circular plots
142 were installed along transects with 80 to 200m between plots. In the field, Fettig et al. (2019) mapped all
143 stem locations relative to the center of each plot using azimuth/distance measurements. Tree identity to
144 species, tree height, and diameter at breast height (DBH) were recorded if DBH was greater than 6.35cm.
145 Year of mortality was estimated based on needle color and retention if it occurred prior to plot establishment,
146 and was directly observed thereafter during annual site visits. A small section of bark (approximately 625
147 cm²) on both north and south aspects was removed from dead trees to determine if bark beetle galleries
148 were present. The shape, distribution, and orientation of galleries are commonly used to distinguish among
149 bark beetle species (Fettig 2016). In some cases, deceased bark beetles were present beneath the bark to
150 supplement identifications based on gallery formation. During the spring and early summer of 2018, all field
151 plots were revisited to assess whether dead trees had fallen (Fettig et al. 2019).

- 152 In the typical life cycle of WPBs, females initiate host colonization by tunneling through the outer bark and
153 into the phloem and outer xylem where they rupture resin canals.
- 154 As a result, oleoresin exudes and collects on the bark surface, as is commonly observed with other bark beetle




Figure 1: The network of field plots spanned a 350-km latitudinal gradient from the Eldorado National Forest in the north to the Sequoia National Forest in the south. Plots were stratified by three elevation bands in each forest, with the plots in the Sequoia National Forest (the southern-most National Forest) occupying elevation bands 305 m above the three bands in the other National Forests in order to capture a similar community composition.


155 species. During the early stages of attack, females release an aggregation pheromone component which, in
156 combination with host monoterpenes released from pitch tubes, is attractive to conspecifics (Bedard et al.
157 1969). An antiaggregation pheromone component is produced during latter stages of host colonization by
158 several pathways, and is thought to reduce intraspecific competition by altering adult behavior to minimize
159 overcrowding of developing brood within the host (Byers and Wood 1980). Volatiles from several nonhosts
160 sympatric with ponderosa pine have been demonstrated to inhibit attraction of WPB to its aggregation
161 pheromones (Shepherd et al. 2007, Fettig and Hilszczański 2015). In California, WPB generally has 2-3
162 generations in a single year and can often out-compete other primary bark beetles such as the mountain pine
163 beetle (*Dendroctonus ponderosae*), in ponderosa pines, especially in larger trees (Miller and Keen 1960).

164 **Aerial data collection and processing**

165 Nadir-facing imagery was captured using a gimbal-stabilized DJI Zenmuse X3 broad-band red/green/blue
166 (RGB) camera (DJI 2015a) and a fixed-mounted Micasense Rededge3 multispectral camera with five narrow
167 bands (Micasense 2015) on a DJI Matrice 100 aircraft (DJI 2015b). Imagery was captured from both cameras
168 along preprogrammed aerial transects over ~40 hectares surrounding each of the 32 sites (each of these
169 containing five field plots) and was processed in a series of steps to yield local forest structure and composition
170 data suitable for our statistical analyses. All images were captured in 2018 during a 3-month period between
171 early April and early July, and thus our work represents a postmortem investigation into the drivers of
172 cumulative tree mortality through the course of the hot drought.

173 Following the call by Wyngaard et al. (2019), we establish “data product levels” to reflect the image processing
174 pipeline from raw imagery (Level 0) to calibrated, fine-scale forest structure and composition information on
175 regular grids (Level 4), with each new data level derived from levels below it. Here, we outline the steps in the
176 processing and calibration pipeline visualized in Figure 2, and include additional details in the Supplemental
177 Information.


Level 0: raw data from sensors




Level 1: basic outputs from photogrammetric processing




Level 2: corrected outputs from photogrammetric processing



Level 3: domain-specific information extraction



Level 4: aggregations to regular grids



179 Figure 2. Schematic of the data processing workflow for a single site with each new data product level derived
180 from data at lower levels.

181 Level 0 represents raw data from the sensors. From left to right: example broad-band RGB photo from
182 DJI Zenmuse X3 camera, example blue photo from Rededge3 (centered on 475nm), example green photo
183 from Rededge3 (centered on 560nm), example red photo from Rededge3 (centered on 668nm), example near
184 infrared photo from Rededge3 (centered on 840nm), and example red edge photo from Rededge3 (centered on
185 717nm).

186 Level 1 represents basic outputs from the photogrammetric workflow, in this case implemented with
187 Pix4Dmapper. From left to right: a dense point cloud visualized in CloudCompare (<https://www.danielgm.net/cc/>), an orthophoto generated from the RGB camera, and a digital surface model representing the
188 altitude above sea level (ground height + vegetation height) for every cell.

190 Level 2 represents outputs from photogrammetric processing that have been corrected radiometrically or
191 geometrically. From left to right: a radiometrically-corrected surface reflectance map of the red narrow band
192 from the Rededge3 camera, a radiometrically-corrected surface reflectance map of the near infrared narrow
193 band from the Rededge3 camera, a rasterized version of the digital terrain model derived by a geometric
194 correction of the dense point cloud, and a canopy height model derived by subtracting the terrain height
195 from the digital surface model.

196 Level 3 represents domain-specific information extraction from Level 2 products and is divided into two
197 sub-levels. Level 3a products are derived using only spectral or only geometric data. From left to right: a
198 reflectance map of Normalized Difference Vegetation Index (NDVI; Rouse et al. (1973)) derived using the red
199 and near infrared Level 2 reflectance products, a map of points representing detected trees from the canopy
200 height model with a red polygon highlighting the area presented in more detail for the next two images, a
201 close-up of points representing detected trees, and a close-up of polygons representing segmented tree crowns.
202 Level 3b products are derived using both spectral and geometric data. From left to right: a map of the point
203 locations of detected trees that have been classified as alive or dead based on the pixel values within each
204 segmented tree crown and a map of the point locations of detected trees classified to WPB host/non-host
205 using the same spectral information. Note that our study relies on the generation of Level 3a products in
206 order to combine them and create Level 3b products, but this need not be the case. For instance, deep
207 learning/neural net methods may be able to use both the spectral and geometric information from Level 2
208 simultaneously to locate and classify trees in a scene and directly generate Level 3b products without a need
209 to first generate the Level 3a products shown in this schematic (Weinstein et al. 2019, dos Santos et al. 2019).

210 Level 4 represents aggregations of Level 3 products to regular grids which might better reflect the grain size
211 of the data for which we have the best calibration and thus the most confidence or which might provide
212 new information not possible at an individual-tree level (e.g., average distance between trees in a small
213 neighborhood). From left to right: aggregation of live/dead classified trees as fraction of dead trees in a 20 x
214 20-m cell, aggregation of host/non-host classified trees as fraction of hosts in a 20 x 20-m cell, aggregation of
215 mean host height in a 20 x 20-m cell, and aggregation of tree count (including all species), in a 20 x 20-m
216 cell. In our case, the 20 x 20-m aggregation produces a grid cell with an area of 400 m², which most closely
217 matches the 404-m² area of the ground-based vegetation plots whose data we used in an aggregated form to
218 calibrate our derivation of Level 3 products.

219 **Level 0: Raw data from sensors**

220 Raw data comprised approximately 1900 images per camera lens (one broad-band RGB lens and five narrow-
221 band multispectral lenses) for each of the 32 sites (Figure 2; Level 0). Prior to the aerial survey, two strips of
222 bright orange drop cloth (~100 x 15 cm) were positioned as an “X” over the permanent monuments marking
223 the center of the 5 field plots from Fettig et al. (2019) (see Supplemental Information).

224 We preprogrammed north-south aerial transects using Map Pilot for DJI on iOS flight software (Drones-
225 MadeEasy 2018) at an altitude of 120 m above ground level (with “ground” defined using a 1-arc-second
226 digital elevation model (Farr et al. 2007)). The resulting ground sampling distance was approximately 5
227 cm/px for the Zenmuse X3 RGB camera and approximately 8 cm/px for the Rededge3 multispectral camera.
228 We used 91.6% image overlap (both forward and side) at the ground for the Zenmuse X3 RGB camera and
229 83.9% overlap (forward and side) for the Rededge3 multispectral camera.

230 **Level 1: Basic outputs from photogrammetric processing**

231 We used SfM photogrammetry implemented in Pix4Dmapper Cloud (www.pix4d.com) to generate dense point
232 clouds (Figure 2; Level 1, left), orthophotos (Figure 2; Level 1, center), and digital surface models (Figure 2;
233 Level 1, right) for each field site (Frey et al. 2018). For 29 sites, we processed the Rededge3 multispectral
234 imagery alone to generate these products. For three sites, we processed the RGB and the multispectral
235 imagery together to enhance the point density of the dense point cloud. All SfM projects resulted in a single
236 processing “block,” indicating that all images in the project were optimized and processed together. The
237 dense point cloud represents x, y, and z coordinates as well as the color of millions of points per site. The
238 orthophoto represents a radiometrically uncalibrated, top-down view of the survey site that preserves the
239 relative x-y positions of objects in the scene. The digital surface model is a rasterized version of the dense

240 point cloud that shows the altitude above sea level for each pixel in the scene at the ground sampling distance
241 of the camera that generated the Level 0 data.

242 **Level 2: Corrected outputs from photogrammetric processing**

243 **Radiometric corrections**

244 A radiometrically-corrected reflectance map (Figure 2; Level 2, left two figures; i.e., a corrected version of the
245 Level 1 orthophoto) was generated using the Pix4D software by incorporating incoming light conditions for
246 each narrow band of the Rededge3 camera (captured simultaneously with the Rededge3 camera using an
247 integrated downwelling light sensor) as well as a pre-flight image of a calibration panel of known reflectance
248 (see Supplemental Information for camera and calibration panel details).

249 **Geometric corrections**

250 We implemented a geometric correction to the Level 1 dense point cloud and digital surface model by
251 normalizing these data for the terrain underneath the vegetation. We generated the digital terrain model
252 representing the ground underneath the vegetation at 1-m resolution (Figure 2; Level 2, third image) by
253 classifying each survey area's dense point cloud into "ground" and "non-ground" points using a cloth simulation
254 filter algorithm (Zhang et al. 2016) implemented in the `lidR` (Roussel et al. 2019) package and rasterizing
255 the ground points using the `raster` package (Hijmans et al. 2019). We generated a canopy height model
256 (Figure 2; Level 2, fourth image) by subtracting the digital terrain model from the digital surface model.

257 **Level 3: Domain-specific information extraction**

258 **Level 3a: Data derived from spectral OR geometric Level 2 product**

259 Using just the spectral information from the radiometrically-corrected reflectance maps, we calculated several
260 vegetation indices including the normalized difference vegetation index (NDVI; Rouse et al. (1973); Figure
261 2; Level 3a, first image), the normalized difference red edge (NDRE; Gitelson and Merzlyak (1994)), the
262 red-green index (RGI; Coops et al. (2006)), the red edge chlorophyll index ($CI_{red\ edge}$; Clevers and Gitelson
263 (2013)), and the green chlorophyll index (CI_{green} ; Clevers and Gitelson (2013)).

Table 1: Algorithm name, number of parameter sets tested for each algorithm, and references.

Algorithm	Parameter sets tested	Reference(s)
li2012	131	Li et al. (2012); Jakubowski et al. (2013); Shin et al. (2018)

Algorithm	Parameter sets tested	Reference(s)
lmfx	30	Roussel (2019)
localMaxima	6	Roussel et al. (2019)
multichm	1	Eysn et al. (2015)
ptrees	3	Vega et al. (2014)
vwf	3	Plowright (2018)
watershed	3	Pau et al. (2010)

Using just the geometric information from the canopy height model or terrain-normalized dense point cloud, we generated maps of detected trees (Figure 2; Level 3a, second and third images) by testing a total of 7 automatic tree detection algorithms and a total of 177 parameter sets (Table 1). We used the field plot data to assess each tree detection algorithm/parameter set by converting the distance-from-center and azimuth measurements of the trees in the field plots to x-y positions relative to the field plot centers distinguishable in the Level 2 reflectance maps as the orange fabric X's that we laid out prior to each flight. In the reflectance maps, we located 220 out of 160 field plot centers while some plot centers were obscured due to dense interlocking tree crowns or because a plot center was located directly under a single tree crown. For each of the 220 field plots with identifiable plot centers— the “validation field plots”, we calculated 7 forest structure metrics using the ground data collected by Fettig et al. (2019): total number of trees, number of trees greater than 15 m in height, mean height of trees, 25th percentile tree height, 75th percentile tree height, mean distance to nearest tree neighbor, and mean distance to second nearest neighbor. For each tree detection algorithm and parameter set described above, we calculated the same set of 7 structure metrics within the footprint of the validation field plots. We calculated the Pearson’s correlation and root mean square error (RMSE) between the ground data and the aerial data for each of the 7 structure metrics for each of the 177 automatic tree detection algorithms/parameter sets. For each algorithm and parameter set, we calculated its performance relative to other algorithms as whether its Pearson’s correlation was within 5% of the highest Pearson’s correlation as well as whether its RMSE was within 5% of the lowest RMSE. We summed the number of forest structure metrics for which it reached these 5% thresholds for each algorithm/parameter set. For automatically detecting trees across the whole study, we selected the algorithm/parameter set that performed well across the most number of forest metrics (see Results).

We delineated individual tree crowns (Figure 2; Level 3a, fourth image) with a marker controlled watershed segmentation algorithm (Meyer and Beucher 1990) implemented in the **ForestTools** package (Plowright

287 2018) using the detected treetops as markers. If the automatic segmentation algorithm failed to generate
288 a crown segment for a detected tree (e.g., often snags with a very small crown footprint), a circular crown
289 was generated with a radius of 0.5 m. If the segmentation generated multiple polygons for a single detected
290 tree, only the polygon containing the detected tree was retained. Because image overlap decreases near the
291 edges of the overall flight path and reduces the quality of the SfM processing in those areas, we excluded
292 segmented crowns within 35 m of the edge of the survey area. Given the narrower field of view of the
293 Rededge3 multispectral camera versus the X3 RGB camera whose optical parameters were used to define the
294 ~40 hectare survey area around each site, as well as the 35 m additional buffering, the survey area at each
295 site was ~30 ha (see Supplemental Information).

296 **Level 3b: Data derived from spectral AND geometric information**

297 We overlaid the segmented crowns on the reflectance maps from 20 sites spanning the latitudinal and elevation
298 gradient in the study. Using QGIS (<https://qgis.org/en/site/>), we hand classified 564 trees as live/dead
299 (Figure 3) and as one of 5 dominant species in the study area (ponderosa pine, *Pinus lambertiana*, *Abies*
300 *concolor*, *Calocedrus decurrens*, or *Quercus kelloggii*) using the mapped ground data as a guide. Each tree was
301 further classified as “host” for ponderosa pine or “non-host” for all other species (Fettig 2016). We extracted
302 all the pixel values within each segmented crown polygon from the five, Level 2 orthorectified reflectance
303 maps (one per narrow band on the Rededge3 camera) as well as from the five, Level 3a vegetation index
304 maps using the **velox** package (Hunziker 2017). For each crown polygon, we calculated the mean value of the
305 extracted Level 2 and Level 3a pixels and used them as ten independent variables in a five-fold cross validated
306 boosted logistic regression model to predict whether the hand classified trees were alive or dead. For just the
307 living trees, we similarly used all 10 mean reflectance values per crown polygon to predict tree species using a
308 five-fold cross validated regularized discriminant analysis. The boosted logistic regression and regularized
309 discriminant analysis were implemented using the **caret** package in R (Kuhn 2008). We used these models to
310 classify all tree crowns in the data set as alive or dead (Figure 2; Level 3b, first image) as well as the species
311 of living trees (Figure 2; Level 3b, second image). Finally, we estimated the basal area of each tree from their
312 photogrammetry-derived height using species-specific simple linear regressions of the relationship between
313 height and diameter at breast height as measured in the coincident field plots from Fettig et al. (2019).

314 **Level 4: Aggregations to regular grids**

315 We rasterized the forest structure and composition data at a spatial resolution similar to that of the field
316 plots to better match the grain size at which we validated the automatic tree detection algorithms. In each
317 raster cell, we calculated: number of dead trees, number of ponderosa pine trees, total number of trees, and

318 mean height of ponderosa pine trees. The values of these variables in each grid cell and derivatives from
319 them were used for visualization and modeling. Here, we show the fraction of dead trees per cell (Figure 2;
320 Level 4, first image), the fraction of host trees per cell (Figure 2; Level 4, second image), the mean height of
321 ponderosa pine trees in each cell (Figure 2; Level 4, third image), and the total count of trees per cell (Figure
322 2; Level 4, fourth image).

323 **Note on assumptions about dead trees**

324 For the purposes of this study, we assumed that all dead trees were ponderosa pine and thus hosts colonized
325 by WPB. This is a reasonably good assumption for our study area; for example, Fettig et al. (2019) found
326 that 73.4% of dead trees in their coincident field plots were ponderosa pine. Mortality was concentrated in
327 the larger-diameter classes and attributed primarily to WPB (see Figure 5 of Fettig et al. 2019). The species
328 contributing to the next highest proportion of dead trees was incense cedar which represented 18.72% of the
329 dead trees in the field plots. While the detected mortality is most likely to be ponderosa pine killed by WPB,
330 it is critical to interpret our results with these limitations in mind.

331 **Environmental data**

332 We used CWD (Stephenson 1998) from the 1981-2010 mean value of the basin characterization model (Flint
333 et al. 2013) as an integrated measure of temperature and moisture conditions for each of the 32 sites. Higher
334 values of CWD correspond to hotter, drier conditions and lower values correspond to cooler, wetter conditions.
335 CWD has been shown to correlate well with broad patterns of tree mortality in the Sierra Nevada (Young
336 et al. 2017) as well as bark beetle-induced tree mortality (Millar et al. 2012). The forests along the entire
337 CWD gradient used in this study experienced exceptional hot drought between 2012 to 2015 (Griffin and
338 Anchukaitis 2014, Robeson 2015). We converted the CWD value for each site into a z-score representing that
339 site's deviation from the mean CWD across the climatic range of Sierra Nevada ponderosa pine as determined
340 from 179 herbarium records described in Baldwin et al. (2017). Thus, a CWD z-score of 1 would indicate
341 that the CWD at that site is one standard deviation hotter/drier than the mean CWD across all geolocated
342 herbarium records for ponderosa pine in the Sierra Nevada.

343 **Statistical model**

344 We used a generalized linear model with a zero-inflated binomial response and a logit link to predict the
345 probability of ponderosa pine mortality within each 20 x 20-m cell using the total number of ponderosa
346 pine trees in each cell as the number of trials, and the number of dead trees in each cell as the number of
347 "successes". As covariates, we used the proportion of trees that are WPB hosts (i.e., ponderosa pine) in each

348 cell, the mean height of ponderosa pine trees in each cell, the count of trees of all species (overall density) in
 349 each cell, and the site-level CWD using Eq. 1. Note that the two-way interaction between the overall density
 350 and the proportion of trees that are hosts is directly proportional to the number of ponderosa pine trees in
 351 the cell. We centered and scaled all predictor values, and used weakly-regularizing default priors from the
 352 `brms` package (Bürkner 2017). To measure and account for spatial autocorrelation underlying ponderosa pine
 353 mortality, we subsampled the data at each site to a random selection of 200, 20 x 20-m cells representing
 354 approximately 27.5% of the surveyed area. Additionally with these subsampled data, we included a separate
 355 exact Gaussian process term per site of the noncentered/nonscaled interaction between the x- and y-position
 356 of each cell using the `gp()` function in the `brms` package (Bürkner 2017). The Gaussian process estimates the
 357 spatial covariance in the response variable (log-odds of ponderosa pine mortality) jointly with the effects of
 358 the other covariates.

$$y_{i,j} \sim \begin{cases} 0, & p \\ Binom(n_i, \pi_i), & 1 - p \end{cases}$$

$$\begin{aligned}
 logit(\pi_i) = & \beta_0 + \\
 & \beta_1 X_{cwd,j} + \beta_2 X_{propHost,i} + \beta_3 X_{PipoHeight,i} + \\
 & \beta_4 X_{overallDensity,i} + \beta_5 X_{overallBA,i} + \\
 & \beta_6 X_{cwd,j} X_{PipoHeight,i} + \beta_7 X_{cwd,j} X_{propHost,i} + \\
 & \beta_8 X_{cwd,j} X_{overallDensity,i} + \beta_9 X_{cwd,j} X_{overallBA,i} + \\
 & \beta_{10} X_{propHost,i} X_{PipoHeight,i} + \beta_{11} X_{propHost,i} X_{overallDensity,i} + \\
 & \beta_{12} X_{cwd,j} X_{propHost,i} X_{PipoHeight,i} + \\
 & \mathcal{GP}_j(x_i, y_i)
 \end{aligned}$$

359 Where y_i is the number of dead trees in cell i , n_i is the sum of the dead trees (assumed to be ponderosa pine)
 360 and live ponderosa pine trees in cell i , π_i is the probability of ponderosa pine tree mortality in cell i , p is the
 361 probability of there being zero dead trees in a cell arising as a result of an independent, unmodeled process,
 362 $X_{cwd,j}$ is the z-score of CWD for site j , $X_{propHost,i}$ is the scaled proportion of trees that are ponderosa pine
 363 in cell i , $X_{PipoHeight,i}$ is the scaled mean height of ponderosa pine trees in cell i , $X_{overallDensity,i}$ is the scaled
 364 density of all trees in cell i , $X_{overallBA,i}$ is the scaled basal area of all trees in cell i , x_i and y_i are the x- and
 365 y- coordinates of the centroid of the cell in an EPSG3310 coordinate reference system, and \mathcal{GP}_j represents
 366 the exact Gaussian process describing the spatial covariance between cells at site j .

367 We fit this model using the `brms` package (Bürkner 2017) which implements the No U-Turn Sampler extension
368 to the Hamiltonian Monte Carlo algorithm (Hoffman and Gelman 2014) in the Stan programming language
369 (Carpenter et al. 2017). We used 4 chains with 4000 iterations each (2000 warmup, 2000 samples), and
370 confirmed chain convergence by ensuring all `Rhat` values were less than 1.1 (Brooks and Gelman 1998) and
371 that the bulk and tail effective sample sizes (ESS) for each estimated parameter were greater than 100 times
372 the number of chains (i.e., greater than 400 in our case). We used posterior predictive checks to visually
373 confirm model performance by overlaying the density curves of the predicted number of dead trees per cell
374 over the observed number (Gabry et al. 2019). For the posterior predictive checks, we used 50 random
375 samples from the model fit to generate 50 density curves and ensured curves were centered on the observed
376 distribution, paying special attention to model performance at capturing counts of zero.

377 **Software and data availability**

378 All data are available via the Open Science Framework. Statistical analyses were performed using the `brms`
379 packages. With the exception of the SfM software (Pix4Dmapper Cloud) and the GIS software QGIS, all
380 data carpentry and analyses were performed using R (R Core Team 2018).

381 **Results**

382 **Tree detection algorithm performance**

383 We found that the experimental `lmfx` algorithm with parameter values of `dist2d = 1` and `ws = 2.5` (Roussel
384 et al. 2019) performed the best across 7 measures of forest structure as measured by Pearson's correlation
385 with ground data (Table 2).

Table 2: Correlation and differences between the best performing tree detection algorithm (`lmfx` with `dist2d = 1` and `ws = 2.5`) and the ground data. An asterisk next to the correlation or RMSE indicates that this value was within 5% of the value of the best-performing algorithm/parameter set. Ground mean represents the mean value of the forest metric across the 220 field plots that were visible from the sUAS-derived imagery. The median error is calculated as the median of the differences between the air and ground values for the 220 visible plots. Thus, a positive number indicates an overestimate by the sUAS workflow and a negative number indicates an underestimate.

Forest structure metric	Ground mean	Correlation with ground	RMSE	Median error
total tree count	19	0.67*	8.68*	2
count of trees > 15 m	9.9	0.43	7.38	0
distance to 1st neighbor (m)	2.8	0.55*	1.16*	0.26
distance to 2nd neighbor (m)	4.3	0.61*	1.70*	0.12

Forest structure metric	Ground mean	Correlation with ground	RMSE	Median error
height (m); 25 th percentile	12	0.16	8.46	-1.2
height (m); mean	18	0.29	7.81*	-2.3
height (m); 75 th percentile	25	0.35	10.33*	-4

386 **Classification accuracy for live/dead and host/non-host**

387 The accuracy of live/dead classification on a withheld test dataset was 96.4%. The accuracy of species
 388 classification on a withheld testing dataset was 64.1%. The accuracy of WPB host/non-WPB-host (i.e.,
 389 ponderosa pine versus other tree species) on a withheld testing dataset was 71.8%.

390 **Site summary based on best tree detection algorithm and classification**

391 Across all study sites, we detected, segmented, and classified 452,413 trees. Of these trees, we classified
 392 118,879 as dead (26.3% mortality). Estimated site-level tree mortality ranged from 6.8% to 53.6%. See
 393 Supplemental Information for site summaries and comparisons to site-level mortality measured from field
 394 data.

395 **Effect of local structure and regional climate on tree mortality attributed to western pine
 396 beetle**

397 We detected a positive main effect of CWD on the probability of ponderosa pine mortality within each 20 x
 398 20-m cell (Figure 4). We found a positive main effect of proportion of host trees per cell (effect size: 0.76;
 399 95% CI: [0.70, 0.82]), with a greater proportion of host trees (i.e., ponderosa pine) in a cell increasing the
 400 probability of ponderosa pine mortality. We detected no effect of overall tree density nor overall basal area
 401 (i.e., including both ponderosa pine and non-host species; tree density effect size: -0.05; 95% CI: [-0.13, 0.03];
 402 basal area effect size: 0.00; 95% CI: [-0.11, 0.11]).

403 We found a positive two-way interaction between the overall tree density per cell and the proportion of trees
 404 that were hosts, which is equivalent to a positive effect of the density of host trees (effect size: 0.08; 95% CI:
 405 [0.03, 0.13]; Figure 4).

406 We found a negative effect of mean height of ponderosa pine on the probability of ponderosa mortality,
 407 suggesting that WPB attacked smaller trees, on average (effect size: -0.40; 95% CI: [-0.50, -0.30]). However,
 408 there was a positive interaction between CWD and ponderosa pine mean height, such that larger trees were




Figure 4: Posterior distributions of effect size from zero-inflated binomial model predicting the probability of ponderosa pine mortality in a 20 x 20-m cell given forest structure characteristics and site-level climatic water deficit (CWD). The gray filled area for each model covariate represents the probability density of the posterior distribution, the point underneath each density curve represents the median of the estimate, the bold interval surrounding the point estimate represents the 66% credible interval, and the thin interval surrounding the point estimate represents the 95% credible interval.

409 more likely to increase the local probability of ponderosa mortality in hotter, drier sites (effect size: 0.29;
410 95% CI: [0.12, 0.46]; Figure 5).

411 We found weakly negative effects of the site-level CWD interactions with both the proportion of host trees
412 and overall tree density (CWD/proportion host interaction effect size: -0.13; 95% CI: [-0.23, -0.03]; Figure 4;
413 CWD/overall tree density interaction effect size: -0.22; 95% CI: [-0.34, -0.11]; Figure 4; Figure 5). We found
414 a positive effect of the interaction between CWD and total basal area (effect size: 0.22; 95% CI: [0.03, 0.42];
415 Figure 4; Figure 5).




Figure 5: Line version of model results with 95% credible intervals showing primary influence of ponderosa pine structure on the probability of ponderosa pine mortality, and the interaction across climatic water deficit. The ‘larger trees’ line represents the mean height of ponderosa pine 0.7 standard deviations above the mean (approximately 24.1 m), and the ‘smaller trees’ line represents the mean height of ponderosa pine 0.7 standard deviations below the mean (approximately 12.1 m).

416 Discussion

417 This study represents a novel use of drones to further our understanding of the simultaneous effects of
418 local forest structure and composition with broad-scale environmental gradients on tree mortality attributed
419 to WPB. We found strong positive effects (effect sizes >0.4) of both the proportion of host trees and the
420 interaction between site CWD and host tree mean size (height) on the probability of ponderosa pine mortality.
421 Conversely, we found a strong negative effect (effect size <-0.4) of mean height of ponderosa pine. Site-level
422 CWD exerted a positive, but relatively weak, main effect on the probability of ponderosa mortality (effect

size: 0.16; 95% CI: [0.03, 0.30]). To that end, we did not measure tree water stress at an individual tree level as in other recent work (Stephenson et al. 2019), and instead treated CWD as a general indicator of tree stress following results of coarser-scale studies (e.g., Young et al. 2017), which may have contributed to our failure to detect a stronger CWD effect. Also, our entire study area experienced the same extreme hot drought between 2012 and 2015 and the variation of mortality explained by a main effect of CWD may be dampened when most trees are experiencing a high degree of water stress (Floyd et al. 2009, Fettig et al. 2019).

Positive effect of host density and a negative effect of overall density

A number of mechanisms associated with the relative abundance of species in a local area might underlie the strong effect of host proportion on the probability of host tree mortality. Frequency-dependent herbivory—whereby mixed-species forests experience less herbivory compared to monocultures (as an extreme example)—is common, especially for oligophagous insect species (Jactel and Brockerhoff 2007). Furthermore, it has been demonstrated that nonhost volatiles reduce attraction of several species of bark beetles to their aggregation pheromones (Seybold et al. 2018), including WPB (Fettig et al. 2005). To that end, combinations of nonhost volatiles and an antiaggregation pheromone have been used successfully to reduce levels of tree mortality attributed to WPB (e.g., Fettig et al. 2012). In general, Hayes et al. (2009) and Fettig et al. (2019) found that measures of host availability explained less variation in mortality than measures of overall tree density, but those conclusions were based on a response variable of “total number of dead host trees,” rather than the number of dead host trees conditional on the total number of host trees as in our study (i.e., a binomial response).

The negative relationship between overall tree density, a potential correlate of the local competitive environment, and the probability of ponderosa pine mortality is counter-intuitive but corroborates findings of coincident ground plots (Fettig et al. 2019, in their analysis using proportion of trees killed as a response) and other work during the same hot drought (Restaino et al. 2019). The forest structure (in the absence of management) is itself a product of climate and, with increasing importance at finer spatial scales, topographic conditions (Fricker et al. 2019). Thus, the denser forest patches in our study may indicate greater local water availability, more favorable conditions for tree growth and survivorship, and increased resistance to beetle-induced mortality (Restaino et al. 2019). The negative two-way interaction between site CWD and overall density that amplifies the negative overall density effect in hotter, drier sites (effect size: -0.22; 95% CI: [-0.34, -0.11]) supports this explanation if greater local tree density implies especially favorable growing conditions (and locally resistant trees) when denser patches are found in hot, dry sites.

454 The positive relationship between host density and susceptibility to colonization by bark beetles has been so
455 well-documented at the experimental plot level (e.g., Raffa and Berryman 1987, Oliver 1995) that lowering
456 stand densities through selective harvest of hosts is commonly recommended for reducing future levels of tree
457 mortality attributed to bark beetles (Fettig and Hilszczański 2015), including WPB (Fettig 2016). Greater
458 host density shortens the flight distance required for WPB to disperse to new hosts, which likely facilitates
459 bark beetle spread, however we calibrated our aerial tree detection to ~400 m² areas rather than to individual
460 tree locations, so our data are insufficient to address these relationships. Increased density of ponderosa pine,
461 specifically, may disproportionately increase the competitive environment for host trees (and thus increase
462 their susceptibility to WPB colonization) if intraspecific competition amongst ponderosa pine trees is stronger
463 than interspecific competition as would be predicted with coexistence theory (Chesson 2000). Finally, greater
464 host densities increase the frequency that searching WPB land on hosts, rather than nonhosts, thus reducing
465 the amount of energy expended during host finding and selection as well as the time that searching WPB
466 spend exposed to predators.

467 **Negative main effect of host tree mean size, but strong positive interaction with site CWD**

468 The negative main effect of host tree mean size was surprising, and appears to contradict long-standing
469 wisdom on the dynamics of western pine beetle in the Sierra Nevada. WPB exhibit a preference for trees 50.8
470 to 76.2 cm in diameter at breast height (Person 1928, 1931), and a positive relationship between host tree
471 size and levels of tree mortality attributed to WPB was reported by Fettig et al. (2019) in the coincident field
472 plots as well as in other recent studies (Restaino et al. 2019, Stephenson et al. 2019, Pile et al. 2019). Indeed,
473 Fettig et al. (2019) reported no mortality in ponderosa pine trees <10.0 cm DBH attributable to WPB and
474 found no tree size/mortality relationship for incense cedar or white fir in the coincident field plots. These
475 species represent 22.3% of the total tree mortality observed in their study, yet in our study all dead trees
476 were classified as ponderosa pine (see Methods) which could dampen positive effect of tree size on mortality.
477 Larger trees are more nutritious and are therefore ideal targets if local bark beetle density is high enough
478 to successfully initiate mass attack as can occur when many trees are under severe water stress (Bentz et
479 al. 2010, Kolb et al. 2016). In the recent hot drought, we expected that most trees would be under severe
480 water stress, setting the stage for increasing beetle density, successful mass attacks, and targeting of larger
481 trees. A possible explanation for our finding counter to this expectation is that our observations represent the
482 cumulative mortality of trees during a multi-year drought event and its aftermath. Lower host tree mean size
483 led to a greater probability of host mortality earlier in this drought (Pile et al. 2019, Stovall et al. 2019) and
484 that signal might have persisted even as mortality continued to accumulate driven by other factors. Finally,
485 tree growth rates may be a better predictor of susceptibility to WPB colonization than tree size per se, with

486 slower-growing trees being most vulnerable (Miller and Keen 1960). While slow-growing trees are often also
487 the largest trees, this may not be the case for our study sites especially given the legacy of fire exclusion
488 in the Sierra Nevada and its effect of perturbing forest structure far outside its natural range of variation
489 (Safford and Stevens 2017).

490 In hot, dry sites, larger average host size increased the probability of host mortality while smaller host sizes
491 increased the probability of host mortality in cooler, wetter sites. Notably, a similar pattern was shown
492 by Stovall et al. (2019) with a strong positive tree height/mortality relationship in areas with the greatest
493 vapor pressure deficit and no tree height/mortality relationship in areas with the lowest vapor pressure
494 deficit. Stovall et al. (2019) did not observe that this environmental dependence extended to a negative tree
495 height/mortality relationship (as we did) even at the lowest extremes of their vapor pressure deficit gradient,
496 perhaps because their entire study took place in the southern Sierra Nevada which represents a hotter, drier
497 portion of the more spatially extensive results we present here. Our work suggests that the WPB was cueing
498 into different aspects of forest structure across an environmental gradient in a spatial context in a parallel
499 manner to the temporal context noted by Stovall et al. (2019) and Pile et al. (2019), who observed that
500 mortality was increasingly driven by larger trees as the hot drought proceeded and became more severe.

501 All of our sites were considered in an “epidemic” population phase for WPB (>5 trees killed per hectare; see
502 Supplemental Information; Miller and Keen 1960, Hayes et al. 2009), but our results challenge the notion that
503 outbreak behavior by the WPB and subsequent tree mortality is always driven by greater tree size. Despite a
504 strong tree size/mortality relationship in coincident ground plots across our study area (Fettig et al. 2019),
505 our results from surveying the broader context surrounding those ground plots reveals different effects of
506 host tree size depending on CWD. Thus, it is possible that the massive tree mortality in hotter/drier Sierra
507 Nevada forests (lower latitudes and elevations; Asner et al. 2016, Young et al. 2017) during the 2012 to
508 2015 hot drought arose as a synergistic alignment of environmental conditions and local forest structure that
509 allowed WPB to successfully colonize large trees, rapidly increase in population size, and expand. It follows
510 that the unexpectedly low mortality in cooler/wetter Sierra Nevada forests compared to model predictions
511 based on coarser-scale forest structure data (Young et al. 2017) could be explained by a different WPB
512 response to local forest structure due to a lack of an alignment with favorable climate conditions.

513 **Limitations and future directions**

514 We have demonstrated that drones can be effective means of collecting forest data at multiple, vastly different
515 spatial scales to investigate a single, multi-scale phenomenon– from meters in between trees, to hundreds
516 of meters of elevation, to hundreds of thousands of meters of latitude. Some limitations remain but can be

517 overcome with further refinements in the use of this tool for forest ecology. Most of these limitations arise
518 from tree detection and classification uncertainty, and thus it was imperative to work with field data for
519 calibration and uncertainty reporting.

520 The greatest limitation in our study arising from classification uncertainty is in the assumption that all dead
521 trees were ponderosa pine, which we estimate from coincident field plots is true approximately 73.4% of
522 the time. Because the forest structure factors influencing the likelihood of individual tree mortality during
523 the hot drought depended on tree species (Stephenson et al. 2019), we cannot rule out that some of the
524 ponderosa pine mortality relationships to forest structure that we observed may be partially explained by
525 those relationships in other species that were misclassified as ponderosa pine using our methods. However,
526 the overall community composition across our study area was similar (Fettig et al. 2019) and we are able
527 to reproduce similar forest structure/mortality patterns in drone-derived data when restricting the scope
528 of analysis to only trees detected in the footprints of the coincident field plots with dramatically different
529 patterns observed when including data from the forest surrounding the coincident field plots (see Supplemental
530 information). Thus, we remain confident that the patterns we observed were driven primarily by the dynamic
531 between WPB and ponderosa pine. While spectral information of foliage could help classify living trees to
532 species, the species of standing dead trees were not spectrally distinct. This challenge of classifying standing
533 dead trees to species implies that a conifer forest system with less bark beetle and tree host diversity, such
534 as mountain pine beetle outbreaks in monocultures of lodgepole pine in the Intermountain West, should be
535 particularly amenable to the methods presented here even with minimal further refinement because dead
536 trees will almost certainly belong to a single species and have succumbed to colonization by a single bark
537 beetle species.

538 Some uncertainty surrounded our ability to detect trees using the geometry of the dense point clouds derived
539 with SfM. The horizontal accuracy of the tree detection was better than the vertical accuracy, which may
540 result from a more significant error contribution by the field-based calculations of tree height compared to
541 tree position relative to plot center (Table 2). Both the horizontal and vertical accuracy would likely improve
542 with better SfM point clouds, which can be enhanced with greater overlap between images (Frey et al. 2018)
543 or with oblique (i.e., off-nadir) imagery (James and Robson 2014). Frey et al. (2018) found that 95% overlap
544 was preferable for generating dense point clouds in forested areas, and James and Robson (2014) reduced
545 dense point cloud errors using imagery taken at 30 degrees off-nadir. We only achieved 91.6% overlap with
546 the X3 RGB camera and 83.9% overlap with the multispectral camera, and all imagery was nadir-facing.
547 While our live/dead classification was fairly accurate (96.4% on a withheld dataset), our species classifier
548 would likely benefit from better crown segmentation because the pixel-level reflectance values within each

549 crown are averaged to characterize the “spectral signature” of each tree. With better delineation of each
550 tree crown, the mean value of pixels within each tree crown will likely be more representative of that tree’s
551 spectral signature. Better crown segmentation might most readily be achieved through greater overlap in
552 imagery. Finally, we anticipate that computer vision and deep learning will prove helpful in overcoming some
553 of these detection and classification challenges (Gray et al. 2019).

554 Conclusions

555 Climate change adaptation strategies emphasize management action that considers whole-ecosystem responses
556 to inevitable change (Millar et al. 2007), which requires a macroecological understanding of how phenomena
557 at multiple scales can interact. We’ve shown that drones can be a valuable tool for investigating multi-scalar
558 phenomena, such as how local forest structure combines with environmental conditions to shape forest insect
559 disturbance. Understanding the conditions that drive dry western U.S. forest responses to disturbances such
560 as bark beetle outbreaks will be vital for predicting outcomes from increasing disturbance frequency and
561 intensity exacerbated by climate change. Our study suggests that outcomes will depend on interactions
562 between local forest structure and broad-scale environmental gradients, with the potential for cross-scale
563 interactions to challenge our current understanding of forest insect dynamics.

564 References

- 565 Anderegg, W. R. L., J. A. Hicke, R. A. Fisher, C. D. Allen, J. Aukema, B. Bentz, S. Hood, J. W. Lichstein,
566 A. K. Macalady, N. McDowell, Y. Pan, K. Raffa, A. Sala, J. D. Shaw, N. L. Stephenson, C. Tague, and
567 M. Zeppel. 2015. Tree mortality from drought, insects, and their interactions in a changing climate. *New*
568 *Phytologist* 208:674–683.
- 569 Asner, G. P., P. G. Brodrick, C. B. Anderson, N. Vaughn, D. E. Knapp, and R. E. Martin. 2016. Progressive
570 forest canopy water loss during the 2012-2015 California drought. *Proceedings of the National Academy of*
571 *Sciences* 113:E249–E255.
- 572 Baldwin, B. G., A. H. Thornhill, W. A. Freyman, D. D. Ackerly, M. M. Kling, N. Morueta-Holme, and B. D.
573 Mishler. 2017. Species richness and endemism in the native flora of California. *American Journal of Botany*
574 104:487–501.
- 575 Bedard, W. D., P. E. Tilden, D. L. Wood, R. M. Silverstein, R. G. Brownlee, and J. O. Rodin. 1969.
576 Western pine beetle: Field response to its sex pheromone and a synergistic host terpene, myrcene. *Science*
577 164:1284–1285.

- 578 Bentz, B. J., J. Régnière, C. J. Fettig, E. M. Hansen, J. L. Hayes, J. A. Hicke, R. G. Kelsey, J. F. Negrón,
579 and S. J. Seybold. 2010. Climate change and bark beetles of the western United States and Canada: Direct
580 and indirect effects. BioScience 60:602–613.
- 581 Berryman, A. A. 1982. Population dynamics of bark beetles. Pages 264–314 in Bark Beetles in North
582 American Conifers: A System for the Study of Evolutionary Biology.
- 583 Boone, C. K., B. H. Aukema, J. Bohlmann, A. L. Carroll, and K. F. Raffa. 2011. Efficacy of tree defense
584 physiology varies with bark beetle population density: A basis for positive feedback in eruptive species.
585 Canadian Journal of Forest Research 41:1174–1188.
- 586 Brooks, S. P., and A. Gelman. 1998. General methods for monitoring convergence of iterative simulations.
587 Journal of Computational and Graphical Statistics 7:434.
- 588 Bürkner, P.-C. 2017. **brms**: An *R* package for bayesian multilevel models using *Stan*. Journal of Statistical
589 Software 80:1–28.
- 590 Byers, J. A., and D. L. Wood. 1980. Interspecific inhibition of the response of the bark beetles, *Dendroctonus*
591 *brevicomis* and *Ips paraconfusus*, to their pheromones in the field. Journal of Chemical Ecology 6:149–164.
- 592 Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker, J. Guo, P. Li,
593 and A. Riddell. 2017. Stan: A Probabilistic Programming Language. Journal of Statistical Software 76:1–32.
- 594 Chesson, P. 2000. Mechanisms of maintenance of species diversity. Annual Review of Ecology and Systematics
595 31:343–366.
- 596 Chubaty, A. M., B. D. Roitberg, and C. Li. 2009. A dynamic host selection model for mountain pine beetle,
597 *Dendroctonus ponderosae* Hopkins. Ecological Modelling 220:1241–1250.
- 598 Clevers, J., and A. Gitelson. 2013. Remote estimation of crop and grass chlorophyll and nitrogen content using
599 red-edge bands on Sentinel-2 and -3. International Journal of Applied Earth Observation and Geoinformation
600 23:344–351.
- 601 Coops, N. C., M. Johnson, M. A. Wulder, and J. C. White. 2006. Assessment of QuickBird high spatial
602 resolution imagery to detect red attack damage due to mountain pine beetle infestation. Remote Sensing of
603 Environment 103:67–80.
- 604 DJI. 2015a. Zenmuse X3 - Creativity Unleashed. <https://www.dji.com/zenmuse-x3/info>.
- 605 DJI. 2015b. DJI - The World Leader in Camera Drones/Quadcopters for Aerial Photography. <https://www.dji.com/matrice100/info>.

- 607 DronesMadeEasy. 2018. Map Pilot for DJI on iOS. [https://itunes.apple.com/us/app/map-pilot-for-dji/
608 id1014765000?mt=8](https://itunes.apple.com/us/app/map-pilot-for-dji/id1014765000?mt=8).
- 609 Evenden, M. L., C. M. Whitehouse, and J. Sykes. 2014. Factors influencing flight capacity of the mountain
610 pine beetle (Coleoptera: Curculionidae: Scolytinae). *Environmental Entomology* 43:187–196.
- 611 Eysn, L., M. Hollaus, E. Lindberg, F. Berger, J.-M. Monnet, M. Dalponte, M. Kobal, M. Pellegrini, E.
612 Lingua, D. Mongus, and N. Pfeifer. 2015. A benchmark of LiDAR-based single tree detection methods using
613 heterogeneous forest data from the alpine space. *Forests* 6:1721–1747.
- 614 Farr, T. G., P. A. Rosen, E. Caro, R. Crippen, R. Duren, S. Hensley, M. Kobrick, M. Paller, E. Rodriguez, L.
615 Roth, D. Seal, S. Shaffer, J. Shimada, J. Umland, M. Werner, M. Oskin, D. Burbank, and D. Alsdorf. 2007.
616 The shuttle radar topography mission. *Reviews of Geophysics* 45.
- 617 Fettig, C. J. 2012. Chapter 2: Forest health and bark beetles. *in* Managing Sierra Nevada Forests. PSW-
618 GTR-237. USDA Forest Service.
- 619 Fettig, C. J. 2016. Native bark beetles and wood borers in Mediterranean forests of California. Pages 499–528
620 *in* Insects and diseases of Mediterranean Forest systems. Springer International Publishing, Switzerland.
- 621 Fettig, C. J., and J. Hilszczański. 2015. Management strategies for bark beetles in conifer forests. Pages
622 555–584 *in* Bark Beetles. Elsevier.
- 623 Fettig, C. J., K. D. Klepzig, R. F. Billings, A. S. Munson, T. E. Nebeker, J. F. Negrón, and J. T. Nowak. 2007.
624 The effectiveness of vegetation management practices for prevention and control of bark beetle infestations in
625 coniferous forests of the western and southern United States. *Forest Ecology and Management* 238:24–53.
- 626 Fettig, C. J., S. R. McKelvey, C. P. Dabney, D. P. W. Huber, C. G. Lait, D. L. Fowler, and J. H. Borden. 2012.
627 Efficacy of “Verbenone Plus” for protecting ponderosa pine trees and stands from *Dendroctonus brevicomis*
628 (Coleoptera: Curculionidae) attack in British Columbia and California. *Journal of Economic Entomology*
629 105:1668–1680.
- 630 Fettig, C. J., S. R. McKelvey, and D. P. W. Huber. 2005. Nonhost angiosperm volatiles and Verbenone disrupt
631 response of western pine beetle, *Dendroctonus brevicomis* (Coleoptera: Scolytidae), to attractant-baited traps.
632 *Journal of Economic Entomology* 98:2041–2048.
- 633 Fettig, C. J., L. A. Mortenson, B. M. Bulaon, and P. B. Foulk. 2019. Tree mortality following drought in the
634 central and southern Sierra Nevada, California, U.S. *Forest Ecology and Management* 432:164–178.
- 635 Flint, L. E., A. L. Flint, J. H. Thorne, and R. Boynton. 2013. Fine-scale hydrologic modeling for regional land-

- 636 scape applications: The California Basin Characterization Model development and performance. Ecological
637 Processes 2:25.
- 638 Floyd, M. L., M. Clifford, N. S. Cobb, D. Hanna, R. Delph, P. Ford, and D. Turner. 2009. Relationship of
639 stand characteristics to drought-induced mortality in three Southwestern piñonJuniper woodlands. Ecological
640 Applications 19:1223–1230.
- 641 Franceschi, V. R., P. Krokene, E. Christiansen, and T. Krekling. 2005. Anatomical and chemical defenses of
642 conifer bark against bark beetles and other pests. New Phytologist 167:353–376.
- 643 Frey, J., K. Kovach, S. Stemmler, and B. Koch. 2018. UAV photogrammetry of forests as a vulnerable
644 process. A sensitivity analysis for a structure from motion RGB-image pipeline. Remote Sensing 10:912.
- 645 Fricker, G. A., N. W. Synes, J. M. Serra-Diaz, M. P. North, F. W. Davis, and J. Franklin. 2019. More than
646 climate? Predictors of tree canopy height vary with scale in complex terrain, Sierra Nevada, CA (USA).
647 Forest Ecology and Management 434:142–153.
- 648 Gabry, J., D. Simpson, A. Vehtari, M. Betancourt, and A. Gelman. 2019. Visualization in Bayesian workflow.
649 Journal of the Royal Statistical Society: Series A (Statistics in Society) 182:389–402.
- 650 Gitelson, A., and M. N. Merzlyak. 1994. Spectral reflectance changes associated with autumn senescence of
651 *Aesculus hippocastanum* L. And *Acer platanoides* L. Leaves. Spectral features and relation to chlorophyll
652 estimation. Journal of Plant Physiology 143:286–292.
- 653 Graf, M., M. Reid, B. Aukema, and B. Lindgren. 2012. Association of tree diameter with body size and lipid
654 content of mountain pine beetles. The Canadian Entomologist 144:467–477.
- 655 Gray, P. C., A. B. Fleishman, D. J. Klein, M. W. McKown, V. S. Bézy, K. J. Lohmann, and D. W. Johnston.
656 2019. A convolutional neural network for detecting sea turtles in drone imagery. Methods in Ecology and
657 Evolution 10:345–355.
- 658 Griffin, D., and K. J. Anchukaitis. 2014. How unusual is the 2012-2014 California drought? Geophysical
659 Research Letters 41:9017–9023.
- 660 Hayes, C. J., C. J. Fettig, and L. D. Merrill. 2009. Evaluation of multiple funnel traps and stand characteristics
661 for estimating western pine beetle-caused tree mortality. Journal of Economic Entomology 102:2170–2182.
- 662 Hijmans, R. J., J. van Etten, M. Sumner, J. Cheng, A. Bevan, R. Bivand, L. Busetto, M. Canty, D. Forrest,
663 A. Ghosh, D. Golicher, J. Gray, J. A. Greenberg, P. Hiemstra, I. for M. A. Geosciences, C. Karney, M.
664 Mattiuzzi, S. Mosher, J. Nowosad, E. Pebesma, O. P. Lamigueiro, E. B. Racine, B. Rowlingson, A. Shortridge,

- 665 B. Venables, and R. Wueest. 2019. *Raster*: Geographic data analysis and modeling.
- 666 Hoffman, M. D., and A. Gelman. 2014. The No-U-Turn Sampler: Adaptively setting path lengths in
667 Hamiltonian Monte Carlo. *Journal of Machine Learning Research* 15:31.
- 668 Hunziker, P. 2017. *Velox*: Fast raster manipulation and extraction.
- 669 Jactel, H., and E. G. Brockerhoff. 2007. Tree diversity reduces herbivory by forest insects. *Ecology Letters*
670 10:835–848.
- 671 Jakubowski, M. K., W. Li, Q. Guo, and M. Kelly. 2013. Delineating individual trees from LiDAR data: A
672 comparison of vector- and raster-based segmentation approaches. *Remote Sensing* 5:4163–4186.
- 673 James, M. R., and S. Robson. 2014. Mitigating systematic error in topographic models derived from UAV
674 and ground-based image networks. *Earth Surface Processes and Landforms* 39:1413–1420.
- 675 Kaiser, K. E., B. L. McGlynn, and R. E. Emanuel. 2013. Ecohydrology of an outbreak: Mountain pine beetle
676 impacts trees in drier landscape positions first: ECOHYDROLOGY OF A MOUNTAIN PINE BEETLE
677 OUTBREAK. *Ecohydrology* 6:444–454.
- 678 Kane, V. R., M. P. North, J. A. Lutz, D. J. Churchill, S. L. Roberts, D. F. Smith, R. J. McGaughey, J. T.
679 Kane, and M. L. Brooks. 2014. Assessing fire effects on forest spatial structure using a fusion of Landsat and
680 airborne LiDAR data in Yosemite National Park. *Remote Sensing of Environment* 151:89–101.
- 681 Kolb, T. E., C. J. Fettig, M. P. Ayres, B. J. Bentz, J. A. Hicke, R. Mathiasen, J. E. Stewart, and A. S. Weed.
682 2016. Observed and anticipated impacts of drought on forest insects and diseases in the United States. *Forest
683 Ecology and Management* 380:321–334.
- 684 Kuhn, M. 2008. Building predictive models in R using the caret package. *Journal of Statistical Software*
685 28:1–26.
- 686 Larson, A. J., and D. Churchill. 2012. Tree spatial patterns in fire-frequent forests of western North America,
687 including mechanisms of pattern formation and implications for designing fuel reduction and restoration
688 treatments. *Forest Ecology and Management* 267:74–92.
- 689 Li, W., Q. Guo, M. K. Jakubowski, and M. Kelly. 2012. A new method for segmenting individual trees from
690 the LiDAR point cloud. *Photogrammetric Engineering & Remote Sensing* 78:75–84.
- 691 Logan, J. A., P. White, B. J. Bentz, and J. A. Powell. 1998. Model analysis of spatial patterns in mountain
692 pine beetle outbreaks. *Theoretical Population Biology* 53:236–255.

- 693 Meyer, F., and S. Beucher. 1990. Morphological segmentation. *Journal of Visual Communication and Image*
694 *Representation* 1:21–46.
- 695 Micasense. 2015. MicaSense. <https://support.micasense.com/hc/en-us/articles/215261448-RedEdge-User->
696 *Manual-PDF-Download-*.
- 697 Millar, C. I., N. L. Stephenson, and S. L. Stephens. 2007. Climate change and forests of the future: Managing
698 in the face of uncertainty. *Ecological Applications* 17:2145–2151.
- 699 Millar, C. I., R. D. Westfall, D. L. Delany, M. J. Bokach, A. L. Flint, and L. E. Flint. 2012. Forest mortality in
700 high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern California, USA: Influence of environmental
701 context, bark beetles, climatic water deficit, and warming. *Canadian Journal of Forest Research* 42:749–765.
- 702 Miller, J. M., and F. P. Keen. 1960. Biology and control of the western pine beetle: A summary of the first
703 fifty years of research. US Department of Agriculture.
- 704 Moeck, H. A., D. L. Wood, and K. Q. Lindahl. 1981. Host selection behavior of bark beetles (Coleoptera:
705 Scolytidae) attacking *Pinus ponderosa*, with special emphasis on the western pine beetle, *Dendroctonus*
706 *brevicomis*. *Journal of Chemical Ecology* 7:49–83.
- 707 Morris, J. L., S. Cottrell, C. J. Fettig, W. D. Hansen, R. L. Sherriff, V. A. Carter, J. L. Clear, J. Clement, R.
708 J. DeRose, J. A. Hicke, P. E. Higuera, K. M. Mattor, A. W. R. Seddon, H. T. Seppä, J. D. Stednick, and S.
709 J. Seybold. 2017. Managing bark beetle impacts on ecosystems and society: Priority questions to motivate
710 future research. *Journal of Applied Ecology* 54:750–760.
- 711 Oliver, W. W. 1995. Is self-thinning in ponderosa pine ruled by *Dendroctonus* bark beetles? Page 6 *in* Forest
712 health through silviculture: Proceedings of the 1995 National Silviculture Workshop.
- 713 Pau, G., F. Fuchs, O. Sklyar, M. Boutros, and W. Huber. 2010. EBImage: An R package for image processing
714 with applications to cellular phenotypes. *Bioinformatics* 26:979–981.
- 715 Person, H. L. 1928. Tree selection by the western pine beetle. *Journal of Forestry* 26:564–578.
- 716 Person, H. L. 1931. Theory in explanation of the selection of certain trees by the western pine beetle. *Journal*
717 *of Forestry* 29:696–699.
- 718 Pile, L. S., M. D. Meyer, R. Rojas, O. Roe, and M. T. Smith. 2019. Drought impacts and compounding
719 mortality on forest trees in the southern Sierra Nevada. *Forests* 10:237.
- 720 Plowright, A. 2018. ForestTools: Analyzing remotely sensed forest data.
- 721 Raffa, K. F., B. H. Aukema, B. J. Bentz, A. L. Carroll, J. A. Hicke, M. G. Turner, and W. H. Romme. 2008.

- 722 Cross-scale drivers of natural disturbances prone to anthropogenic amplification: The dynamics of bark beetle
723 eruptions. BioScience 58:501–517.
- 724 Raffa, K. F., and A. A. Berryman. 1983. The role of host plant resistance in the colonization behavior and
725 ecology of bark beetles (Coleoptera: Scolytidae). Ecological Monographs 53:27–49.
- 726 Raffa, K. F., and A. A. Berryman. 1987. Interacting selective pressures in conifer-bark beetle systems: A
727 basis for reciprocal adaptations? The American Naturalist 129:234–262.
- 728 Raffa, K. F., J.-C. Grégoire, and B. Staffan Lindgren. 2015. Natural history and ecology of bark beetles.
729 Pages 1–40 in Bark Beetles. Elsevier.
- 730 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical
731 Computing, Vienna, Austria.
- 732 Restaino, C., D. Young, B. Estes, S. Gross, A. Wuenschel, M. Meyer, and H. Safford. 2019. Forest
733 structure and climate mediate drought-induced tree mortality in forests of the Sierra Nevada, USA. Ecological
734 Applications 0:e01902.
- 735 Robeson, S. M. 2015. Revisiting the recent California drought as an extreme value. Geophysical Research
736 Letters 42:6771–6779.
- 737 Rouse, W., R. H. Haas, W. Deering, and J. A. Schell. 1973. Monitoring the vernal advancement and
738 retrogradation (green wave effect) of natural vegetation. Type II Report, Goddard Space Flight Center,
739 Greenbelt, MD, USA.
- 740 Roussel, J.-R. 2019. lidRplugins: Extra functions and algorithms for lidR package.
- 741 Roussel, J.-R., D. Auty, F. De Boissieu, and A. S. Meador. 2019. lidR: Airborne LiDAR data manipulation
742 and visualization for forestry applications.
- 743 Safford, H. D., and J. T. Stevens. 2017. Natural range of variation for yellow pine and mixed-conifer forests
744 in the Sierra Nevada, Southern Cascades, and Modoc and Inyo National Forests, California, USA. Page 241.
- 745 dos Santos, A. A., J. Marcato Junior, M. S. Araújo, D. R. Di Martini, E. C. Tetila, H. L. Siqueira, C. Aoki, A.
746 Eltner, E. T. Matsubara, H. Pistori, R. Q. Feitosa, V. Liesenberg, and W. N. Gonçalves. 2019. Assessment of
747 CNN-Based Methods for Individual Tree Detection on Images Captured by RGB Cameras Attached to UAVs.
748 Sensors (Basel, Switzerland) 19.
- 749 Seidl, R., J. Müller, T. Hothorn, C. Bässler, M. Heurich, and M. Kautz. 2016. Small beetle, large-scale
750 drivers: How regional and landscape factors affect outbreaks of the European spruce bark beetle. The Journal

- 751 of applied ecology 53:530–540.
- 752 Seybold, S. J., B. J. Bentz, C. J. Fettig, J. E. Lundquist, R. A. Progar, and N. E. Gillette. 2018. Management
753 of western North American bark beetles with semiochemicals. Annual Review of Entomology 63:407–432.
- 754 Shepherd, W. P., D. P. W. Huber, S. J. Seybold, and C. J. Fettig. 2007. Antennal responses of the western
755 pine beetle, *Dendroctonus brevicomis* (Coleoptera: Curculionidae), to stem volatiles of its primary host,
756 *Pinus ponderosa*, and nine sympatric nonhost angiosperms and conifers. Chemoecology 17:209–221.
- 757 Shiklomanov, A. N., B. A. Bradley, K. M. Dahlin, A. M. Fox, C. M. Gough, F. M. Hoffman, E. M. Middleton,
758 S. P. Serbin, L. Smallman, and W. K. Smith. 2019. Enhancing global change experiments through integration
759 of remote-sensing techniques. Frontiers in Ecology and the Environment 0.
- 760 Shin, P., T. Sankey, M. Moore, and A. Thode. 2018. Evaluating unmanned aerial vehicle images for estimating
761 forest canopy fuels in a ponderosa pine stand. Remote Sensing 10:1266.
- 762 Stephenson, N. 1998. Actual evapotranspiration and deficit: Biologically meaningful correlates of vegetation
763 distribution across spatial scales. Journal of Biogeography 25:855–870.
- 764 Stephenson, N. L., A. J. Das, N. J. Ampersee, and B. M. Bulaon. 2019. Which trees die during drought?
765 The key role of insect host-tree selection. Journal of Ecology:75.
- 766 Stovall, A. E. L., H. Shugart, and X. Yang. 2019. Tree height explains mortality risk during an intense
767 drought. Nature Communications 10:1–6.
- 768 Thistle, H. W., H. Peterson, G. Allwine, B. Lamb, T. Strand, E. H. Holsten, and P. J. Shea. 2004. Surrogate
769 pheromone plumes in three forest trunk spaces: Composite statistics and case studies. Forest Science 50.
- 770 USDAFS. 2019, February 11. Press Release: Survey finds 18 million trees died in California in 2018.
771 https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/FSEPRD609321.pdf.
- 772 Vega, C., A. Hamrouni, S. El Mokhtari, J. Morel, J. Bock, J. P. Renaud, M. Bouvier, and S. Durrieu. 2014.
773 PTrees: A point-based approach to forest tree extraction from LiDAR data. International Journal of Applied
774 Earth Observation and Geoinformation 33:98–108.
- 775 Wallin, K. F., and K. F. Raffa. 2004. Feedback between Individual Host Selection Behavior and Population
776 Dynamics in an Eruptive Herbivore. Ecological Monographs 74:101–116.
- 777 Weinstein, B. G., S. Marconi, S. Bohlman, A. Zare, and E. White. 2019. Individual Tree-Crown Detection in
778 RGB Imagery Using Semi-Supervised Deep Learning Neural Networks. Remote Sensing 11:1309.
- 779 Wyngaard, J., L. Barbieri, A. Thomer, J. Adams, D. Sullivan, C. Crosby, C. Parr, J. Klump, S. Raj Shrestha,

- 780 and T. Bell. 2019. Emergent challenges for science sUAS data management: Fairness through community
781 engagement and best practices development. *Remote Sensing* 11:1797.
- 782 Young, D. J. N., J. T. Stevens, J. M. Earles, J. Moore, A. Ellis, A. L. Jirka, and A. M. Latimer. 2017.
783 Long-term climate and competition explain forest mortality patterns under extreme drought. *Ecology Letters*
784 20:78–86.
- 785 Zhang, W., J. Qi, P. Wan, H. Wang, D. Xie, X. Wang, and G. Yan. 2016. An easy-to-use airborne LiDAR
786 data filtering method based on cloth simulation. *Remote Sensing* 8:501.