

¹ **Cross-scale interaction of host tree size and climate governs bark
2 beetle-induced tree mortality**

³ Michael J. Koontz^{1,2,3*}, Andrew M. Latimer^{1,2}, Leif A. Mortenson⁴, Christopher J. Fettig⁵, Malcolm P.
⁴ North^{1,2,6}

⁵ ¹Graduate Group in Ecology, University of California, Davis, CA, USA

⁶ ²Department of Plant Sciences, University of California, Davis, CA, USA

⁷ ³Earth Lab, University of Colorado-Boulder; Boulder, CO, USA

⁸ ⁴USDA Forest Service, Pacific Southwest Research Station, Placerville, CA, USA

⁹ ⁵USDA Forest Service, Pacific Southwest Research Station, Davis, CA, USA

¹⁰ ⁶USDA Forest Service, Pacific Southwest Research Station, Mammoth Lakes, CA, USA

¹¹ *Correspondence: michael.koontz@colorado.edu

¹² *Keywords:* *Dendroctonus brevicomis*, disturbance, drones, *Pinus ponderosa*, Sierra Nevada, structure from motion, forest structure, climate change-type drought, macroecology

¹⁴ *Abstract word count:* 310

¹⁵ *Overall .docx word count:* 11499

¹⁶ *Main text word count:* 4034 (Intro: 1264 Results: 368 (41+110+217); Discussion: 2402)

¹⁷ *Methods word count:* 3093 (661+642+1463+327)

¹⁸ *Text boxes word count:* 0

¹⁹ Date report generated: March 08, 2020

²⁰ **Abstract**

²¹ The Californian hot drought of 2012 to 2015 created favorable conditions for unprecedented ponderosa pine
²² (*Pinus ponderosa*) mortality in the Sierra Nevada mountain range, largely attributable to the western pine
²³ beetle (*Dendroctonus brevicomis*; WPB). Climate conditions and forest density may interact to affect tree
²⁴ mortality, but density is a coarse gauge of forest structure that can affect WPB behavior in a number of
²⁵ ways. Measuring broad-scale climate conditions simultaneously with local forest composition and structure–
²⁶ the spatial distribution and size of trees– will refine our understanding of how these variables interact, but is
²⁷ generally expensive and/or labor-intensive. We use drone surveys over a network of 160 field plots along a 350–
²⁸ km latitudinal and 1000-m elevational gradient in western slope Sierra Nevada ponderosa pine/mixed-conifer

29 forests and structure from motion (SfM) processing to segment and classify more than 450,000 trees over 9
30 km² of forest with WPB-induced tree mortality. We modeled the probability of ponderosa pine mortality as a
31 function of forest structure and composition and their interaction with site-level climatic water deficit (CWD),
32 accounting for spatial covariance using exact Gaussian processes. A greater local proportion of host trees
33 strongly increased the probability of host mortality, with greater host density amplifying this effect. Further,
34 we found a strong interaction between host size and CWD such that larger trees increased the probability of
35 host mortality at hot/dry sites, but smaller trees tended to drive mortality in cool/wet sites.
36 Our results demonstrate a variable response of WPB to local forest structure and composition across an
37 environmental gradient, which may help reconcile differences between observed ecosystem-wide tree mortality
38 patterns and predictions from models based on coarser-scale forest structure. Climate change adaptation
39 strategies should consider that future disturbance outcomes may depend on interactions between local forest
40 structure and broad-scale environmental gradients, with the potential for cross-scale interactions that challenge
41 our current understanding of forest insect dynamics.

42 Introduction

43 Bark beetles dealt the final blow to many of the nearly 150 million trees killed in the California hot drought
44 of 2012 to 2015 and its aftermath (USDAFS 2019). A harbinger of climate change effects to come, record
45 high temperatures exacerbated the drought (Griffin and Anchukaitis 2014), which increased water stress in
46 trees (Asner et al. 2016), making them more susceptible to colonization by bark beetles (Fettig 2012, Kolb
47 et al. 2016). Further, a century of fire suppression policy has enabled forests to grow into dense stands,
48 which can also makes them more vulnerable to bark beetles (Fettig 2012). This combination of environmental
49 conditions and forest structural characteristics led to tree mortality events of unprecedented size in the
50 driest, densest forests across the state (Young et al. 2017). The mechanisms underlying the link between
51 tree susceptibility to colonization by insects and hot, dry conditions are often directly attributed to tree
52 physiology (Bentz et al. 2010, Kolb et al. 2016), while the link to forest density is multifaceted (Fettig 2012).
53 Because forest density is a coarse metric of the forest features to which bark beetles respond (Raffa et al.
54 2008), our understanding of the connection between forest density and insect disturbance severity could
55 be enhanced with more finely-resolved measures of forest structure as well as explicit consideration of tree
56 species composition (Stephenson et al. 2019, Fettig et al. 2019). Finally, the challenge of simultaneously
57 measuring the effects of both local-scale forest features (such as structure and composition) and broad-scale
58 environmental conditions (such as climatic water deicit; CWD) on forest insect disturbance leaves their
59 interaction effect relatively underexplored (Seidl et al. 2016, Stephenson et al. 2019, Fettig et al. 2019).

60 The ponderosa pine/mixed-conifer forests in California's Sierra Nevada region are characterized by regular
61 bark beetle disturbances, primarily by the influence of western pine beetle (*Dendroctonus brevicomis*; WPB)
62 on its host ponderosa pine (*Pinus ponderosa*) (Fettig 2016). WPB is a "primary" bark beetle— its reproductive
63 success is contingent upon host tree mortality, which itself requires enough beetles to "mass attack" the
64 host tree and overwhelm its defenses (Raffa and Berryman 1983). This Allee effect creates a strong coupling
65 between beetle selection behavior of host trees and host tree susceptibility to colonization (Raffa and Berryman
66 1983, Logan et al. 1998). A key defense mechanism of conifers to bark beetle attack is to flood beetle bore
67 holes with resin, which physically expels beetles and may interrupt beetle communication (Franceschi et al.
68 2005, Raffa et al. 2015). Under normal conditions, weakened trees with compromised defenses are the most
69 susceptible to colonization and will be the main targets of primary bark beetles like WPB (Bentz et al. 2010,
70 Raffa et al. 2015). Under severe water stress, many trees no longer have the resources available to mount
71 a defense (Kolb et al. 2016) and thus prolonged drought can often trigger increased bark beetle-induced
72 tree mortality as average tree vigor declines (Bentz et al. 2010). As the local population density of beetles
73 increases due to successful reproduction within spatially-aggregated weakened trees, as might occur during
74 drought, mass attacks grow in size and become capable of overwhelming formidable tree defenses such that
75 even healthy trees may be susceptible to colonization and mortality (Bentz et al. 2010, Raffa et al. 2015).
76 Thus, water stress can be a key determinant of whether individual trees are susceptible to bark beetles under
77 many conditions, and this environmental condition may interact with beetle population dynamics to drive
78 tree susceptibility under extreme conditions (Bentz et al. 2010, Stephenson et al. 2019).

79 WPB activity is strongly influenced by forest structure— the spatial distribution and size of trees— and tree
80 species composition. Taking forest structure alone, high-density forests are more prone to bark beetle-induced
81 tree mortality (Fettig 2012) which may arise as greater competition for water resources amongst crowded trees
82 and thus average tree resistance is lower (Hayes et al. 2009), or because smaller gaps between trees protect
83 pheromone plumes from dissipation by the wind and thus enhance intraspecific beetle communication (Thistle
84 et al. 2004). Tree size is another aspect of forest structure that affects bark beetle host selection behavior
85 with smaller trees tending to have lower capacity for resisting attack, and larger trees being more desirable
86 targets on account of their thicker phloem providing greater nutritional content (Chubaty et al. 2009, Graf et
87 al. 2012). Taking forest composition alone, WPB activity in the Sierra Nevada mountain range of California
88 is necessarily tied to the regional distribution of its exclusive host, ponderosa pine (Fettig 2016). Colonization
89 by primary bark beetles can also depend on the relative frequencies of tree species in a more local area, akin
90 to reduced oligophagous insect herbivory in forests comprising taxonomically-distinct tree species compared
91 to monocultures (Jactel and Brockerhoff 2007). The interaction between forest structure and composition

92 also drives WPB activity. For instance, high-density forests with high host availability may experience greater
93 beetle-induced tree mortality because dispersal distances between potential host trees are shorter reducing
94 predation of adults searching for hosts and facilitating higher rates of colonization (Miller and Keen 1960,
95 Berryman 1982, Fettig et al. 2007) or because high host availability reduces the chance of individual beetles
96 wasting their limited resources flying to and landing on a non-host tree (Moeck et al. 1981, Evenden et
97 al. 2014). Stand-scale measures of forest structure and composition thus paint a fundamentally limited
98 picture of the mechanisms by which these forest characteristics affect bark beetle disturbance, but finer-grain
99 information explicitly recognizing tree size, tree species, and local tree density should more appropriately
100 capture the ecological processes underlying insect-induced tree mortality. Additionally, considering the
101 effects of local forest structure and composition with the effects of environmental conditions may help refine
102 our understanding of tree mortality patterns in widespread events such as during the recent California hot
103 drought.

104 The vast spatial extent of tree mortality in the 2012 to 2015 California hot drought challenges our ability to
105 simultaneously consider how broad-scale environmental conditions may interact with local forest structure
106 and composition to affect the dynamic between bark beetle selection and colonization of host trees, and host
107 tree susceptibility to attack (Anderegg et al. 2015, Stephenson et al. 2019). Measuring local forest structure
108 generally requires expensive instrumentation (Kane et al. 2014, Asner et al. 2016) or labor-intensive field
109 surveys (Larson and Churchill 2012, Stephenson et al. 2019, Fettig et al. 2019), which constrains survey
110 extent and frequency. Small, unhumanned aerial systems (sUAS) enable relatively fast and cheap remote
111 imaging over dozens of hectares of forest, which can be used to measure complex forest structure at the
112 individual tree scale (Morris et al. 2017, Shiklomanov et al. 2019). Distributing such surveys across an
113 environmental gradient can overcome the data acquisition challenge inherent in investigating phenomena
114 with both a strong local- and a strong broad-scale component.

115 We used ultra-high resolution, sUAS-derived remote sensing data over a network of 32 sites in Sierra Nevada
116 ponderosa pine/mixed-conifer forests spanning 1000 m of elevation and 350 km of latitude (see Fettig et
117 al. 2019) and covering a total of 9 km² to ask how broad-scale environmental conditions interacted with
118 local forest structure and composition to affect the probability of tree mortality during the cumulative tree
119 mortality event of 2012 to 2018. We asked:

- 120 1. How does the proportion of host trees in a local area and average host tree size affect WPB-induced
121 tree mortality?
- 122 2. How does the density of all tree species (hereafter “overall density”) affect WPB-induced tree mortality?

- 123 3. How does environmentally-driven tree moisture stress affect WPB-induced tree mortality?
- 124 4. Do the effects of forest structure, forest composition, and environmental condition interact to influence
- 125 WPB-induced tree mortality?

126 **Methods**

127 **Study system**

128 We built our study coincident with 160 vegetation/forest insect monitoring plots at 32 sites established
129 between 2016 and 2017 by Fettig et al. (2019) (Figure 1). The study sites were chosen to reflect typical
130 west-side Sierra Nevada yellow pine/mixed-conifer forests and were dominated by ponderosa pine (Fettig
131 et al. 2019). Plots were located in WPB-attacked, yellow pine/mixed-conifer forests across the Eldorado,
132 Stanislaus, Sierra and Sequoia National Forests and were stratified by elevation (914-1219 m, 1219-1524
133 m, 1524-1829 m above sea level). In the Sequoia National Forest, the southernmost National Forest in our
134 study, plots were stratified with the lowest elevation band of 1219-1524 m and extended to an upper elevation
135 band of 1829-2134 m to capture a more similar forest community composition as at the more northern
136 National Forests. The sites have variable forest structure and plot locations were selected in areas with >35%
137 ponderosa pine basal area and >10% ponderosa pine mortality. At each site, five 0.041 ha circular plots
138 were installed along transects with 80 to 200m between plots. In the field, Fettig et al. (2019) mapped all
139 stem locations relative to the center of each plot using azimuth/distance measurements. Tree identity to
140 species, tree height, and diameter at breast height (DBH) were recorded if DBH was greater than 6.35cm.
141 Year of mortality was estimated based on needle color and retention if it occurred prior to plot establishment,
142 and was directly observed thereafter during annual site visits. A small section of bark (approximately 625
143 cm²) on both north and south aspects was removed from dead trees to determine if bark beetle galleries
144 were present. The shape, distribution, and orientation of galleries are commonly used to distinguish among
145 bark beetle species (Fettig 2016). In some cases, deceased bark beetles were present beneath the bark to
146 supplement identifications based on gallery formation. During the spring and early summer of 2018, all field
147 plots were revisited to assess whether dead trees had fallen (Fettig et al. 2019).

148 In the typical life cycle of WPBs, females initiate host colonization by tunneling through the outer bark and
149 into the phloem and outer xylem where they rupture resin canals.

150 As a result, oleoresin exudes and collects on the bark surface, as is commonly observed with other bark beetle
151 species. During the early stages of attack, females release an aggregation pheromone component which, in
152 combination with host monoterpenes released from pitch tubes, is attractive to conspecifics (Bedard et al.





Figure 1: The network of field plots spanned a 350-km latitudinal gradient from the Eldorado National Forest in the north to the Sequoia National Forest in the south. Plots were stratified by three elevation bands in each forest, with the plots in the Sequoia National Forest (the southern-most National Forest) occupying elevation bands 305 m above the three bands in the other National Forests in order to capture a similar community composition.

153 1969). An antiaggregation pheromone component is produced during latter stages of host colonization by
154 several pathways, and is thought to reduce intraspecific competition by altering adult behavior to minimize
155 overcrowding of developing brood within the host (Byers and Wood 1980). Volatiles from several nonhosts
156 sympatric with ponderosa pine have been demonstrated to inhibit attraction of WPB (Shepherd et al. 2007,
157 Fettig and Hilszczański 2015). In California, WPB generally has 2-3 generations in a single year and can
158 often out-compete its congener, the mountain pine beetle, *Dendroctonus ponderosae*, in ponderosa pines,
159 especially in larger trees (Miller and Keen 1960).

160 **Aerial data collection and processing**

161 Nadir-facing imagery was captured using a gimbal-stabilized DJI Zenmuse X3 broad-band red/green/blue
162 (RGB) camera (DJI 2015a) and a fixed-mounted Micasense Rededge3 multispectral camera with five narrow
163 bands (Micasense 2015) on a DJI Matrice 100 aircraft (DJI 2015b). Imagery was captured from both cameras
164 along preprogrammed aerial transects over ~40 hectares surrounding each of the 32 sites (each of these
165 containing five field plots) and was processed in a series of steps to yield local forest structure and composition
166 data suitable for our statistical analyses. Following the call by Wyngaard et al. (2019), we establish “data
167 product levels” to reflect the image processing pipeline from raw imagery (Level 0) to calibrated, fine-scale
168 forest structure and composition information on regular grids (Level 4), with each new data level derived
169 from levels below it. Here, we outline the steps in the processing and calibration pipeline visualized in Figure
170 2, and include additional details in the Supplemental Information.


Level 0: raw data from sensors




Level 1: basic outputs from photogrammetric processing




Level 2: corrected outputs from photogrammetric processing



Level 3: domain-specific information extraction



Level 4: aggregations to regular grids



172 Figure 2. Schematic of the data processing workflow for a single site with each new data product level derived
173 from data at lower levels.

174 Level 0 represents raw data from the sensors. From left to right: example broad-band RGB photo from
175 DJI Zenmuse X3 camera, example blue photo from Rededge3 (centered on 475nm), example green photo
176 from Rededge3 (centered on 560nm), example red photo from Rededge3 (centered on 668nm), example near
177 infrared photo from Rededge3 (centered on 840nm), and example red edge photo from Rededge3 (centered on
178 717nm).

179 Level 1 represents basic outputs from the photogrammetric workflow, in this case implemented with
180 Pix4Dmapper. From left to right: a dense point cloud visualized in CloudCompare (<https://www.danielgm.net/cc/>), an orthophoto generated from the RGB camera, and a digital surface model representing the
181 altitude above sea level (ground height + vegetation height) for every cell.

183 Level 2 represents outputs from photogrammetric processing that have been corrected radiometrically or
184 geometrically. From left to right: a radiometrically-corrected surface reflectance map of the red narrow band
185 from the Rededge3 camera, a radiometrically-corrected surface reflectance map of the near infrared narrow
186 band from the Rededge3 camera, a rasterized version of the digital terrain model derived by a geometric
187 correction of the dense point cloud, and a canopy height model derived by subtracting the terrain height
188 from the digital surface model.

189 Level 3 represents domain-specific information extraction from Level 2 products and is divided into two
190 sub-levels. Level 3a products are derived using only spectral or only geometric data. From left to right: a
191 reflectance map of Normalized Difference Vegetation Index (NDVI; Rouse et al. (1973)) derived using the red
192 and near infrared Level 2 reflectance products, a map of points representing detected trees from the canopy
193 height model with a red polygon highlighting the area presented in more detail for the next two images, a
194 close-up of points representing detected trees, and a close-up of polygons representing segmented tree crowns.
195 Level 3b products are derived using both spectral and geometric data. From left to right: a map of the point
196 locations of detected trees that have been classified as alive or dead based on the pixel values within each
197 segmented tree crown and a map of the point locations of detected trees classified to WPB host/non-host
198 using the same spectral information. Note that our study relies on the generation of Level 3a products in
199 order to combine them and create Level 3b products, but this need not be the case. For instance, deep
200 learning/neural net methods may be able to use both the spectral and geometric information from Level 2
201 simultaneously to locate and classify trees in a scene and directly generate Level 3b products without a need
202 to first generate the Level 3a products shown in this schematic (Weinstein et al. 2019, dos Santos et al. 2019).

203 Level 4 represents aggregations of Level 3 products to regular grids which might better reflect the grain size
204 of the data for which we have the best calibration and thus the most confidence or which might provide
205 new information not possible at an individual-tree level (e.g., average distance between trees in a small
206 neighborhood). From left to right: aggregation of live/dead classified trees as fraction of dead trees in a 20 x
207 20-m cell, aggregation of host/non-host classified trees as fraction of hosts in a 20 x 20-m cell, aggregation of
208 mean host height in a 20 x 20-m cell, and aggregation of tree count (including all species), in a 20 x 20-m
209 cell. In our case, the 20 x 20-m aggregation produces a grid cell with an area of 400 m², which most closely
210 matches the 404-m² area of the ground-based vegetation plots whose data we used in an aggregated form to
211 calibrate our derivation of Level 3 products.

212 **Level 0: Raw data from sensors**

213 Raw data comprised approximately 1900 images per camera lens (one broad-band RGB lens and five narrow-
214 band multispectral lenses) for each of the 32 sites (Figure 2; Level 0). Prior to the aerial survey, two strips of
215 bright orange drop cloth (~100 x 15 cm) were positioned as an “X” over the permanent monuments marking
216 the center of the 5 field plots from Fettig et al. (2019) (see Supplemental Information).

217 We preprogrammed north-south aerial transects using Map Pilot for DJI on iOS flight software (Drones-
218 MadeEasy 2018) at an altitude of 120 m above ground level (with “ground” defined using a 1-arc-second
219 digital elevation model (Farr et al. 2007)). The resulting ground sampling distance was approximately 5
220 cm/px for the Zenmuse X3 RGB camera and approximately 8 cm/px for the Rededge3 multispectral camera.
221 We used 91.6% image overlap (both forward and side) at the ground for the Zenmuse X3 RGB camera and
222 83.9% overlap (forward and side) for the Rededge3 multispectral camera.

223 **Level 1: Basic outputs from photogrammetric processing**

224 We used SfM photogrammetry implemented in Pix4Dmapper Cloud (www.pix4d.com) to generate dense point
225 clouds (Figure 2; Level 1, left), orthophotos (Figure 2; Level 1, center), and digital surface models (Figure 2;
226 Level 1, right) for each field site (Frey et al. 2018). For 29 sites, we processed the Rededge3 multispectral
227 imagery alone to generate these products. For three sites, we processed the RGB and the multispectral
228 imagery together to enhance the point density of the dense point cloud. All SfM projects resulted in a single
229 processing “block,” indicating that all images in the project were optimized and processed together. The
230 dense point cloud represents x, y, and z coordinates as well as the color of millions of points per site. The
231 orthophoto represents a radiometrically uncalibrated, top-down view of the survey site that preserves the

232 relative x-y positions of objects in the scene. The digital surface model is a rasterized version of the dense
233 point cloud that shows the altitude above sea level for each pixel in the scene at the ground sampling distance
234 of the camera that generated the Level 0 data.

235 **Level 2: Corrected outputs from photogrammetric processing**

236 **Radiometric corrections**

237 A radiometrically-corrected reflectance map (Figure 2; Level 2, left two figures; i.e., a corrected version of the
238 Level 1 orthophoto) was generated using the Pix4D software by incorporating incoming light conditions for
239 each narrow band of the Rededge3 camera (captured simultaneously with the Rededge3 camera using an
240 integrated downwelling light sensor) as well as a pre-flight image of a calibration panel of known reflectance
241 (see Supplemental Information for camera and calibration panel details).

242 **Geometric corrections**

243 We implemented a geometric correction to the Level 1 dense point cloud and digital surface model by
244 normalizing these data for the terrain underneath the vegetation. We generated the digital terrain model
245 representing the ground underneath the vegetation at 1-m resolution (Figure 2; Level 2, third image) by
246 classifying each survey area's dense point cloud into "ground" and "non-ground" points using a cloth simulation
247 filter algorithm (Zhang et al. 2016) implemented in the `lidR` (Roussel et al. 2019) package and rasterizing
248 the ground points using the `raster` package (Hijmans et al. 2019). We generated a canopy height model
249 (Figure 2; Level 2, fourth image) by subtracting the digital terrain model from the digital surface model.

250 **Level 3: Domain-specific information extraction**

251 **Level 3a: Data derived from spectral OR geometric Level 2 product**

252 Using just the spectral information from the radiometrically-corrected reflectance maps, we calculated several
253 vegetation indices including the normalized difference vegetation index (NDVI; Rouse et al. (1973); Figure
254 2; Level 3a, first image), the normalized difference red edge (NDRE; Gitelson and Merzlyak (1994)), the
255 red-green index (RGI; Coops et al. (2006)), the red edge chlorophyll index ($CI_{red\ edge}$; Clevers and Gitelson
256 (2013)), and the green chlorophyll index (CI_{green} ; Clevers and Gitelson (2013)).

Table 1: Algorithm name, number of parameter sets tested for each algorithm, and references.

Algorithm	Parameter sets tested	Reference(s)
li2012	131	Li et al. (2012); Jakubowski et al. (2013); Shin et al. (2018)
lmfx	30	Roussel (2019)
localMaxima	6	Roussel et al. (2019)
multichm	1	Eysn et al. (2015)
ptrees	3	Vega et al. (2014)
vwf	3	Plowright (2018)
watershed	3	Pau et al. (2010)

257 Using just the geometric information from the canopy height model or terrain-normalized dense point cloud,
 258 we generated maps of detected trees (Figure 2; Level 3a, second and third images) by testing a total of 7
 259 automatic tree detection algorithms and a total of 177 parameter sets (Table 1). We used the field plot data
 260 to assess each tree detection algorithm/parameter set by converting the distance-from-center and azimuth
 261 measurements of the trees in the field plots to x-y positions relative to the field plot centers distinguishable in
 262 the Level 2 reflectance maps as the orange fabric X’s that we laid out prior to each flight. In the reflectance
 263 maps, we located 110 out of 160 field plot centers while some plot centers were obscured due to dense
 264 interlocking tree crowns or because a plot center was located directly under a single tree crown. For each of
 265 the 110 field plots with identifiable plot centers— the “validation field plots”, we calculated 7 forest structure
 266 metrics using the ground data collected by Fettig et al. (2019): total number of trees, number of trees greater
 267 than 15 m in height, mean height of trees, 25th percentile tree height, 75th percentile tree height, mean
 268 distance to nearest tree neighbor, and mean distance to second nearest neighbor. For each tree detection
 269 algorithm and parameter set described above, we calculated the same set of 7 structure metrics within the
 270 footprint of the validation field plots. We calculated the Pearson’s correlation and root mean square error
 271 (RMSE) between the ground data and the aerial data for each of the 7 structure metrics for each of the 177
 272 automatic tree detection algorithms/parameter sets. For each algorithm and parameter set, we calculated its
 273 performance relative to other algorithms as whether its Pearson’s correlation was within 5% of the highest
 274 Pearson’s correlation as well as whether its RMSE was within 5% of the lowest RMSE. We summed the
 275 number of forest structure metrics for which it reached these 5% thresholds for each algorithm/parameter
 276 set. For automatically detecting trees across the whole study, we selected the algorithm/parameter set that

277 performed well across the most number of forest metrics (see Results).

278 We delineated individual tree crowns (Figure 2; Level 3a, fourth image) with a marker controlled watershed
279 segmentation algorithm (Meyer and Beucher 1990) implemented in the **ForestTools** package (Plowright
280 2018) using the detected treetops as markers. If the automatic segmentation algorithm failed to generate
281 a crown segment for a detected tree (e.g., often snags with a very small crown footprint), a circular crown
282 was generated with a radius of 0.5 m. If the segmentation generated multiple polygons for a single detected
283 tree, only the polygon containing the detected tree was retained. Because image overlap decreases near the
284 edges of the overall flight path and reduces the quality of the SfM processing in those areas, we excluded
285 segmented crowns within 35 m of the edge of the survey area. Given the narrower field of view of the
286 Rededge3 multispectral camera versus the X3 RGB camera whose optical parameters were used to define the
287 ~40 hectare survey area around each site, as well as the 35 m additional buffering, the survey area at each
288 site was ~30 ha (see Supplemental Information).

289 **Level 3b: Data derived from spectral AND geometric information**

290 We overlaid the segmented crowns on the reflectance maps from 20 sites spanning the latitudinal and elevation
291 gradient in the study. Using QGIS (<https://qgis.org/en/site/>), we hand classified 564 trees as live/dead
292 (Figure 3) and as one of 5 dominant species in the study area (ponderosa pine, *Pinus lambertiana*, *Abies*
293 *concolor*, *Calocedrus decurrens*, or *Quercus kelloggii*) using the mapped ground data as a guide. Each tree was
294 further classified as “host” for ponderosa pine or “non-host” for all other species (Fettig 2016). We extracted
295 all the pixel values within each segmented crown polygon from the five, Level 2 orthorectified reflectance
296 maps (one per narrow band on the Rededge3 camera) as well as from the five, Level 3a vegetation index
297 maps using the **velox** package (Hunziker 2017). For each crown polygon, we calculated the mean value of
298 the extracted Level 2 and Level 3a pixels and used them as ten independent variables in a five-fold cross
299 validated boosted logistic regression model to predict whether the hand classified trees were alive or dead.
300 For just the living trees, we similarly used all 10 mean reflectance values per crown polygon to predict tree
301 species using a five-fold cross validated regularized discriminant analysis. The boosted logistic regression and
302 regularized discriminant analysis were implemented using the **caret** package in R (Kuhn 2008). Finally, we
303 used these models to classify all tree crowns in the data set as alive or dead (Figure 2; Level 3b, first image)
304 as well as the species of living trees (Figure 2; Level 3b, second image).

305 **Level 4: Aggregations to regular grids**

306 We rasterized the forest structure and composition data at a spatial resolution similar to that of the field
307 plots to better match the grain size at which we validated the automatic tree detection algorithms. In each
308 raster cell, we calculated: number of dead trees, number of ponderosa pine trees, total number of trees, and
309 mean height of ponderosa pine trees. The values of these variables in each grid cell and derivatives from
310 them were used for visualization and modeling. Here, we show the fraction of dead trees per cell (Figure 2;
311 Level 4, first image), the fraction of host trees per cell (Figure 2; Level 4, second image), the mean height of
312 ponderosa pine trees in each cell (Figure 2; Level 4, third image), and the total count of trees per cell (Figure
313 2; Level 4, fourth image).

314 **Note on assumptions about dead trees**

315 For the purposes of this study, we assumed that all dead trees were ponderosa pine and thus hosts colonized
316 by WPB. This is a reasonably good assumption for our study area; for example, Fettig et al. (2019) found
317 that 73.4% of dead trees in their coincident field plots were ponderosa pine. Mortality was concentrated in
318 the larger-diameter classes and attributed primarily to WPB (see Figure 5 of Fettig et al. 2019). The species
319 contributing to the next highest proportion of dead trees was incense cedar which represented 18.72% of the
320 dead trees in the field plots. While the detected mortality is most likely to be ponderosa pine killed by WPB,
321 it is critical to interpret our results with these limitations in mind.

322 **Environmental data**

323 We used CWD (Stephenson 1998) from the 1981-2010 mean value of the basin characterization model (Flint
324 et al. 2013) as an integrated measure of temperature and moisture conditions for each of the 32 sites. Higher
325 values of CWD correspond to hotter, drier conditions and lower values correspond to cooler, wetter conditions.
326 CWD has been shown to correlate well with broad patterns of tree mortality in the Sierra Nevada (Young et
327 al. 2017) as well as bark beetle-induced tree mortality (Millar et al. 2012). We converted the CWD value for
328 each site into a z-score representing that site's deviation from the mean CWD across the climatic range of
329 Sierra Nevada ponderosa pine as determined from 179 herbarium records described in Baldwin et al. (2017).
330 Thus, a CWD z-score of 1 would indicate that the CWD at that site is one standard deviation hotter/drier
331 than the mean CWD across all geolocated herbarium records for ponderosa pine in the Sierra Nevada.

332 **Statistical model**

333 We used a generalized linear model with a zero-inflated binomial response and a logit link to predict the
 334 probability of ponderosa pine mortality within each 20 x 20-m cell using the total number of ponderosa
 335 pine trees in each cell as the number of trials, and the number of dead trees in each cell as the number of
 336 “successes”. As covariates, we used the proportion of trees that are WPB hosts (i.e., ponderosa pine) in each
 337 cell, the mean height of ponderosa pine trees in each cell, the count of trees of all species (overall density) in
 338 each cell, and the site-level CWD using Eq. 1. Note that the two-way interaction between the overall density
 339 and the proportion of trees that are hosts is equivalent to the number of ponderosa pine trees in the cell.
 340 To measure and account for spatial autocorrelation underlying ponderosa pine mortality, we subsampled
 341 the data at each site to a random selection of 200, 20 x 20-m cells representing approximately 27.5% of the
 342 surveyed area. Additionally with these subsampled data, we included a separate exact Gaussian process term
 343 per site of the interaction between the x- and y-position of each cell using the `gp()` function in the `brms`
 344 package (Bürkner 2017). The Gaussian process estimates the spatial covariance in the response variable
 345 (log-odds of ponderosa pine mortality) jointly with the effects of the other covariates.

$$y_{i,j} \sim \begin{cases} 0, & p \\ Binom(n_i, \pi_i), & 1 - p \end{cases}$$

$$\begin{aligned} logit(\pi_i) = \beta_0 + \\ \beta_1 X_{cwd,j} + \beta_2 X_{propHost,i} + \beta_3 X_{PIPOheight,i} + \beta_4 X_{overallDensity,i} + \\ \beta_5 X_{cwd,j} X_{PIPOheight,i} + \beta_6 X_{cwd,j} X_{propHost,i} + \beta_7 X_{cwd,j} X_{overallDensity,i} + \\ \beta_8 X_{propHost,i} X_{PIPOheight,i} + \beta_9 X_{propHost,i} X_{overallDensity,i} + \\ \beta_{10} X_{cwd,j} X_{propHost,i} X_{PIPOheight,i} + \\ \mathcal{GP}_j(x_i, y_i) \end{aligned}$$

346 Where y_i is the number of dead trees in cell i , n_i is the sum of the dead trees (assumed to be ponderosa pine)
 347 and live ponderosa pine trees in cell i , π_i is the probability of ponderosa pine tree mortality in cell i , p is
 348 the probability of there being zero dead trees in a cell arising as a result of an unmodeled process, $X_{cwd,j}$
 349 is the z-score of CWD for site j , $X_{propHost,i}$ is the scaled proportion of trees that are ponderosa pine in
 350 cell i , $X_{PIPOheight,i}$ is the scaled mean height of ponderosa pine trees in cell i , $X_{overallDensity,i}$ is the scaled
 351 density of all trees in cell i , x_i and y_i are the x- and y- coordinates of the centroid of the cell in an EPSG3310
 352 coordinate reference system, and \mathcal{GP}_j represents the exact Gaussian process describing the spatial covariance

353 between cells at site j .

354 We used 4 chains with 4000 iterations each (2000 warmup, 2000 samples), and confirmed chain convergence
355 by ensuring all R_{hat} values were less than 1.1 (Brooks and Gelman 1998) and that the bulk and tail effective
356 sample sizes (ESS) for each estimated parameter were greater than 100 times the number of chains (i.e.,
357 greater than 400 in our case). We used posterior predictive checks to visually confirm model performance by
358 overlaying the density curves of the predicted number of dead trees per cell over the observed number (Gabry
359 et al. 2019). For the posterior predictive checks, we used 50 random samples from the model fit to generate
360 50 density curves and ensured curves were centered on the observed distribution, paying special attention to
361 model performance at capturing counts of zero.

362 Software and data availability

363 All data are available via the Open Science Framework. Statistical analyses were performed using the **brms**
364 packages. With the exception of the SfM software (Pix4Dmapper Cloud) and the GIS software QGIS, all
365 data carpentry and analyses were performed using R (R Core Team 2018).

366 Results

367 Tree detection algorithm performance

368 We found that the experimental **lmfx** algorithm with parameter values of **dist2d = 1** and **ws = 2.5** (Roussel
369 et al. 2019) performed the best across 7 measures of forest structure as measured by Pearson's correlation
370 with ground data (Table 2).

Table 2: Correlation and differences between the best performing tree detection algorithm (**lmfx** with **dist2d = 1** and **ws = 2.5**) and the ground data. An asterisk next to the correlation or RMSE indicates that this value was within 5% of the value of the best-performing algorithm/parameter set. Ground mean represents the mean value of the forest metric across the 110 field plots that were visible from the sUAS-derived imagery. The median error is calculated as the median of the differences between the air and ground values for the 110 visible plots. Thus, a positive number indicates an overestimate by the sUAS workflow and a negative number indicates an underestimate.

Forest structure metric	Ground mean	Correlation with ground	RMSE	Median error
total tree count	19	0.67*	8.68*	2
count of trees > 15 m	9.9	0.43	7.38	0
distance to 1st neighbor (m)	2.8	0.55*	1.16*	0.26

Forest structure metric	Ground mean	Correlation with ground	RMSE	Median error
distance to 2nd neighbor (m)	4.3	0.61*	1.70*	0.12
height (m); 25 th percentile	12	0.16	8.46	-1.2
height (m); mean	18	0.29	7.81*	-2.3
height (m); 75 th percentile	25	0.35	10.33*	-4

371 **Classification accuracy for live/dead and host/non-host**

372 The accuracy of live/dead classification on a withheld test dataset was 97.3%. The accuracy of species
 373 classification on a withheld testing dataset was 66.7%. The accuracy of WPB host/non-WPB-host (i.e.,
 374 ponderosa pine versus other tree species) on a withheld testing dataset was 74.4%.

375 **Site summary based on best tree detection algorithm and classification**

376 Across all study sites, we detected, segmented, and classified 452,413 trees (see Supplemental Information for
 377 site summaries). Of these trees, we classified 118,879 as dead (26.3% mortality). Estimated site-level tree
 378 mortality ranged from 6.8% to 53.6%.

379 **Effect of local structure and regional climate on tree mortality attributed to western pine
 380 beetle**

381 We detected a positive main effect of CWD on the probability of ponderosa pine mortality within each 20
 382 x 20-m cell (Figure 4). We found a positive main effect of proportion of host trees per cell, with a greater
 383 proportion of host trees (i.e., ponderosa pine) in a cell increasing the probability of ponderosa pine mortality.
 384 Conversely, we found a negative effect of overall tree density (i.e., including both ponderosa pine and non-host
 385 species) such that greater tree density in a 20 x 20-m cell (for the same proportion of host trees) would
 386 decrease the probability of ponderosa pine mortality. We found a positive two-way interaction between the
 387 overall tree density per cell and the proportion of trees that were hosts, which is equivalent to a positive
 388 effect of the density of host trees (Figure 4).

389 We found a negative effect of mean height of ponderosa pine on the probability of ponderosa mortality,
 390 suggesting that WPB attacked smaller trees, on average. However, there was a positive interaction between
 391 CWD and ponderosa pine mean height, such that larger trees were more likely to increase the probability of




Figure 4: Posterior distributions of effect size from zero-inflated binomial model predicting the probability of ponderosa pine mortality in a 20 x 20-m cell given forest structure characteristics and site-level climatic water deficit (CWD). The gray density distribution for each model covariate represents the density of the posterior distribution, the point underneath each density curve represents the median of the estimate, the bold interval surrounding the point estimate represents the 66% credible interval, and the thin interval surrounding the point estimate represents the 95% credible interval.

392 ponderosa mortality in hotter, drier sites (Figure 5).

393 We found weakly negative effects of the site-level CWD interactions with both the proportion of host trees
394 and overall tree density (Figure 4).




Figure 5: Line version of model results with 95% credible intervals showing primary influence of ponderosa pine structure on the probability of ponderosa pine mortality, and the interaction across climatic water deficit. The 'larger trees' line represents the mean height of ponderosa pine 0.7 standard deviations above the mean (approximately 24.1 m), and the 'smaller trees' line represents the mean height of ponderosa pine 0.7 standard deviations below the mean (approximately 12.1 m).

395 Discussion

396 This study represents a novel use of drones to further our understanding of the simultaneous effects of
397 local forest structure and composition with broad-scale environmental gradients on tree mortality attributed
398 to WPB. We found strong positive effects (effect sizes >0.4) of both the proportion of host trees and the
399 interaction between site CWD and host tree mean size (height) on the probability of ponderosa pine mortality.
400 Conversely, we found a strong negative effect (effect size <-0.4) of mean height of ponderosa pine. Site-level
401 CWD exerted a positive, but relatively weak, main effect on the probability of ponderosa mortality (effect
402 size: 0.16; 95% CI: [0.03, 0.29]). To that end, we did not measure tree water stress at an individual tree level
403 as in other recent work (Stephenson et al. 2019), and instead treated CWD as a general indicator of tree
404 stress following results of coarser-scale studies (e.g., Asner et al. 2016, Young et al. 2017), which may have

405 contributed to our failure to detect a stronger CWD effect. Also, our entire study area experienced the same
406 extreme hot drought between 2012 and 2015 and the variation of mortality explained by a main effect of
407 CWD may be dampened when most trees are experiencing a high degree of water stress (Floyd et al. 2009,
408 Fettig et al. 2019).

409 **Positive effect of host density and a negative effect of overall density**

410 The strongest effect on the probability of ponderosa pine mortality was the positive effect of the proportion
411 of trees in each 20 x 20-m cell that were ponderosa pine– the host of the WPB (effect size: 0.76; 95% CI:
412 [0.70, 0.82]).

413 A number of mechanisms associated with the relative abundance of species in a local area might underlie
414 this relationship. Frequency-dependent herbivory–whereby mixed-species forests experience less herbivory
415 compared to monocultures (as an extreme example)– is common, especially for oligophagous insect species
416 (Jactel and Brockerhoff 2007). Furthermore, it has been demonstrated that nonhost volatiles reduce attraction
417 of several species of bark beetles to their aggregation pheromones (Seybold et al. 2018), including WPB
418 (Fettig et al. 2005). To that end, combinations of nonhost volatiles and an antiaggregation pheromone have
419 been used successfully to reduce levels of tree mortality attributed to WPB (e.g., Fettig et al. 2012). In
420 general, Hayes et al. (2009) and Fettig et al. (2019) found that measures of host availability explained less
421 variation in mortality than measures of overall tree density, but those conclusions were based on a response
422 variable of “total number of dead host trees,” rather than the number of dead host trees conditional on the
423 total number of host trees as in our study (i.e., a binomial response).

424 The negative relationship between overall tree density, a potential correlate of the local competitive envi-
425 ronment, and the probability of ponderosa pine mortality is counter-intuitive but corroborates findings of
426 coincident ground plots (Fettig et al. 2019, in their analysis using proportion of trees killed as a response)
427 and other work during the same hot drought (Restaino et al. 2019). In the absence of management, the forest
428 structure is itself a product of climate and, with increasing importance at finer spatial scales, topographic
429 conditions (Fricker et al. 2019). Thus, the denser forest patches in our study may indicate greater local
430 water availability, more favorable conditions for tree growth and survivorship, and increased resistance to
431 beetle-induced mortality (Restaino et al. 2019). The negative two-way interaction between site CWD and
432 overall density that amplifies the negative overall density effect in hotter, drier sites (effect size: -0.16; 95%
433 CI: [-0.24, -0.07]) supports this explanation if greater local tree density implies especially favorable growing
434 conditions (and locally resistant trees) when denser patches are found in hot, dry sites.

435 We found a positive two-way interaction between overall tree density (host and non-host) within each cell
436 and proportion of host trees, which is equivalent to a positive effect of host density (effect size: 0.08; 95% CI:
437 [0.03, 0.12]). The relationship between host density and susceptibility to colonization by bark beetles has
438 been so well-documented at the experimental plot level (e.g., Raffa and Berryman 1987, Oliver 1995) that
439 lowering stand densities through selective harvest of hosts is commonly recommended for reducing future
440 levels of tree mortality attributed to bark beetles (Fettig and Hilszczański 2015), including WPB (Fettig
441 2016). Greater host density shortens the flight distance required for WPB to disperse to new hosts, which
442 likely facilitates bark beetle spread, however we calibrated our aerial tree detection to ~400 m² areas rather
443 than to individual tree locations, so our data are insufficient to address these relationships. Increased density
444 of ponderosa pine, specifically, may disproportionately increase the competitive environment for host trees
445 (and thus increase their susceptibility to WPB colonization) if intraspecific competition amongst ponderosa
446 pine trees is stronger than interspecific competition as would be predicted with coexistence theory (Chesson
447 2000). Finally, greater host densities increase the frequency that searching WPB land on hosts, rather than
448 nonhosts, thus reducing the amount of energy expended during host finding and selection as well as the time
449 that searching WPB spend exposed to predators.

450 **Negative main effect of host tree mean size, but strong positive interaction with site CWD**

451 Counter to our expectations, we found an overall negative effect of host tree mean size on the probability of
452 host mortality (effect size: -0.40; 95% CI: [-0.46, -0.34]). WPB exhibit a preference for trees 50.8 to 76.2
453 cm in diameter at breast height (Person 1928, 1931), and a positive relationship between host tree size and
454 levels of tree mortality attributed to WPB was reported by Fettig et al. (2019) in the coincident field plots
455 as well as in other recent studies (Restaino et al. 2019, Stephenson et al. 2019, Pile et al. 2019). Indeed,
456 Fettig et al. (2019) reported no mortality in ponderosa pine trees <10.0 cm DBH attributable to WPB and
457 found no tree size/mortality relationship for incense cedar or white fir in the coincident field plots. These
458 species represent 22.3% of the total tree mortality observed in their study, yet in our study all dead trees
459 were classified as ponderosa pine (see Methods) which could dampen positive effect of tree size on mortality.
460 Larger trees are more nutritious and are therefore ideal targets if local bark beetle density is high enough
461 to successfully initiate mass attack as can occur when many trees are under severe water stress (Bentz et
462 al. 2010, Kolb et al. 2016). In the recent hot drought, we expected that most trees would be under severe
463 water stress, setting the stage for increasing beetle density, successful mass attacks, and targeting of larger
464 trees. A possible explanation for our finding counter to this expectation is that our observations represent the
465 cumulative mortality of trees during a multi-year drought event and its aftermath. Lower host tree mean size

466 led to a greater probability of host mortality earlier in this drought (Pile et al. 2019, Stovall et al. 2019) and
467 that signal might have persisted even as mortality continued to accumulate driven by other factors. Finally,
468 tree growth rates may be a better predictor of susceptibility to WPB colonization than tree size per se, with
469 slower-growing trees being most vulnerable (Miller and Keen 1960). While slow-growing trees are often also
470 the largest trees, this may not be the case for our study sites especially given the legacy of fire exclusion
471 in the Sierra Nevada and its effect of perturbing forest structure far outside its natural range of variation
472 (Safford and Stevens 2017).

473 We did observe a strong host tree size effect in its interaction with site CWD (effect size: 0.44; 95% CI: [0.34,
474 0.54]). In hot, dry sites, larger average host size increased the probability of host mortality while smaller host
475 sizes increased the probability of host mortality in cooler, wetter sites. Notably, a similar pattern was shown
476 by Stovall et al. (2019) with a strong positive tree height/mortality relationship in areas with the greatest
477 vapor pressure deficit and no tree height/mortality relationship in areas with the lowest vapor pressure
478 deficit. Stovall et al. (2019) did not observe that this environmental dependence extended to a negative tree
479 height/mortality relationship (as we did) even at the lowest extremes of their vapor pressure deficit gradient,
480 perhaps because their entire study took place in the southern Sierra Nevada which represents a hotter, drier
481 portion of the more spatially extensive results we present here. Our work suggests that the WPB was cueing
482 into different aspects of forest structure across an environmental gradient in a spatial context in a parallel
483 manner to the temporal context noted by Stovall et al. (2019) and Pile et al. (2019), who observed that
484 mortality was increasingly driven by larger trees as the hot drought proceeded and became more severe.

485 All of our sites were considered in an “epidemic” population phase for WPB (>5 trees killed per hectare; see
486 Supplemental Information; Miller and Keen 1960, Hayes et al. 2009), but our results challenge the notion that
487 outbreak behavior by the WPB and subsequent tree mortality is always driven by greater tree size. Despite a
488 strong tree size/mortality relationship in coincident ground plots across our study area (Fettig et al. 2019),
489 our results from surveying the broader context surrounding those ground plots reveals different effects of
490 host tree size depending on CWD. Thus, it is possible that the massive tree mortality in hotter/drier Sierra
491 Nevada forests (lower latitudes and elevations; Asner et al. 2016, Young et al. 2017) during the 2012 to
492 2015 hot drought arose as a synergistic alignment of environmental conditions and local forest structure that
493 allowed WPB to successfully colonize large trees, rapidly increase in population size, and expand. Conversely,
494 our results may suggest that the unexpectedly low mortality in cooler/wetter Sierra Nevada forests compared
495 to model predictions based on coarser-scale forest structure data (Young et al. 2017) could be explained
496 by a different WPB response to local forest structure due to a lack of an alignment with favorable climate
497 conditions.

498 **Limitations and future directions**

499 We have demonstrated that drones can be effective means of collecting forest data at multiple, vastly different
500 spatial scales to investigate a single, multi-scale phenomenon— from meters in between trees, to hundreds
501 of meters of elevation, to hundreds of thousands of meters of latitude. Some limitations remain but can be
502 overcome with further refinements in the use of this tool for forest ecology. Most of these limitations arise
503 from tree detection and classification uncertainty, and thus it was imperative to work with field data for
504 calibration and uncertainty reporting.

505 The greatest limitation in our study arising from classification uncertainty is in the assumption that all dead
506 trees were ponderosa pine, which we estimate from coincident field plots is true approximately 73.4% of
507 the time. Because the forest structure factors influencing the likelihood of individual tree mortality during
508 the hot drought depended on tree species (Stephenson et al. 2019), we cannot rule out that some of the
509 ponderosa pine mortality relationships to forest structure that we observed may be partially explained by
510 those relationships in other species that were misclassified as ponderosa pine using our methods. However,
511 the overall community composition across our study area was similar (Fettig et al. 2019) and we are able
512 to reproduce similar forest structure/mortality patterns in drone-derived data when restricting the scope
513 of analysis to only trees detected in the footprints of the coincident field plots with dramatically different
514 patterns observed when including data from the forest surrounding the coincident field plots (see Supplemental
515 information). Thus, we remain confident that the patterns we observed were driven primarily by the dynamic
516 between WPB and ponderosa pine. While spectral information of foliage could help classify living trees to
517 species, the species of standing dead trees were not spectrally distinct. This challenge of classifying standing
518 dead trees to species implies that a conifer forest system with less bark beetle and tree host diversity, such
519 as mountain pine beetle outbreaks in monocultures of lodgepole pine in the Intermountain West, should be
520 particularly amenable to the methods presented here even with minimal further refinement because dead
521 trees will almost certainly belong to a single species and have succumbed to colonization by a single bark
522 beetle species.

523 Some uncertainty surrounded our ability to detect trees using the geometry of the dense point clouds derived
524 with SfM. The horizontal accuracy of the tree detection was better than the vertical accuracy, which may
525 result from a more significant error contribution by the field-based calculations of tree height compared to
526 tree position relative to plot center (Table 2). Both the horizontal and vertical accuracy would likely improve
527 with better SfM point clouds, which can be enhanced with greater overlap between images (Frey et al. 2018)
528 or with oblique (i.e., off-nadir) imagery (James and Robson 2014). Frey et al. (2018) found that 95% overlap

529 was preferable for generating dense point clouds in forested areas, and James and Robson (2014) reduced
530 dense point cloud errors using imagery taken at 30 degrees off-nadir. We only achieved 91.6% overlap with
531 the X3 RGB camera and 83.9% overlap with the multispectral camera, and all imagery was nadir-facing.
532 While our live/dead classification was fairly accurate (97.3% on a withheld dataset), our species classifier
533 would likely benefit from better crown segmentation because the pixel-level reflectance values within each
534 crown are averaged to characterize the “spectral signature” of each tree. With better delineation of each
535 tree crown, the mean value of pixels within each tree crown will likely be more representative of that tree’s
536 spectral signature. Better crown segmentation might most readily be achieved through greater overlap in
537 imagery. Finally, we anticipate that computer vision and deep learning will prove helpful in overcoming some
538 of these detection and classification challenges (Gray et al. 2019).

539 **Conclusions**

540 Climate change adaptation strategies emphasize management action that considers whole-ecosystem responses
541 to inevitable change (Millar et al. 2007), which requires a macroecological understanding of how phenomena
542 at multiple scales can interact. We’ve shown that drones can be a valuable tool for investigating multi-scalar
543 phenomena, such as how local forest structure combines with environmental conditions to shape forest insect
544 disturbance. Understanding the conditions that drive dry western U.S. forest responses to disturbances such
545 as bark beetle outbreaks will be vital for predicting outcomes from increasing disturbance frequency and
546 intensity exacerbated by climate change. Our study suggests that outcomes will depend on interactions
547 between local forest structure and broad-scale environmental gradients, with the potential for cross-scale
548 interactions to challenge our current understanding of forest insect dynamics.

549 **References**

- 550 Anderegg, W. R. L., J. A. Hicke, R. A. Fisher, C. D. Allen, J. Aukema, B. Bentz, S. Hood, J. W. Lichstein,
551 A. K. Macalady, N. McDowell, Y. Pan, K. Raffa, A. Sala, J. D. Shaw, N. L. Stephenson, C. Tague, and
552 M. Zeppel. 2015. Tree mortality from drought, insects, and their interactions in a changing climate. *New*
553 *Phytologist* 208:674–683.
- 554 Asner, G. P., P. G. Brodrick, C. B. Anderson, N. Vaughn, D. E. Knapp, and R. E. Martin. 2016. Progressive
555 forest canopy water loss during the 2012–2015 California drought. *Proceedings of the National Academy of*
556 *Sciences* 113:E249–E255.

- 557 Baldwin, B. G., A. H. Thornhill, W. A. Freyman, D. D. Ackerly, M. M. Kling, N. Morueta-Holme, and B. D.
558 Mishler. 2017. Species richness and endemism in the native flora of California. American Journal of Botany
559 104:487–501.
- 560 Bedard, W. D., P. E. Tilden, D. L. Wood, R. M. Silverstein, R. G. Brownlee, and J. O. Rodin. 1969.
561 Western pine beetle: Field response to its sex pheromone and a synergistic host terpene, myrcene. Science
562 164:1284–1285.
- 563 Bentz, B. J., J. Régnière, C. J. Fettig, E. M. Hansen, J. L. Hayes, J. A. Hicke, R. G. Kelsey, J. F. Negrón,
564 and S. J. Seybold. 2010. Climate change and bark beetles of the western United States and Canada: Direct
565 and indirect effects. BioScience 60:602–613.
- 566 Berryman, A. A. 1982. Population dynamics of bark beetles. Pages 264–314 in Bark Beetles in North
567 American Conifers: A System for the Study of Evolutionary Biology.
- 568 Brooks, S. P., and A. Gelman. 1998. General methods for monitoring convergence of iterative simulations.
569 Journal of Computational and Graphical Statistics 7:434.
- 570 Bürkner, P.-C. 2017. **brms**: An *R* package for bayesian multilevel models using *Stan*. Journal of Statistical
571 Software 80:1–28.
- 572 Byers, J. A., and D. L. Wood. 1980. Interspecific inhibition of the response of the bark beetles, *Dendroctonus*
573 *brevicomis* and *Ips paraconfusus*, to their pheromones in the field. Journal of Chemical Ecology 6:149–164.
- 574 Chesson, P. 2000. Mechanisms of maintenance of species diversity. Annual Review of Ecology and Systematics
575 31:343–366.
- 576 Chubaty, A. M., B. D. Roitberg, and C. Li. 2009. A dynamic host selection model for mountain pine beetle,
577 *Dendroctonus ponderosae* Hopkins. Ecological Modelling 220:1241–1250.
- 578 Clevers, J., and A. Gitelson. 2013. Remote estimation of crop and grass chlorophyll and nitrogen content using
579 red-edge bands on Sentinel-2 and -3. International Journal of Applied Earth Observation and Geoinformation
580 23:344–351.
- 581 Coops, N. C., M. Johnson, M. A. Wulder, and J. C. White. 2006. Assessment of QuickBird high spatial
582 resolution imagery to detect red attack damage due to mountain pine beetle infestation. Remote Sensing of
583 Environment 103:67–80.
- 584 DJI. 2015a. Zenmuse X3 - Creativity Unleashed. <https://www.dji.com/zenmuse-x3/info>.
- 585 DJI. 2015b. DJI - The World Leader in Camera Drones/Quadcopters for Aerial Photography. [https:](https://)

- 586 //www.dji.com/matrice100/info.
- 587 DronesMadeEasy. 2018. Map Pilot for DJI on iOS. [https://itunes.apple.com/us/app/map-pilot-for-dji/
id1014765000?mt=8](https://itunes.apple.com/us/app/map-pilot-for-dji/id1014765000?mt=8).
- 589 Evenden, M. L., C. M. Whitehouse, and J. Sykes. 2014. Factors influencing flight capacity of the mountain
590 pine beetle (Coleoptera: Curculionidae: Scolytinae). *Environmental Entomology* 43:187–196.
- 591 Eysn, L., M. Hollaus, E. Lindberg, F. Berger, J.-M. Monnet, M. Dalponte, M. Kobal, M. Pellegrini, E.
592 Lingua, D. Mongus, and N. Pfeifer. 2015. A benchmark of LiDAR-based single tree detection methods using
593 heterogeneous forest data from the alpine space. *Forests* 6:1721–1747.
- 594 Farr, T. G., P. A. Rosen, E. Caro, R. Crippen, R. Duren, S. Hensley, M. Kobrick, M. Paller, E. Rodriguez, L.
595 Roth, D. Seal, S. Shaffer, J. Shimada, J. Umland, M. Werner, M. Oskin, D. Burbank, and D. Alsdorf. 2007.
596 The shuttle radar topography mission. *Reviews of Geophysics* 45.
- 597 Fettig, C. J. 2012. Chapter 2: Forest health and bark beetles. *in* Managing Sierra Nevada Forests. PSW-
598 GTR-237. USDA Forest Service.
- 599 Fettig, C. J. 2016. Native bark beetles and wood borers in Mediterranean forests of California. Pages 499–528
600 *in* Insects and diseases of Mediterranean Forest systems. Springer International Publishing, Switzerland.
- 601 Fettig, C. J., and J. Hilszczański. 2015. Management strategies for bark beetles in conifer forests. Pages
602 555–584 *in* Bark Beetles. Elsevier.
- 603 Fettig, C. J., K. D. Klepzig, R. F. Billings, A. S. Munson, T. E. Nebeker, J. F. Negrón, and J. T. Nowak. 2007.
604 The effectiveness of vegetation management practices for prevention and control of bark beetle infestations in
605 coniferous forests of the western and southern United States. *Forest Ecology and Management* 238:24–53.
- 606 Fettig, C. J., S. R. McKelvey, C. P. Dabney, D. P. W. Huber, C. G. Lait, D. L. Fowler, and J. H. Borden. 2012.
607 Efficacy of “Verbenone Plus” for protecting ponderosa pine trees and stands from *Dendroctonus brevicomis*
608 (Coleoptera: Curculionidae) attack in British Columbia and California. *Journal of Economic Entomology*
609 105:1668–1680.
- 610 Fettig, C. J., S. R. McKelvey, and D. P. W. Huber. 2005. Nonhost angiosperm volatiles and Verbenone disrupt
611 response of western pine beetle, *Dendroctonus brevicomis* (Coleoptera: Scolytidae), to attractant-baited traps.
612 *Journal of Economic Entomology* 98:2041–2048.
- 613 Fettig, C. J., L. A. Mortenson, B. M. Bulaon, and P. B. Foulk. 2019. Tree mortality following drought in the
614 central and southern Sierra Nevada, California, U.S. *Forest Ecology and Management* 432:164–178.

- 615 Flint, L. E., A. L. Flint, J. H. Thorne, and R. Boynton. 2013. Fine-scale hydrologic modeling for regional land-
616 scape applications: The California Basin Characterization Model development and performance. Ecological
617 Processes 2:25.
- 618 Floyd, M. L., M. Clifford, N. S. Cobb, D. Hanna, R. Delph, P. Ford, and D. Turner. 2009. Relationship of
619 stand characteristics to drought-induced mortality in three Southwestern piñonJuniper woodlands. Ecological
620 Applications 19:1223–1230.
- 621 Franceschi, V. R., P. Krokene, E. Christiansen, and T. Krekling. 2005. Anatomical and chemical defenses of
622 conifer bark against bark beetles and other pests. New Phytologist 167:353–376.
- 623 Frey, J., K. Kovach, S. Stemmler, and B. Koch. 2018. UAV photogrammetry of forests as a vulnerable
624 process. A sensitivity analysis for a structure from motion RGB-image pipeline. Remote Sensing 10:912.
- 625 Fricker, G. A., N. W. Synes, J. M. Serra-Diaz, M. P. North, F. W. Davis, and J. Franklin. 2019. More than
626 climate? Predictors of tree canopy height vary with scale in complex terrain, Sierra Nevada, CA (USA).
627 Forest Ecology and Management 434:142–153.
- 628 Gabry, J., D. Simpson, A. Vehtari, M. Betancourt, and A. Gelman. 2019. Visualization in Bayesian workflow.
629 Journal of the Royal Statistical Society: Series A (Statistics in Society) 182:389–402.
- 630 Gitelson, A., and M. N. Merzlyak. 1994. Spectral reflectance changes associated with autumn senescence of
631 *Aesculus hippocastanum* L. And *Acer platanoides* L. Leaves. Spectral features and relation to chlorophyll
632 estimation. Journal of Plant Physiology 143:286–292.
- 633 Graf, M., M. Reid, B. Aukema, and B. Lindgren. 2012. Association of tree diameter with body size and lipid
634 content of mountain pine beetles. The Canadian Entomologist 144:467–477.
- 635 Gray, P. C., A. B. Fleishman, D. J. Klein, M. W. McKown, V. S. Bézy, K. J. Lohmann, and D. W. Johnston.
636 2019. A convolutional neural network for detecting sea turtles in drone imagery. Methods in Ecology and
637 Evolution 10:345–355.
- 638 Griffin, D., and K. J. Anchukaitis. 2014. How unusual is the 2012-2014 California drought? Geophysical
639 Research Letters 41:9017–9023.
- 640 Hayes, C. J., C. J. Fettig, and L. D. Merrill. 2009. Evaluation of multiple funnel traps and stand characteristics
641 for estimating western pine beetle-caused tree mortality. Journal of Economic Entomology 102:2170–2182.
- 642 Hijmans, R. J., J. van Etten, M. Sumner, J. Cheng, A. Bevan, R. Bivand, L. Busetto, M. Canty, D. Forrest,
643 A. Ghosh, D. Golicher, J. Gray, J. A. Greenberg, P. Hiemstra, I. for M. A. Geosciences, C. Karney, M.

- 644 Mattiuzzi, S. Mosher, J. Nowosad, E. Pebesma, O. P. Lamigueiro, E. B. Racine, B. Rowlingson, A. Shortridge,
645 B. Venables, and R. Wueest. 2019. *Raster*: Geographic data analysis and modeling.
- 646 Hunziker, P. 2017. *Velox*: Fast raster manipulation and extraction.
- 647 Jactel, H., and E. G. Brockerhoff. 2007. Tree diversity reduces herbivory by forest insects. *Ecology Letters*
648 10:835–848.
- 649 Jakubowski, M. K., W. Li, Q. Guo, and M. Kelly. 2013. Delineating individual trees from LiDAR data: A
650 comparison of vector- and raster-based segmentation approaches. *Remote Sensing* 5:4163–4186.
- 651 James, M. R., and S. Robson. 2014. Mitigating systematic error in topographic models derived from UAV
652 and ground-based image networks. *Earth Surface Processes and Landforms* 39:1413–1420.
- 653 Kane, V. R., M. P. North, J. A. Lutz, D. J. Churchill, S. L. Roberts, D. F. Smith, R. J. McGaughey, J. T.
654 Kane, and M. L. Brooks. 2014. Assessing fire effects on forest spatial structure using a fusion of Landsat and
655 airborne LiDAR data in Yosemite National Park. *Remote Sensing of Environment* 151:89–101.
- 656 Kolb, T. E., C. J. Fettig, M. P. Ayres, B. J. Bentz, J. A. Hicke, R. Mathiasen, J. E. Stewart, and A. S. Weed.
657 2016. Observed and anticipated impacts of drought on forest insects and diseases in the United States. *Forest
658 Ecology and Management* 380:321–334.
- 659 Kuhn, M. 2008. Building predictive models in R using the caret package. *Journal of Statistical Software*
660 28:1–26.
- 661 Larson, A. J., and D. Churchill. 2012. Tree spatial patterns in fire-frequent forests of western North America,
662 including mechanisms of pattern formation and implications for designing fuel reduction and restoration
663 treatments. *Forest Ecology and Management* 267:74–92.
- 664 Li, W., Q. Guo, M. K. Jakubowski, and M. Kelly. 2012. A new method for segmenting individual trees from
665 the LiDAR point cloud. *Photogrammetric Engineering & Remote Sensing* 78:75–84.
- 666 Logan, J. A., P. White, B. J. Bentz, and J. A. Powell. 1998. Model analysis of spatial patterns in mountain
667 pine beetle outbreaks. *Theoretical Population Biology* 53:236–255.
- 668 Meyer, F., and S. Beucher. 1990. Morphological segmentation. *Journal of Visual Communication and Image
669 Representation* 1:21–46.
- 670 Micasense. 2015. MicaSense. <https://support.micasense.com/hc/en-us/articles/215261448-RedEdge-User->
671 Manual-PDF-Download-.

- 672 Millar, C. I., N. L. Stephenson, and S. L. Stephens. 2007. Climate change and forests of the future: Managing
673 in the face of uncertainty. *Ecological Applications* 17:2145–2151.
- 674 Millar, C. I., R. D. Westfall, D. L. Delany, M. J. Bokach, A. L. Flint, and L. E. Flint. 2012. Forest mortality in
675 high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern California, USA: Influence of environmental
676 context, bark beetles, climatic water deficit, and warming. *Canadian Journal of Forest Research* 42:749–765.
- 677 Miller, J. M., and F. P. Keen. 1960. Biology and control of the western pine beetle: A summary of the first
678 fifty years of research. US Department of Agriculture.
- 679 Moeck, H. A., D. L. Wood, and K. Q. Lindahl. 1981. Host selection behavior of bark beetles (Coleoptera:
680 Scolytidae) attacking *Pinus ponderosa*, with special emphasis on the western pine beetle, *Dendroctonus*
681 *brevicomis*. *Journal of Chemical Ecology* 7:49–83.
- 682 Morris, J. L., S. Cottrell, C. J. Fettig, W. D. Hansen, R. L. Sherriff, V. A. Carter, J. L. Clear, J. Clement, R.
683 J. DeRose, J. A. Hicke, P. E. Higuera, K. M. Mattor, A. W. R. Seddon, H. T. Seppä, J. D. Stednick, and S.
684 J. Seybold. 2017. Managing bark beetle impacts on ecosystems and society: Priority questions to motivate
685 future research. *Journal of Applied Ecology* 54:750–760.
- 686 Oliver, W. W. 1995. Is self-thinning in ponderosa pine ruled by *Dendroctonus* bark beetles? Page 6 in Forest
687 health through silviculture: Proceedings of the 1995 National Silviculture Workshop.
- 688 Pau, G., F. Fuchs, O. Sklyar, M. Boutros, and W. Huber. 2010. EBImage: An R package for image processing
689 with applications to cellular phenotypes. *Bioinformatics* 26:979–981.
- 690 Person, H. L. 1928. Tree selection by the western pine beetle. *Journal of Forestry* 26:564–578.
- 691 Person, H. L. 1931. Theory in explanation of the selection of certain trees by the western pine beetle. *Journal*
692 *of Forestry* 29:696–699.
- 693 Pile, L. S., M. D. Meyer, R. Rojas, O. Roe, and M. T. Smith. 2019. Drought impacts and compounding
694 mortality on forest trees in the southern Sierra Nevada. *Forests* 10:237.
- 695 Plowright, A. 2018. ForestTools: Analyzing remotely sensed forest data.
- 696 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical
697 Computing, Vienna, Austria.
- 698 Raffa, K. F., B. H. Aukema, B. J. Bentz, A. L. Carroll, J. A. Hicke, M. G. Turner, and W. H. Romme. 2008.
699 Cross-scale drivers of natural disturbances prone to anthropogenic amplification: The dynamics of bark beetle
700 eruptions. *BioScience* 58:501–517.

- 701 Raffa, K. F., and A. A. Berryman. 1983. The role of host plant resistance in the colonization behavior and
702 ecology of bark beetles (Coleoptera: Scolytidae). Ecological Monographs 53:27–49.
- 703 Raffa, K. F., and A. A. Berryman. 1987. Interacting selective pressures in conifer-bark beetle systems: A
704 basis for reciprocal adaptations? The American Naturalist 129:234–262.
- 705 Raffa, K. F., J.-C. Grégoire, and B. Staffan Lindgren. 2015. Natural history and ecology of bark beetles.
706 Pages 1–40 in Bark Beetles. Elsevier.
- 707 Restaino, C., D. Young, B. Estes, S. Gross, A. Wuenschel, M. Meyer, and H. Safford. 2019. Forest
708 structure and climate mediate drought-induced tree mortality in forests of the Sierra Nevada, USA. Ecological
709 Applications 0:e01902.
- 710 Rouse, W., R. H. Haas, W. Deering, and J. A. Schell. 1973. Monitoring the vernal advancement and
711 retrogradation (green wave effect) of natural vegetation. Type II Report, Goddard Space Flight Center,
712 Greenbelt, MD, USA.
- 713 Roussel, J.-R. 2019. lidRplugins: Extra functions and algorithms for lidR package.
- 714 Roussel, J.-R., D. Auty, F. De Boissieu, and A. S. Meador. 2019. lidR: Airborne LiDAR data manipulation
715 and visualization for forestry applications.
- 716 Safford, H. D., and J. T. Stevens. 2017. Natural range of variation for yellow pine and mixed-conifer forests
717 in the Sierra Nevada, Southern Cascades, and Modoc and Inyo National Forests, California, USA. Page 241.
- 718 dos Santos, A. A., J. Marcato Junior, M. S. Araújo, D. R. Di Martini, E. C. Tetila, H. L. Siqueira, C. Aoki, A.
719 Eltner, E. T. Matsubara, H. Pistori, R. Q. Feitosa, V. Liesenberg, and W. N. Gonçalves. 2019. Assessment of
720 CNN-Based Methods for Individual Tree Detection on Images Captured by RGB Cameras Attached to UAVs.
721 Sensors (Basel, Switzerland) 19.
- 722 Seidl, R., J. Müller, T. Hothorn, C. Bässler, M. Heurich, and M. Kautz. 2016. Small beetle, large-scale
723 drivers: How regional and landscape factors affect outbreaks of the European spruce bark beetle. The Journal
724 of applied ecology 53:530–540.
- 725 Seybold, S. J., B. J. Bentz, C. J. Fettig, J. E. Lundquist, R. A. Progar, and N. E. Gillette. 2018. Management
726 of western North American bark beetles with semiochemicals. Annual Review of Entomology 63:407–432.
- 727 Shepherd, W. P., D. P. W. Huber, S. J. Seybold, and C. J. Fettig. 2007. Antennal responses of the western
728 pine beetle, *Dendroctonus brevicomis* (Coleoptera: Curculionidae), to stem volatiles of its primary host,
729 *Pinus ponderosa*, and nine sympatric nonhost angiosperms and conifers. Chemoecology 17:209–221.

- 730 Shiklomanov, A. N., B. A. Bradley, K. M. Dahlin, A. M. Fox, C. M. Gough, F. M. Hoffman, E. M. Middleton,
731 S. P. Serbin, L. Smallman, and W. K. Smith. 2019. Enhancing global change experiments through integration
732 of remote-sensing techniques. *Frontiers in Ecology and the Environment* 0.
- 733 Shin, P., T. Sankey, M. Moore, and A. Thode. 2018. Evaluating unmanned aerial vehicle images for estimating
734 forest canopy fuels in a ponderosa pine stand. *Remote Sensing* 10:1266.
- 735 Stephenson, N. 1998. Actual evapotranspiration and deficit: Biologically meaningful correlates of vegetation
736 distribution across spatial scales. *Journal of Biogeography* 25:855–870.
- 737 Stephenson, N. L., A. J. Das, N. J. Ampersee, and B. M. Bulaon. 2019. Which trees die during drought?
738 The key role of insect host-tree selection. *Journal of Ecology*:75.
- 739 Stovall, A. E. L., H. Shugart, and X. Yang. 2019. Tree height explains mortality risk during an intense
740 drought. *Nature Communications* 10:1–6.
- 741 Thistle, H. W., H. Peterson, G. Allwine, B. Lamb, T. Strand, E. H. Holsten, and P. J. Shea. 2004. Surrogate
742 pheromone plumes in three forest trunk spaces: Composite statistics and case studies. *Forest Science* 50.
- 743 USDAFS. 2019, February 11. Press Release: Survey finds 18 million trees died in California in 2018.
744 https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/FSEPRD609321.pdf.
- 745 Vega, C., A. Hamrouni, S. El Mokhtari, J. Morel, J. Bock, J. P. Renaud, M. Bouvier, and S. Durrieu. 2014.
746 PTrees: A point-based approach to forest tree extraction from LiDAR data. *International Journal of Applied
747 Earth Observation and Geoinformation* 33:98–108.
- 748 Weinstein, B. G., S. Marconi, S. Bohlman, A. Zare, and E. White. 2019. Individual Tree-Crown Detection in
749 RGB Imagery Using Semi-Supervised Deep Learning Neural Networks. *Remote Sensing* 11:1309.
- 750 Wyngaard, J., L. Barbieri, A. Thomer, J. Adams, D. Sullivan, C. Crosby, C. Parr, J. Klump, S. Raj Shrestha,
751 and T. Bell. 2019. Emergent challenges for science sUAS data management: Fairness through community
752 engagement and best practices development. *Remote Sensing* 11:1797.
- 753 Young, D. J. N., J. T. Stevens, J. M. Earles, J. Moore, A. Ellis, A. L. Jirka, and A. M. Latimer. 2017.
754 Long-term climate and competition explain forest mortality patterns under extreme drought. *Ecology Letters*
755 20:78–86.
- 756 Zhang, W., J. Qi, P. Wan, H. Wang, D. Xie, X. Wang, and G. Yan. 2016. An easy-to-use airborne LiDAR
757 data filtering method based on cloth simulation. *Remote Sensing* 8:501.