

Mis-Classified, Binary, Endogenous Regressors: Identification and Inference

Francis J. DiTraglia¹ Camilo García-Jimeno^{2,3}

¹University of Pennsylvania

²Emory University

³NBER

October 11th, 2018

- Thank you for inviting me. Joint work with Camilo Garcia-Jimeno.
- Intro. 'metrics students learn that a valid IV serves double duty: correct for endogeneity and classical measurement error
- Classical measurement error is a special case: requires true value of regressor indep. of or at least uncorrelated with measurement error
- Applied work often involves endogenous binary regressor: smoker/non-smoker or union/non-union. Binary \implies non-classical error. True 0 \implies can only mis-measure *upwards* as 1; true 1 \implies can only mis-measure *downwards* as 0. Error *negatively correlated* with truth.
- To accommodate this, consider *non-diff* error. Say more later, but roughly non-diff means *conditionally classical*: condition on truth and controls, remaining component of error unrelated to everything else.
- Today pose simple question: binary, endog. regressor subject to non-diff. error. Can valid IV correct for *both* measurement error and endog?

What is the effect of T^* ?

$$y = c(\mathbf{x}) + \beta(\mathbf{x})T^* + \varepsilon$$

- ▶ y – Outcome of interest
- ▶ T^* – Unobserved, endogenous binary regressor
- ▶ T – Observed, mis-measured binary surrogate for T^*
- ▶ \mathbf{x} – Exogenous covariates
- ▶ z – Discrete (typically binary) instrumental variable

$$y = c(x) + \beta(x)T^* + \varepsilon$$

- y = Outcome of interest
- T^* = Unobserved, endogenous binary regressor
- T = Observed, mis-measured binary surrogate for T^*
- x = Exogenous covariates
- z = Discrete (typically binary) instrumental variable

└ What is the effect of T^* ?

- Here is the specific model I will focus on today. Additively separable model, want to learn the causal effect of binary regressor T^* on y . Unfortunately T^* is unobserved. Observe only mis-measured binary surrogate T . To make matters worse, T^* is endogenous, but we have a discrete instrument z .
- Additive separability is an assumption. Allow very general forms of observed heterogeneity through x but restricts unobserved heterogeneity.
- Conditionally linear model. This is without loss of generality since the model is additively separable and T^* is binary.
- Mainly focus on additively separable case today, but will also discuss implications of our results for a LATE model.

Using a discrete IV to learn about $\beta(\mathbf{x})$

$$y = c(\mathbf{x}) + \beta(\mathbf{x})T^* + \varepsilon$$

Contributions of This Paper

1. Show that only existing point identification result for mis-classified, endogenous T^* is incorrect.
2. Sharp identified set for β under standard assumptions.
3. Point identification of β under slightly stronger assumptions.
4. Point out problem of weak identification in mis-classification models, develop identification-robust inference for β .

$$y = c(\mathbf{x}) + \beta(\mathbf{x})T^* + \varepsilon$$

Contributions of This Paper

1. Show that only existing point identification result for mis-classified, endogenous T^* is incorrect.
2. Sharp identified set for β under standard assumptions.
3. Point identification of β under slightly stronger assumptions.
4. Point out problem of weak identification in mis-classification models, develop identification-robust inference for β .

└ Using a discrete IV to learn about $\beta(\mathbf{x})$

- Here are the main contributions of paper that I will discuss today.
- Many papers consider using IV to identify effect of exog. mis-measured binary regressor, but little work on endog. case. First: show only point identification result for this case incorrect: ident. is an open question.
- Next: use standard assumptions to derive the “sharp identified set” for β . This means *fully* exploit all information in the data and our assumptions to derive tightest possible bounds for β . If bounds contain a single point, β is point identified. Otherwise partially identified.
- Novel and informative bounds for β , but not point identified. Then consider slightly stronger assumptions that allow us to exploit additional features of the data and show that these suffice to point identify β .
- Next consider inference. Show that mis-classification models, suffer from potential weak identification. Propose procedure for robust inference.
- Now a motivating example. . .

Example: Smoking and Birthweight (SNAP Trial)

Coleman et al. (N Engl J Med, 2012)

RCT with pregnant smokers in England: half given nicotine patches, the rest given placebo patches. Some given nicotine fail to quit; some given placebo quit.

- ▶ y – Birthweight
- ▶ T^* – True smoking behavior
- ▶ T – Self-reported smoking behavior
- ▶ x – Mother characteristics
- ▶ z – Indicator of nicotine patch

Baseline Assumptions I – Model & Instrument

Additively Separable Model

$$y = c(\mathbf{x}) + \beta(\mathbf{x})T^* + \varepsilon, \quad \mathbb{E}[\varepsilon] = 0$$

Valid & Relevant Instrument: $z \in \{0, 1\}$

- ▶ $\mathbb{P}(T^* = 1 | \mathbf{x}, z = 1) \neq \mathbb{P}(T^* = 1 | \mathbf{x}, z = 0)$
- ▶ $\mathbb{E}[\varepsilon | \mathbf{x}, z] = 0$
- ▶ $0 < \mathbb{P}(z = 1 | \mathbf{x}) < 1$

Baseline Assumptions II – Measurement Error

Notation

► $\alpha_0(\mathbf{x}, z) \equiv \mathbb{P}(T = 1 | T^* = 0, \mathbf{x}, z)$

► $\alpha_1(\mathbf{x}, z) \equiv \mathbb{P}(T = 0 | T^* = 1, \mathbf{x}, z)$

Mis-classification unaffected by z

$$\alpha_0(\mathbf{x}, z) = \alpha_0(\mathbf{x}), \quad \alpha_1(\mathbf{x}, z) = \alpha_1(\mathbf{x})$$

Extent of Mis-classification

$$\alpha_0(\mathbf{x}) + \alpha_1(\mathbf{x}) < 1 \quad (T \text{ is positively correlated with } T^*)$$

Non-differential Mis-classification

$$\mathbb{E}[\varepsilon | \mathbf{x}, z, T^*, T] = \mathbb{E}[\varepsilon | \mathbf{x}, z, T^*]$$

Existing Results

Correct: Exogenous T^*

- ▶ Mahajan (2006), Frazis & Loewenstein (2003)
- ▶ $\mathbb{E}[\varepsilon|\mathbf{x}, z, T^*] = 0$ + “Baseline” $\Rightarrow \beta(\mathbf{x})$ identified.

Incorrect: Endogenous T^*

- ▶ Mahajan (2006) A.2
- ▶ $\mathbb{E}[\varepsilon|\mathbf{x}, z, T^*, T] = \mathbb{E}[\varepsilon|\mathbf{x}, T^*] + \text{“Baseline”} \Rightarrow \beta(\mathbf{x})$ identified.

We show: Mahajan's assumptions imply that the instrument z is uncorrelated with T^* unless T^* is in fact *exogenous*.

Existing Results

Existing Results

Correct: Exogenous T^*

- Mahajan (2006), Frazis & Loewenstein (2003)
- $E[y|x, z, T^*] = 0 + \text{"Baseline"} \Rightarrow \beta(x) \text{ identified.}$

Incorrect: Endogenous T^*

- Mahajan (2006) A.2
- $E[y|x, z, T^*] = E[y|x, T^*] + \text{"Baseline"} \Rightarrow \beta(x) \text{ identified.}$

We show: Mahajan's assumptions imply that the instrument z is uncorrelated with T^* unless T^* is in fact exogenous.

- Point out that the FL estimator is a nonlinear GMM rather than IV and note that they require *joint* exogeneity of T^* and z .
- 1st contribution: show that only existing point identification result for mis-measured, binary, endog. regressor is false
- As mentioned a few minutes ago, main result from Mahajan (2006; Ecta) is for T^* , but paper also contains a result for the endogenous case [READ THE RESULT]
- Exotic-looking assumption is needed to leverage Mahajan's result for the exogenous case. Unfortunately we show that it leads to a contradiction. [READ THE RESULT]
- Identification in this model is an open question: though Mahajan's proof fails, this does not establish that β is unidentified under the baseline assumptions.
- Next step show you two known results: simple bounds for α_0, α_1 , and relationship between IV estimator and α_0, α_1 , yielding bounds for β
- Then our 2nd contribution: sharp identified set for β under baseline

“Weak” Bounds

First-Stage

$$p_k(\mathbf{x}) \equiv \mathbb{P}(T = 1 | \mathbf{x}, z = k)$$

IV Estimand

$$\frac{\mathbb{E}[y | \mathbf{x}, z = 1] - \mathbb{E}[y | \mathbf{x}, z = 0]}{p_1(\mathbf{x}) - p_0(\mathbf{x})} = \frac{\beta(\mathbf{x})}{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})}$$

Bounds for (α_0, α_1)

$$\alpha_0(\mathbf{x}) \leq \min_k \{p_k(\mathbf{x})\}, \quad \alpha_1(\mathbf{x}) \leq \min_k \{1 - p_k(\mathbf{x})\} \quad \text{▶ more}$$

Bounds for β

$\beta(\mathbf{x})$ is between IV and Reduced form; same sign as IV. ▶ more

Binary Regressors

└ “Weak” Bounds

“Weak” Bounds

First-Stage

$$\rho_k(\mathbf{x}) = \mathbb{P}(T = 1 | \mathbf{x}, z = k)$$

IV Estimand

$$\frac{\mathbb{E}[y | \mathbf{x}, z = 1] - \mathbb{E}[y | \mathbf{x}, z = 0]}{\rho_1(\mathbf{x}) - \rho_0(\mathbf{x})} = \frac{\beta(\mathbf{x})}{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})}$$


Bounds for (α_0, α_1)

$$\alpha_0(\mathbf{x}) \leq \min_k \{\rho_k(\mathbf{x})\}, \quad \alpha_1(\mathbf{x}) \leq \min_k \{1 - \rho_k(\mathbf{x})\}$$

Bounds for β

$\beta(\mathbf{x})$ is between IV and Reduced form; same sign as IV

This doesn't rely on non-diff assumption or additive separability. Mention F&L (2003) and Ura (2016). But the point identification results from the literature rely on non-diff, and these bounds do not in fact impose that. Do these contain any additional information about β ? Perhaps they even point identify it!



Restrictions from Non-differential Mis-classification?

(Suppress \mathbf{x} for simplicity)

Notation

- ▶ $r_{tk} \equiv \mathbb{P}(T^* = 1 | T = t, z = k)$
- ▶ z_k is shorthand for $z = k$

Iterated Expectations over T^*

$$\mathbb{E}(y | T = 0, z_k) = (1 - r_{0k})\mathbb{E}(y | T^* = 0, T = 0, z_k) + r_{0k}\mathbb{E}(y | T^* = 1, T = 0, z_k)$$

$$\mathbb{E}(y | T = 1, z_k) = (1 - r_{1k})\mathbb{E}(y | T^* = 0, T = 1, z_k) + r_{1k}\mathbb{E}(y | T^* = 1, T = 1, z_k)$$

Restrictions from Non-differential Mis-classification?

(Suppress \mathbf{x} for simplicity)

Notation

- ▶ $r_{tk} \equiv \mathbb{P}(T^* = 1 | T = t, z = k)$
- ▶ z_k is shorthand for $z = k$

Adding Non-differential Assumption

$$\mathbb{E}(y | T = 0, z_k) = (1 - r_{0k})\mathbb{E}(y | T^* = 0, z_k) + r_{0k}\mathbb{E}(y | T^* = 1, z_k)$$

$$\mathbb{E}(y | T = 1, z_k) = (1 - r_{1k})\mathbb{E}(y | T^* = 0, z_k) + r_{1k}\mathbb{E}(y | T^* = 1, z_k)$$

2 equations in 2 unknowns \Rightarrow solve for $\mathbb{E}(y T^* = t^*, z = k)$ given (r_{0k}, r_{1k}) .

Restrictions from Non-differential Mis-classification?

Mixture Representation

$$F_{tk} = (1 - r_{tk})F_{tk}^0 + r_{tk}F_{tk}^1$$

$$F_{tk} \equiv y | (T = t, z = k)$$

$$F_{tk}^{t^*} \equiv y | (T^* = t^*, T = t, z = k)$$

Restrictions

- ▶ $\mathbb{E}(y | T^*, T, z) = \mathbb{E}(y | T^*, z)$ observable given (α_0, α_1)
- ▶ r_{tk} observable given (α_0, α_1)

Question

Given (α_0, α_1) can we always find (F_{tk}^0, F_{tk}^1) to satisfy the mixture model?

Restrictions from Non-differential Mis-classification?

Equivalent Problem




Given a specified CDF F , for what values of p and μ do there exist valid CDFs (G, H) with $F = (1 - p)G + pH$ and $\mu = \text{mean}(H)$?


Necessary and Sufficient Condition if F is Continuous

$$\underline{\mu}(F, p) \leq \mu \leq \bar{\mu}(F, p)$$




$$\underline{\mu}(F, p) \equiv \int_{-\infty}^{\infty} x \left[p^{-1} f(x) \mathbf{1}\{x < F^{-1}(p)\} \right] dx = \int_{-\infty}^{\infty} x \underline{h}(x) dx$$




$$\bar{\mu}(F, p) \equiv \int_{-\infty}^{\infty} x \left[p^{-1} f(x) \mathbf{1}\{x > F^{-1}(1 - p)\} \right] dx = \int_{-\infty}^{\infty} x \bar{h}(x) dx$$














Picture very simple: for given weight p on H , top panel shows the smallest mean that H can have and the bottom shows the largest mean it can take to yield a valid mixture in which H has weight p . As you change p , you change the range of values that the mean of H can take. In this example the observed distribution F is a simple mixture of normals. If it were a different distribution we'd get different restrictions: picture shows how shape of F leads to the upper and lower bounds for μ .
































As p approaches 1, the mean of H is more tightly constrained: must be close to the mean of the observed distribution F , namely -0.8 . As p approaches zero, it is less and less constrained: since it contributes very little to the overall mixture, it can take on nearly any mean.

Sharp Identified Set under Baseline Assumptions

Theorem

- (i) As long as $\mathbb{E}[y|\mathbf{x}, T = 0, z = k] \neq \mathbb{E}[y|\mathbf{x}, T = 1, z = k]$ for some k , non-differential measurement error strictly improves the weak bounds for α_0, α_1 , and β .
- (ii) Under the baseline assumptions, β is not point identified, regardless of how many (discrete) values z takes on.

Corollary

Our bounds for α_0, α_1 , and β remain valid in a LATE model, although they may not be sharp, since they do not incorporate the testable implications of the LATE assumptions.

└ Sharp Identified Set under Baseline

Theorem

- (i) As long as $\mathbb{E}[y|x, T=0, z=k] \neq \mathbb{E}[y|x, T=1, z=k]$ for some k , non-differential measurement error strictly improves the weak bounds for α_0, α_1 , and β .
- (ii) Under the baseline assumptions, β is not point identified, regardless of how many (discrete) values z takes on.

Corollary

Our bounds for α_0, α_1 , and β remain valid in a LATE model, although they may not be sharp, since they do not incorporate the testable implications of the LATE assumptions.

Second contribution. Simple bounds I showed you earlier are not sharp: in other words, they're not the best bounds you can get under our assumptions. Even when we get the best bounds (the “sharp” bounds) they're not enough to point identify β . Describe the intuition for why not: no mis-classification means r_{tk} is either zero or 1 so it is trivial to form the required mixture in this case. Also point out that the restrictions from non-differential measurement error can be very informative in practice! Now transition to point identification argument. Can we obtain point identification under stronger but credible assumptions?

Point Identification: 1st Ingredient

Reparameterization

$$\theta_1(\mathbf{x}) = \beta(\mathbf{x}) / [1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})]$$

$$\theta_2(\mathbf{x}) = [\theta_1(\mathbf{x})]^2 [1 + \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})]$$

$$\theta_3(\mathbf{x}) = [\theta_1(\mathbf{x})]^3 \left[\{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})\}^2 + 6\alpha_0(\mathbf{x}) \{1 - \alpha_1(\mathbf{x})\} \right]$$

Lemma

Baseline Assumptions $\implies \text{Cov}(y, z|\mathbf{x}) = \theta_1(\mathbf{x})\text{Cov}(z, T|\mathbf{x})$.

Binary Regressors

└ Point Identification: 1st Ingredient

Note that $\beta = 0$ iff $\theta_1 = \theta_2 = \theta_3 = 0$.

Point Identification: 1st Ingredient

Reparameterization

$$\theta_1(\mathbf{x}) = \beta(\mathbf{x}) / [1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})]$$

$$\theta_2(\mathbf{x}) = [\theta_1(\mathbf{x})]^2 [1 + \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})]$$

$$\theta_3(\mathbf{x}) = [\theta_1(\mathbf{x})]^3 \left[(1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x}))^2 + 6\alpha_0(\mathbf{x}) (1 - \alpha_1(\mathbf{x})) \right]$$

Lemma

Baseline Assumptions $\implies \text{Cov}(y, z|\mathbf{x}) = \theta_1(\mathbf{x})\text{Cov}(z, T|\mathbf{x})$.

Point Identification: 2nd Ingredient

Assumption (II)

$$\mathbb{E}[\varepsilon^2|\mathbf{x}, z] = \mathbb{E}[\varepsilon^2|\mathbf{x}]$$

Lemma

(Baseline) + (II) \implies

$$\text{Cov}(y^2, z|\mathbf{x}) = 2\text{Cov}(yT, z|\mathbf{x})\theta_1(\mathbf{x}) - \text{Cov}(T, z|\mathbf{x})\theta_2(\mathbf{x})$$

Corollary

(Baseline) + (II) + $[\beta(\mathbf{x}) \neq 0] \implies [\alpha_1(\mathbf{x}) - \alpha_0(\mathbf{x})]$ is identified.

Binary Regressors

└ Point Identification: 2nd Ingredient

Point Identification: 2nd Ingredient

Assumption (II)

$$\mathbb{E}[z^2 | \mathbf{x}, z] = \mathbb{E}[z^2 | \mathbf{x}]$$

Lemma

$$(\text{Baseline}) + (\text{I}) \implies$$

$$\text{Cov}(y^2, z | \mathbf{x}) = 2 \text{Cov}(y, z | \mathbf{x}) \theta_1(\mathbf{x}) - \text{Cov}(T, z | \mathbf{x}) \theta_2(\mathbf{x})$$

Corollary

$$(\text{Baseline}) + (\text{I}) + [\beta(\mathbf{x}) \neq 0] \implies [\alpha_1(\mathbf{x}) - \alpha_0(\mathbf{x})] \text{ is identified.}$$

Notice that the corollary implies that β is point identified if mis-classification is one-sided, as it might well be in the smoking example.

Point Identification: 3rd Ingredient

Assumption (III)

$$(i) \mathbb{E}[\varepsilon^2 | \mathbf{x}, z, T^*, T] = \mathbb{E}[\varepsilon^2 | \mathbf{x}, z, T^*]$$

$$(ii) \mathbb{E}[\varepsilon^3 | \mathbf{x}, z] = \mathbb{E}[\varepsilon^3 | \mathbf{x}]$$

Lemma

(Baseline) + (II) + (III) \implies

$$\text{Cov}(y^3, z | \mathbf{x}) = 3\text{Cov}(y^2 T, z | \mathbf{x})\theta_1(\mathbf{x}) - 3\text{Cov}(yT, z | \mathbf{x})\theta_2(\mathbf{x}) + \text{Cov}(T, z | \mathbf{x})\theta_3(\mathbf{x})$$

Point Identification Result

Theorem

(Baseline) + (II) + (III) $\implies \beta(\mathbf{x})$ is point identified. If $\beta(\mathbf{x}) \neq 0$, then $\alpha_0(\mathbf{x})$ and $\alpha_1(\mathbf{x})$ are likewise point identified.

Explicit Solution

$$\beta(\mathbf{x}) = \text{sign} [\theta_1(\mathbf{x})] \sqrt{3 [\theta_2(\mathbf{x})/\theta_1(\mathbf{x})]^2 - 2 [\theta_3(\mathbf{x})/\theta_1(\mathbf{x})]}$$

Sufficient for (II) and (III)

- (a) T is conditionally independent of (ε, z) given (T^*, \mathbf{x})
- (b) z is conditionally independent of ε given \mathbf{x}

Binary Regressors

└ Point Identification Result

Point Identification Result

Theorem

(Baseline) + (II) + (III) $\implies \beta(x)$ is point identified. If $\beta(x) \neq 0$, then $\alpha_0(x)$ and $\alpha_1(x)$ are likewise point identified.

Explicit Solution

$$\beta(x) = \text{sign}[\theta_1(x)] \sqrt{3[\theta_2(x)/\theta_1(x)]^2 - 2[\theta_3(x)/\theta_1(x)]}$$

Sufficient for (II) and (III)

- (a) T is conditionally independent of (x, z) given (T^*, x)
- (b) z is conditionally independent of x given x

Comment on the sufficient conditions: say that we really think these are what people have in mind in a natural experiment setting. Explain about reporting results in both logs and levels.

Inference for a Mis-classified Regressor

Weak Identification

- ▶ β small \Rightarrow moment equalities uninformative about (α_0, α_1) [▶ more](#)
- ▶ (α_0, α_1) could be on the boundary of the parameter space
- ▶ Also true of existing estimators that assume T^* exogenous

Our Approach

- ▶ Sharp identified set yields *inequality* moment restrictions that remain informative even if $\beta \approx 0$. [▶ more](#)
- ▶ Identification-robust inference with equality and inequality MCs.

Inference with Moment Equalities and Inequalities

Moment Conditions

$$\mathbb{E}[m_j(\mathbf{w}_i, \vartheta_0)] \geq 0, \quad j = 1, \dots, J$$

$$\mathbb{E}[m_j(\mathbf{w}_i, \vartheta_0)] = 0, \quad j = J+1, \dots, J+K$$

Test Statistic

$$T_n(\vartheta) = \sum_{j=1}^J \left[\frac{\sqrt{n} \bar{m}_{n,j}(\vartheta)}{\hat{\sigma}_{n,j}(\vartheta)} \right]^2 + \sum_{j=J+1}^{J+K} \left[\frac{\sqrt{n} \bar{m}_{n,j}(\vartheta)}{\hat{\sigma}_{n,j}(\vartheta)} \right]^2$$

Critical Value

- ▶ $\sqrt{n} \bar{\mathbf{m}}_n(\vartheta_0) \rightarrow_d$ normal limit with covariance matrix $\Sigma(\vartheta_0)$
- ▶ Use this to bootstrap the limit dist. of $T_n(\vartheta)$ under $H_0: \vartheta = \vartheta_0$

Binary Regressors

└ Inference with Moment Equalities and

Moment Conditions

$$E[m_j(w_i, \theta_0)] \geq 0, \quad j = 1, \dots, J$$

$$E[m_j(w_i, \theta_0)] = 0, \quad j = J+1, \dots, J+K$$

Test Statistic

$$T_n(\hat{\theta}) = \sum_{j=1}^J \left[\frac{\sqrt{n} \hat{m}_{nj}(\hat{\theta})}{\hat{\sigma}_{nj}(\hat{\theta})} \right]^2 + \sum_{j=J+1}^{J+K} \left[\frac{\sqrt{n} \hat{m}_{nj}(\hat{\theta})}{\hat{\sigma}_{nj}(\hat{\theta})} \right]^2$$

Critical Value

- $\sqrt{n} \hat{m}_{nj}(\theta_0) \rightarrow_d$ normal limit with covariance matrix $\Sigma(\theta_0)$
- Use this to bootstrap the limit dist. of $T_n(\hat{\theta})$ under $H_0: \theta = \theta_0$

Explain about the meaning of the m-var, the sigma-hat and the “minus” subscript

Generalized Moment Selection

Andrews & Soares (2010)

- ▶ Inequalities that don't bind reduce power of test, so eliminate those that are “far from binding” before calculating critical value:

$$\text{Drop inequality } j \text{ if } \frac{\sqrt{n} \bar{m}_{n,j}(\vartheta_0)}{\hat{\sigma}_{n,j}(\vartheta_0)} > \sqrt{\log n}$$

- ▶ Uniformly valid test of $H_0: \vartheta = \vartheta_0$ even if ϑ_0 is not point identified.
- ▶ Not asymptotically conservative.

Problem

Joint test for the whole parameter vector but we're only interested in β .
Projection is conservative and computationally intensive.

Generalized Moment Selection

- Inequalities that don't bind reduce power of test, so eliminate those that are "far from binding" before calculating critical value:

$$\text{Drop inequality } j \text{ if } \frac{\sqrt{n} \hat{m}_{n,j}(\hat{\theta}_n)}{\hat{\sigma}_{n,j}(\hat{\theta}_n)} > \sqrt{\log n}$$

- Uniformly valid test of $H_0: \theta = \theta_0$ even if θ_0 is not point identified.
- Not asymptotically conservative.

Problem

Joint test for the whole parameter vector but we're only interested in β .

Projection is conservative and computationally intensive.

Explain what not asymptotically conservative means. Explain what projection is and why it's conservative and computationally intensive.

Our Solution: Bonferroni-Based Inference

Special Structure

- ▶ β only enters MCs through $\theta_1 = \beta / (1 - \alpha_0 - \alpha_1)$
- ▶ Strong instrument \Rightarrow inference for θ_1 is standard.
- ▶ Nuisance pars γ strongly identified under null for (α_0, α_1)

Procedure

1. Concentrate out $(\theta_1, \gamma) \Rightarrow$ joint GMS test for (α_0, α_1)
2. Invert test $\Rightarrow (1 - \delta_1) \times 100\%$ confidence set for (α_0, α_1)
3. Project \Rightarrow CI for $(1 - \alpha_0 - \alpha_1)$
4. Construct standard $(1 - \delta_2) \times 100\%$ IV CI for θ_1
5. Bonferroni $\Rightarrow (1 - \delta_1 - \delta_2) \times 100\%$ CI for β

Binary Regressors

└ Our Solution: Bonferroni-Based Inference

Our Solution: Bonferroni-Based Inference

Special Structure

- β only enters MCs through $\theta_1 \Rightarrow \beta/(1 - \alpha_0 - \alpha_1)$
- Strong instrument \Rightarrow inference for θ_1 is standard.
- Nuisance pars γ strongly identified under null for (α_0, α_1)

Procedure


1. Concentrate out $(\theta_1, \gamma) \Rightarrow$ joint GMS test for (α_0, α_1)
2. Invert test $\Rightarrow (1 - \delta_1) \times 100\%$ confidence set for (α_0, α_1)
3. Project \Rightarrow CI for $(1 - \alpha_0 - \alpha_1)$
4. Construct standard $(1 - \delta_2) \times 100\%$ IV CI for θ_1
5. Bonferroni $\Rightarrow (1 - \delta_1 - \delta_2) \times 100\%$ CI for β

Explain that the procedure works well in simulations etc. Possibly add link to simulation here.

Example

(sim data: $\beta = 1, \alpha_0 = 0.1, \alpha_1 = 0.2, n = 5000$)

97.5% GMS Confidence Region for (α_0, α_1)



Bonferroni Interval

1. 97.5% CI for $(1 - \alpha_0 - \alpha_1) = (0.64, 0.82)$
2. 97.5% CI for $\theta_1 = (1.20, 1.47)$
3. $> 95\%$ CI for β :
 $(0.64 \times 1.20, 0.82 \times 1.47) = (0.77, 1.21)$

Comparisons

- ▶ $(0.88, 1.04)$ for IV if T^* were observed
- ▶ $(1.22, 1.45)$ for naive IV interval using T

Conclusion

This Paper

- ▶ Partial and point identification results for effect of binary, endogenous regressor using a valid instrument.
- ▶ Identification-robust inference in models with mis-classification

Related Work

- ▶ Relaxing Instrument Validity: “A Framework for Eliticing, Incorporating, and Disciplining Identification Beliefs in Linear Models” (with Camilo Garcia-Jimeno)
- ▶ Relaxing Non-differential Measurement Error: “Estimating the Returns to Lying” (with Arthur Lewbel)

Simple Bounds for Mis-classification from First-stage

Unobserved	Observed
$p_k^*(\mathbf{x}) \equiv \mathbb{P}(T^* = 1 \mathbf{x}, z = k)$	$p_k(\mathbf{x}) \equiv \mathbb{P}(T = 1 \mathbf{x}, z = k)$

Relationship

$$p_k^*(\mathbf{x}) = \frac{p_k(\mathbf{x}) - \alpha_0(\mathbf{x})}{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})}, \quad k = 0, 1$$

z does not affect (α_0, α_1) ; denominator $\neq 0$

Bounds for Mis-classification

$$\alpha_0(\mathbf{x}) \leq p_k(\mathbf{x}) \leq 1 - \alpha_1(\mathbf{x}), \quad k = 0, 1$$

$$\alpha_0(\mathbf{x}) + \alpha_1(\mathbf{x}) < 1$$

What does IV estimate under mis-classification?

Unobserved

$$\beta(\mathbf{x}) = \frac{\mathbb{E}[y|\mathbf{x}, z = 1] - \mathbb{E}[y|\mathbf{x}, z = 0]}{p_1^*(\mathbf{x}) - p_0^*(\mathbf{x})}$$

Wald (Observed)

$$\frac{\mathbb{E}[y|\mathbf{x}, z = 1] - \mathbb{E}[y|\mathbf{x}, z = 0]}{p_1(\mathbf{x}) - p_0(\mathbf{x})} = \beta(\mathbf{x}) \left[\frac{p_1^*(\mathbf{x}) - p_0^*(\mathbf{x})}{p_1(\mathbf{x}) - p_0(\mathbf{x})} \right] = \frac{\beta(\mathbf{x})}{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})}$$

$$p_1^*(\mathbf{x}) - p_0^*(\mathbf{x}) = \frac{p_1(\mathbf{x}) - \alpha_0(\mathbf{x})}{1 - \alpha_0 - \alpha_1(\mathbf{x})} - \frac{p_0(\mathbf{x}) - \alpha_0(\mathbf{x})}{1 - \alpha_0 - \alpha_1(\mathbf{x})} = \frac{p_1(\mathbf{x}) - p_0(\mathbf{x})}{1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})}$$

Partial Identification Bounds for $\beta(\mathbf{x})$

$$\beta(\mathbf{x}) = [1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x})] \left[\frac{\mathbb{E}[y|\mathbf{x}, z = 1] - \mathbb{E}[y|\mathbf{x}, z = 0]}{p_1(\mathbf{x}) - p_0(\mathbf{x})} \right]$$

$$0 \leq \alpha_0 \leq \min_k \{p_k(\mathbf{x})\}, \quad 0 \leq \alpha_1 \leq \min_k \{1 - p_k(\mathbf{x})\}$$

No Mis-classification

$$\alpha_0(\mathbf{x}) = \alpha_1(\mathbf{x}) = 0 \implies \beta(\mathbf{x}) = \text{Wald}$$

Maximum Mis-classification

$$\alpha_0(\mathbf{x}) = p_{\min}(\mathbf{x}), \quad \alpha_1(\mathbf{x}) = 1 - p_{\max}(\mathbf{x})$$

$$\implies 1 - \alpha_0(\mathbf{x}) - \alpha_1(\mathbf{x}) = p_{\max}(\mathbf{x}) - p_{\min}(\mathbf{x}) = |p_1(\mathbf{x}) - p_0(\mathbf{x})|$$

$$\implies \beta(\mathbf{x}) = \text{sign}\{p_1(\mathbf{x}) - p_0(\mathbf{x})\} \times (\text{Reduced Form})$$

Just-Identified System of Moment Equalities

Suppress dependence on $\mathbf{x} \dots$

$$\mathbb{E} \left[\{ \boldsymbol{\Psi}(\boldsymbol{\theta}) \mathbf{w}_i - \boldsymbol{\kappa} \} \otimes \begin{pmatrix} 1 \\ z \end{pmatrix} \right] = \mathbf{0}$$

$$\boldsymbol{\Psi}(\boldsymbol{\theta}) \equiv \begin{bmatrix} -\theta_1 & 1 & 0 & 0 & 0 & 0 \\ \theta_2 & 0 & -2\theta_1 & 1 & 0 & 0 \\ -\theta_3 & 0 & 3\theta_2 & 0 & -3\theta_1 & 1 \end{bmatrix}$$

$$\mathbf{w}_i = (T_i, y_i, y_i T_i, y_i^2, y_i^2 T_i, y_i^3)' \quad \theta_1 = \beta / (1 - \alpha_0 - \alpha_1)$$

$$\boldsymbol{\kappa} = (\kappa_1, \kappa_2, \kappa_3)' \quad \theta_2 = \theta_1^2 (1 + \alpha_0 - \alpha_1)$$

$$\theta_3 = \theta_1^3 [(1 - \alpha_0 - \alpha_1)^2 + 6\alpha_0(1 - \alpha_1)]$$

Moment Inequalities I – First-stage Probabilities

$\alpha_0 \leq p_k \leq 1 - \alpha_1$ becomes $\mathbb{E}[m(\mathbf{w}_i, \boldsymbol{\vartheta})] \geq \mathbf{0}$ for all k where

$$m(\mathbf{w}_i, \boldsymbol{\vartheta}) \equiv \begin{bmatrix} \mathbf{1}(z_i = k)(T - \alpha_0) \\ \mathbf{1}(z_i = k)(1 - T_i - \alpha_1) \end{bmatrix}$$

Moment Inequalities II – Non-differential Assumption

For all k , we have $\mathbb{E}[m(\mathbf{w}_i, \vartheta, \mathbf{q}_k)] \geq 0$ where

$$m(\mathbf{w}_i, \vartheta, \mathbf{q}_k) \equiv \begin{bmatrix} y_i \mathbf{1}(z_i = k) \left\{ (T_i - \alpha_0) - \mathbf{1}(y_i \leq \underline{q}_{0k})(1 - T_i) \left(\frac{1 - \alpha_0 - \alpha_1}{\alpha_1} \right) \right\} \\ -y_i \mathbf{1}(z_i = k) \left\{ (T_i - \alpha_0) - \mathbf{1}(y_i > \bar{q}_{0k})(1 - T_i) \left(\frac{1 - \alpha_0 - \alpha_1}{\alpha_1} \right) \right\} \\ y_i \mathbf{1}(z_i = k) \left\{ (T_i - \alpha_0) - \mathbf{1}(y_i \leq \underline{q}_{1k}) T_i \left(\frac{1 - \alpha_0 - \alpha_1}{1 - \alpha_1} \right) \right\} \\ -y_i \mathbf{1}(z_i = k) \left\{ (T_i - \alpha_0) - \mathbf{1}(y_i > \bar{q}_{1k}) T_i \left(\frac{1 - \alpha_0 - \alpha_1}{1 - \alpha_1} \right) \right\} \end{bmatrix}$$

and $\mathbf{q}_k \equiv (\underline{q}_{0k}, \bar{q}_{0k}, \underline{q}_{1k}, \bar{q}_{1k})'$ defined by $\mathbb{E}[h(\mathbf{w}_i, \vartheta, \mathbf{q}_k)] = 0$ with

$$h(\mathbf{w}_i, \vartheta, \mathbf{q}_k) = \begin{bmatrix} \mathbf{1}(y_i \leq \underline{q}_{0k}) \mathbf{1}(z_i = k)(1 - T_i) - \left(\frac{\alpha_1}{1 - \alpha_0 - \alpha_1} \right) \mathbf{1}(z_i = k)(T_i - \alpha_0) \\ \mathbf{1}(y_i \leq \bar{q}_{0k}) \mathbf{1}(z_i = k)(1 - T_i) - \left(\frac{1 - \alpha_0}{1 - \alpha_0 - \alpha_1} \right) \mathbf{1}(z_i = k)(1 - T_i - \alpha_1) \\ \mathbf{1}(y_i \leq \underline{q}_{1k}) \mathbf{1}(z_i = k) T_i - \left(\frac{1 - \alpha_1}{1 - \alpha_0 - \alpha_1} \right) \mathbf{1}(z_i = k)(T_i - \alpha_0) \\ \mathbf{1}(y_i \leq \bar{q}_{1k}) \mathbf{1}(z_i = k) T_i - \left(\frac{\alpha_0}{1 - \alpha_0 - \alpha_1} \right) \mathbf{1}(z_i = k)(1 - T_i - \alpha_1) \end{bmatrix}$$