

# Econ 722 – Advanced Econometrics IV

Francis J. DiTraglia

University of Pennsylvania

# Lecture #1 – Decision Theory

# Lecture #2 – Model Selection I

# Lecture #3 – Model Selection II

# Lecture #4 – Asymptotic Properties

# Lecture #5 – Andrews (1999) Moment Selection Criteria

# Lecture #6 – Focused Moment Selection

# Lecture #7 – High-Dimensional Regression I



# Lecture #8 – High-Dimensional Regression II

# Lecture #10 – Selective Inference

Optimal Inference After Model Selection (Fithian et al., 2017)

# How Statistics is Done In Reality

## Step 1: Selection – Decide what questions to ask.

“The analyst chooses a statistical model for the data at hand, and formulates testing, estimation, or other problems in terms of unknown aspects of that model.”

## Step 2: Inference – Answer the Questions.

“The analyst investigates the chosen problems using the data and the selected model.”

## Problem – “Data-snooping”

Standard techniques for (frequentist) statistical inference assume that we choose our questions **before** observing the data.

## Simple Example: “File Drawer Problem”

$Y_i \sim \text{iid } N(\mu_i, 1)$  for  $i = 1, \dots, n$

- ▶ I want to know which  $\mu_i \neq 0$ , but I'm busy and  $n$  is big.
- ▶ My RA looks at each  $Y_i$  and finds the “interesting” ones, namely  $\hat{\mathcal{I}} = \{i: |Y_i| > 1\}$ .
- ▶ I test  $H_{0,i}: \mu_i = 0$  against the two-sided alternative at the 5% significance level for each  $i \in \hat{\mathcal{I}}$ .

### Two Questions

1. What is the probability of falsely rejecting  $H_{0,i}$ ?
2. Among all  $H_{0,i}$  that I test, what fraction are false rejections?

## Simple Example: “File Drawer Problem”

$$\begin{aligned}\mathbb{P}_{H_{0,i}}(\{\text{Reject } H_{0,i}\}) &= \mathbb{P}_{H_{0,i}}(\{\text{Test } H_{0,i}\} \cap \{\text{Reject } H_{0,i}\}) \\&= \mathbb{P}_{H_{0,i}}(\{\text{Reject } H_{0,i}\} | \{\text{Test } H_{0,i}\}) \mathbb{P}_{H_{0,i}}(\{\text{Test } H_{0,i}\}) \\&= \mathbb{P}_{H_{0,i}}(|Y_i| > 1.96 | |Y_i| > 1) \mathbb{P}_{H_{0,i}}(|Y_i| > 1) \\&= \frac{2\Phi(-1.96)}{2\Phi(-1)} \times 2\Phi(-1) \\&\approx 0.16 \times 0.32 \approx 0.05\end{aligned}$$

$$\begin{aligned}\mathbb{P}_{H_{0,i}}(\{\text{Reject } H_{0,i}\} | \{\text{Test } H_{0,i}\}) &= \mathbb{P}_{H_{0,i}}(|Y_i| > 1.96 | |Y_i| > 1) \\&= \frac{\Phi(-1.96)}{\Phi(-1)} \approx 0.16\end{aligned}$$

# Simple Example: “File Drawer Problem”

## Conditional vs. Unconditional Type I Error Rates

- ▶ The **conditional** probability of falsely rejecting  $H_{0,i}$ , given that I have tested it, is about 0.16.
- ▶ The **unconditional** probability of falsely rejecting  $H_{0,i}$  is 0.05 since I only test a false null with probability 0.32.

## Idea for Post-Selection Inference

Control the Type I Error Rate **conditional on selection**: “The answer must be valid, given that the question was asked.”

## Simple Example: “File Drawer Problem”

### Conditional Type I Error Rate

Solve  $\mathbb{P}_{H_{0,i}}(\{|Y_i| > c\}|\{|Y_i| > 1\}) = 0.05$  for  $c$ .

$$\mathbb{P}_{H_{0,i}}(\{|Y_i| > c\}|\{|Y_i| > 1\}) = \frac{\Phi(-c)}{\Phi(-1)} = 0.05$$

$$c = -\Phi^{-1}(\Phi(-1) \times 0.05)$$

$$c \approx 2.41$$

### Notice:

To account for the first-stage selection step, we need a larger critical value: 2.41 vs. 1.96. This means the test is less powerful.

# Selective Inference vs. Sample-Splitting

## Classical Inference

Control the Type I error under model  $M$ :  $\mathbb{P}_{M, H_0}(\text{reject } H_0) \leq \alpha$ .

## Selective Inference

Control the Type I error under model  $M$ , **given** that  $M$  and  $H_0$  were selected:  $\mathbb{P}_{M, H_0}(\text{reject } H_0 | \{M, H_0 \text{ selected}\}) \leq \alpha$ .

## Sample-Splitting

Use different datasets to choose  $(M, H_0)$  and carry out inference:

$$\mathbb{P}_{M, H_0}(\text{reject } H_0 | \{M, H_0 \text{ selected}\}) = \mathbb{P}_{M, H_0}(\text{reject } H_0).$$



# Selective Inference in Exponential Family Models

## Questions

1. Recipe for selective inference in realistic examples?
2. How to construct the “best” selective test in a given example?
3. How does selective inference compare to sample-splitting?

## Fithian, Sun & Taylor (2017)

- ▶ Use classical theory for exponential family models (Lehmann & Scheffé).
- ▶ Computational procedure for UMPU selective test/CI after arbitrary model/hypothesis selection.
- ▶ Sample-splitting is typically inadmissible (wastes information).
- ▶ Example: post-selection inference for high-dimensional regression

# A Prototype Example of Selective Inference

This is my own example, but uses the same idea that underlies Fithian et al.

- ▶ Choose between models M1 and M2 based on parameter  $\delta$ .
  - ▶ If  $\delta \neq 0$ , choose M1
  - ▶ If  $\delta = 0$ , choose M2
  - ▶ E.g.  $\delta$  is the endogeneity of  $X$ , M1 is IV and M2 is OLS
- ▶ Observe  $Y_\delta \sim N(\delta, 1)$  and use this to choose a model.
  - ▶ Selection Event:  $A \equiv \{|Y_\delta| > c\}$ , for some critical value  $c$
  - ▶ If  $A$ , then choose M1. Otherwise, choose M2.
- ▶ After choosing a model, carry out inference for  $\beta$ .
  - ▶ Under a particular model  $M$ ,  $Y_\beta \sim N(\beta, 1)$
  - ▶  $\beta$  is a *model-specific* parameter