

# Periodic solutions of Euler-Lagrange equations with “sublinear nonlinearity” in an Orlicz-Sobolev space setting

Sonia Acinas \*

Instituto de Matemática Aplicada San Luis (CONICET-UNSL)  
(5700) San Luis, Argentina  
Universidad Nacional de La Pampa  
(6300) Santa Rosa, La Pampa, Argentina  
sonia.acinas@gmail.com

Fernando D. Mazzone †

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales  
Universidad Nacional de Río Cuarto  
(5800) Río Cuarto, Córdoba, Argentina,  
fmazzone@exa.unrc.edu.ar

## Abstract

In this paper we....

## 1 Introduction

This paper is concerned with the existence of periodic solutions of the problem

$$\begin{cases} \frac{d}{dt} D_y \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) = D_x \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) & \text{a.e. } t \in (0, T) \\ \mathbf{u}(0) - \mathbf{u}(T) = \dot{\mathbf{u}}(0) - \dot{\mathbf{u}}(T) = 0 \end{cases} \quad (1)$$

---

\*SECyT-UNRC, UNSL and CONICET

†SECyT-UNRC and CONICET

**2010 AMS Subject Classification.** Primary: . Secondary: .

**Keywords and phrases.** .

where  $T > 0$ ,  $\mathbf{u} : [0, T] \rightarrow \mathbb{R}^d$  is absolutely continuous and the *Lagrangian*  $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function satisfying the conditions

$$|\mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( b(t) + \Phi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right), \quad (2)$$

$$|D_{\mathbf{x}}\mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( b(t) + \Phi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right), \quad (3)$$

$$|D_{\mathbf{y}}\mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( c(t) + \varphi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right). \quad (4)$$

In these inequalities we assume that  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $\lambda > 0$ ,  $\Phi$  is an  $N$ -function (see section Preliminaries for definitions),  $\varphi$  is the right continuous derivative of  $\Phi$  and the non negative functions  $b$ ,  $c$  and  $f$  belong to certain Banach spaces that will be introduced later.

It is well known that problem (1) comes from a variational one, that is, a solution of (1) is a critical point of the *action integral*

$$I(\mathbf{u}) = \int_0^T \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) dt. \quad (5)$$

Variational problems and hamiltonian systems have been studied extensively. Classic references of these subjects are [21, 26, 13]. Problems like (1) have maintained the interest of researchers as the recent literature on the topic testifies. For lagrangian functions of the type  $\mathcal{L}(t, \mathbf{x}, \mathbf{y}) = \frac{|\mathbf{y}|^2}{2} + F(t, \mathbf{x})$  many solvability conditions have been given expanding the results of [21]. In [28] the function  $F$  was split up into two potentials, one of them with a property of subadditivity and the other with a bounded gradient. In [27] it was required a certain sublinearity condition on the gradient of the potential  $F$ ; and, in [32] it was considered a potential  $F$  given by a sum of a subconvex function and a subquadratic one. In [29] the uniform coercivity of  $\int_0^T F(t, \mathbf{x}) dt$  was replaced by local coercivity of  $F$  in some positive measure subset of  $[0, T]$ . In [35], the authors took a similar potential to that in [32] getting new solvability conditions and they also studied the case in which the two potentials do not have any convexity.

The Lagrangian  $\mathcal{L}(t, \mathbf{x}, \mathbf{y}) = \frac{|\mathbf{y}|^p}{p} + F(t, \mathbf{x})$  for  $p > 1$  was treated in more recent papers. By using the dual least action principle, in [31] it was performed the extension of some results given in [21]; and, in [30] the authors improved the work done in [32]. On the other hand, by the minimax methods in critical point theory some existence theorems were obtained. In [33] it was employed a subquadratic potential  $F$  in Rabinowitz's sense and in [34]  $F$  was taken as in [27].

Another source of problems, close to our proposal, is the one in which a *p-laplacian-like* operator is involved. Assuming that the function  $\varphi$  is a homeomorphism from  $\mathbb{R}^d$  into itself, it is considered the differential operator  $\mathbf{u} \mapsto (\varphi(\mathbf{u}'))'$ . In [4, 5, 6, 19, 20], using the Leray-Schauder degree theory, some existence results of solutions of equations like  $(\varphi(\mathbf{u}'))' = \mathbf{f}(t, \mathbf{u}(t), \mathbf{u}'(t))$  were obtained under different boundary conditions (periodic, Dirichlet, von Neumann) and where  $\mathbf{f}$  is not necessarily a gradient. We point out that our approach differs from that of previous articles because we tackle the direct method of the calculus of variations.

In the Orlicz-Sobolev space setting, in [14] a constrained minimization problem associated to the existence of eigenvalues for certain differential operators involving

$N$ -functions was studied. Slightly away from the problems to be treated in this paper, we can mention [8, 10] where A. Cianchi dealt with the regularity of minimizers of action integrals defined on several variable functions.

In this article we consider lagrangian functions defined on Orlicz-Sobolev spaces  $W^1 L^\Phi$  (see [3, 17, 22, 23]) and we use the direct method of calculus of variations. The exposition is organized as follows. In Section 2 we enumerate results related to Orlicz spaces, Orlicz-Sobolev spaces and composition operators. Almost all results in this section are essentially known. Conditions (2), (3) and (4) are the means to ensure that  $I$  is finitely defined on a non trivial subset of  $W^1 L_d^\Phi$  and  $I$  is Gâteaux differentiable in this subset. We develop these issues in Theorem 2.5 of Section ???. In Section ??? we prove that critical points of (5) are solutions of (1). Conditions to guarantee the coercitivity of action integrals are discussed in Section 3. Finally, our main theorem about existence of solutions of (1) is introduced and proved in Section ???.

We lay emphasis on that we use  $\Delta_2$ -condition only when necessary in a certain sense (see, for example, Lemma 3.2).

## 2 Preliminaries

For reader convenience, we give a short introduction to Orlicz and Orlicz-Sobolev spaces of vector valued functions and a list of results that we will use throughout the article. Classic references for Orlicz spaces of real valued functions are [3, 17, 22]. For Orlicz spaces of vector valued functions, see [25] and the references therein.

Hereafter we denote by  $\mathbb{R}^+$  the set of all non negative real numbers. A function  $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is called an  $N$ -function if  $\Phi$  is given by

$$\Phi(t) = \int_0^t \varphi(\tau) d\tau, \quad \text{for } t \geq 0,$$

where  $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a right continuous non decreasing function satisfying  $\varphi(0) = 0$ ,  $\varphi(t) > 0$  for  $t > 0$  and  $\lim_{t \rightarrow \infty} \varphi(t) = +\infty$ .

Given a function  $\varphi$  as above, we consider the so-called right inverse function  $\psi$  of  $\varphi$  which is defined by  $\psi(s) = \sup_{\varphi(t) \leq s} t$ . The function  $\psi$  satisfies the same properties as the function  $\varphi$ , therefore we have an  $N$ -function  $\Psi$  such that  $\Psi' = \psi$ . The function  $\Psi$  is called the *complementary function* of  $\Phi$ .

We say that  $\Phi$  satisfies the  $\Delta_2$ -condition, denoted by  $\Phi \in \Delta_2$ , if there exist constants  $K > 0$  and  $t_0 \geq 0$  such that

$$\Phi(2t) \leq K\Phi(t) \tag{6}$$

for every  $t \geq t_0$ . If  $t_0 = 0$ , we say that  $\Phi$  satisfies the  $\Delta_2$ -condition globally ( $\Phi \in \Delta_2$  globally).

Let  $d$  be a positive integer. We denote by  $\mathcal{M}_d := \mathcal{M}_d([0, T])$  the set of all measurable functions defined on  $[0, T]$  with values on  $\mathbb{R}^d$  and we write  $\mathbf{u} = (u_1, \dots, u_d)$  for  $\mathbf{u} \in \mathcal{M}_d$ . In this paper we adopt the convention that bold symbols denote points in  $\mathbb{R}^d$ .

Given an  $N$ -function  $\Phi$  we define the *modular function*  $\rho_\Phi : \mathcal{M}_d \rightarrow \mathbb{R}^+ \cup \{+\infty\}$  by

$$\rho_\Phi(\mathbf{u}) := \int_0^T \Phi(|\mathbf{u}|) dt.$$

Here  $|\cdot|$  is the euclidean norm of  $\mathbb{R}^d$ . The *Orlicz class*  $C_d^\Phi = C_d^\Phi([0, T])$  is given by

$$C_d^\Phi := \{\mathbf{u} \in \mathcal{M}_d | \rho_\Phi(\mathbf{u}) < \infty\}. \quad (7)$$

The *Orlicz space*  $L_d^\Phi = L_d^\Phi([0, T])$  is the linear hull of  $C_d^\Phi$ ; equivalently,

$$L_d^\Phi := \{\mathbf{u} \in \mathcal{M}_d | \exists \lambda > 0 : \rho_\Phi(\lambda \mathbf{u}) < \infty\}. \quad (8)$$

The Orlicz space  $L_d^\Phi$  equipped with the *Orlicz norm*

$$\|\mathbf{u}\|_{L^\Phi} := \sup \left\{ \int_0^T \mathbf{u} \cdot \mathbf{v} dt \mid \rho_\Psi(\mathbf{v}) \leq 1 \right\},$$

is a Banach space. By  $\mathbf{u} \cdot \mathbf{v}$  we denote the usual dot product in  $\mathbb{R}^d$  between  $\mathbf{u}$  and  $\mathbf{v}$ . The following alternative expression for the norm, known as *Amemiya norm*, will be useful (see [17, Thm. 10.5] and [15]). For every  $\mathbf{u} \in L^\Phi$ ,

$$\|\mathbf{u}\|_{L^\Phi} = \inf_{k>0} \frac{1}{k} \{1 + \rho_\Phi(k\mathbf{u})\}. \quad (9)$$

The subspace  $E_d^\Phi = E_d^\Phi([0, T])$  is defined as the closure in  $L_d^\Phi$  of the subspace  $L_d^\infty$  of all  $\mathbb{R}^d$ -valued essentially bounded functions. It is shown that  $E_d^\Phi$  is the only one maximal subspace contained in the Orlicz class  $C_d^\Phi$ , i.e.  $\mathbf{u} \in E_d^\Phi$  if and only if  $\rho_\Phi(\lambda \mathbf{u}) < \infty$  for any  $\lambda > 0$ .

A generalized version of *Hölder's inequality* holds in Orlicz spaces. Namely, if  $\mathbf{u} \in L_d^\Phi$  and  $\mathbf{v} \in L_d^\Psi$  then  $\mathbf{u} \cdot \mathbf{v} \in L_1^1$  and

$$\int_0^T \mathbf{v} \cdot \mathbf{u} dt \leq \|\mathbf{u}\|_{L^\Phi} \|\mathbf{v}\|_{L^\Psi}. \quad (10)$$

If  $X$  and  $Y$  are Banach spaces such that  $Y \subset X^*$ , we denote by  $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{R}$  the bilinear pairing map given by  $\langle x^*, x \rangle = x^*(x)$ . Hölder's inequality shows that  $L_d^\Psi \subset [L_d^\Phi]^*$ , where the pairing  $\langle \mathbf{v}, \mathbf{u} \rangle$  is defined by

$$\langle \mathbf{v}, \mathbf{u} \rangle = \int_0^T \mathbf{v} \cdot \mathbf{u} dt \quad (11)$$

with  $\mathbf{u} \in L_d^\Phi$  and  $\mathbf{v} \in L_d^\Psi$ . Unless  $\Phi \in \Delta_2$ , the relation  $L_d^\Psi = [L_d^\Phi]^*$  will not hold. In general, it is true that  $[E_d^\Phi]^* = L_d^\Psi$ .

Like in [17], we will consider the subset  $\Pi(E_d^\Phi, r)$  of  $L_d^\Phi$  given by

$$\Pi(E_d^\Phi, r) := \{\mathbf{u} \in L_d^\Phi | d(\mathbf{u}, E_d^\Phi) < r\}.$$

This set is related to the Orlicz class  $C_d^\Phi$  by means of inclusions, namely,

$$\Pi(E_d^\Phi, r) \subset rC_d^\Phi \subset \overline{\Pi(E_d^\Phi, r)} \quad (12)$$

for any positive  $r$ . If  $\Phi \in \Delta_2$ , then the sets  $L_d^\Phi$ ,  $E_d^\Phi$ ,  $\Pi(E_d^\Phi, r)$  and  $C_d^\Phi$  are equal.

We define the *Sobolev-Orlicz space*  $W^1 L_d^\Phi$  (see [3]) by

$$W^1 L_d^\Phi := \{u \mid u \text{ is absolutely continuous and } u, \dot{u} \in L_d^\Phi\}.$$

$W^1 L_d^\Phi$  is a Banach space when equipped with the norm

$$\|u\|_{W^1 L^\Phi} = \|u\|_{L^\Phi} + \|\dot{u}\|_{L^\Phi}.$$

For a function  $u \in L_d^1([0, T])$ , we write  $u = \bar{u} + \tilde{u}$  where  $\bar{u} = \frac{1}{T} \int_0^T u(t) dt$  and  $\tilde{u} = u - \bar{u}$ .

As usual, if  $(X, \|\cdot\|_X)$  is a Banach space and  $(Y, \|\cdot\|_Y)$  is a subspace of  $X$ , we write  $Y \hookrightarrow X$  and we say that  $Y$  is *embedded* in  $X$  when the restricted identity map  $i_Y : Y \rightarrow X$  is bounded. That is, there exists  $C > 0$  such that for any  $y \in Y$  we have  $\|y\|_X \leq C\|y\|_Y$ . With this notation, Hölder's inequality states that  $L_d^\Psi \hookrightarrow [L_d^\Phi]^*$ ; and, it is easy to see that for every  $N$ -function  $\Phi$  we have that  $L_d^\infty \hookrightarrow L_d^\Phi \hookrightarrow L_d^1$ .

Recall that a function  $w : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is called a *modulus of continuity* if  $w$  is a continuous increasing function which satisfies  $w(0) = 0$ . For example, it can be easily shown that  $w(s) = s\Phi^{-1}(1/s)$  is a modulus of continuity for every  $N$ -function  $\Phi$ . We say that  $u : [0, T] \rightarrow \mathbb{R}^d$  has modulus of continuity  $w$  when there exists a constant  $C > 0$  such that

$$|u(t) - u(s)| \leq Cw(|t - s|). \quad (13)$$

We denote by  $C^w([0, T], \mathbb{R}^d)$  the space of  $w$ -Hölder continuous functions. This is the space of all functions satisfying (13) for some  $C > 0$  and it is a Banach space with norm

$$\|u\|_{C^w([0, T], \mathbb{R}^d)} := \|u\|_{L^\infty} + \sup_{t \neq s} \frac{|u(t) - u(s)|}{w(|t - s|)}.$$

An important aspect of the theory of Sobolev spaces is related to embedding theorems. There is an extensive literature on this question in the Orlicz-Sobolev space setting, see for example [7, 9, 11, 12, 16]. The next simple lemma is essentially known and we will use it systematically. For the sake of completeness, we include a brief proof of it.

**Lemma 2.1.** *Let  $w(s) := s\Phi^{-1}(1/s)$ . Then, the following statements hold:*

1.  $W^1 L^\Phi \hookrightarrow C^w([0, T], \mathbb{R}^d)$  and for every  $u \in W^1 L^\Phi$

$$|u(t) - u(s)| \leq \|\dot{u}\|_{L^\Phi} w(|t - s|), \quad (14)$$

$$\|u\|_{L^\infty} \leq \Phi^{-1}\left(\frac{1}{T}\right) \max\{1, T\} \|u\|_{W^1 L^\Phi} \quad (15)$$

2. For every  $u \in W^1 L^\Phi$  we have  $\tilde{u} \in L_d^\infty$  and

$$\|\tilde{u}\|_{L^\infty} \leq T\Phi^{-1}\left(\frac{1}{T}\right) \|\dot{u}\|_{L^\Phi} \quad (\text{Sobolev's inequality}). \quad (16)$$

*Proof.* For  $0 \leq s \leq t \leq T$ , we get

$$\begin{aligned} |\mathbf{u}(t) - \mathbf{u}(s)| &\leq \int_s^t |\dot{\mathbf{u}}(\tau)| \, d\tau \\ &\leq \|\chi_{[s,t]}\|_{L^\Psi} \|\dot{\mathbf{u}}\|_{L^\Phi} \\ &= \|\dot{\mathbf{u}}\|_{L^\Phi} (t-s) \Phi^{-1}\left(\frac{1}{t-s}\right), \end{aligned} \tag{17}$$

using Hölder's inequality and [17, Eq. (9.11)]. This proves the inequality (14).

Since  $u_i$  is continuous, from Mean Value Theorem for integrals, there exists  $s_i \in [0, T]$  such that  $u_i(s_i) = \bar{u}_i$ . Using this  $s_i$  value in (14) with  $u_i$  instead of  $\mathbf{u}$  and taking into account that  $s\Phi^{-1}(1/s)$  is increasing, we obtain Sobolev's inequality for each  $u_i$ . The inequality (16) follows easily from the corresponding result for each component of  $\mathbf{u}$ .

On the other hand, again by Hölder's inequality and [17, Eq. (9.11)], we have

$$|\bar{\mathbf{u}}| = \frac{1}{T} \int_0^T |\mathbf{u}(s)| \, ds \leq \Phi^{-1}\left(\frac{1}{T}\right) \|\mathbf{u}\|_{L^\Phi}. \tag{18}$$

From (16), (18) and the fact that  $\mathbf{u} = \bar{\mathbf{u}} + \tilde{\mathbf{u}}$ , we obtain (15). This completes the proof of item 1.  $\square$

The next result is analogous to some lemmata in  $W^1 L_d^p$ , see Xu-Tang-2007 y otros...

**Lemma 2.2.** *If  $\|\mathbf{u}\|_{W^1 L^\Phi} \rightarrow \infty$ , then  $(|\bar{\mathbf{u}}| + \|\dot{\mathbf{u}}\|_{L^\Phi}) \rightarrow \infty$ .*

*Proof.* We have

$$\|\mathbf{u}\|_{L^\Phi} = \|\bar{\mathbf{u}} + \tilde{\mathbf{u}}\|_{L^\Phi} \leq \|\bar{\mathbf{u}}\|_{L^\Phi} + \|\tilde{\mathbf{u}}\|_{L^\Phi} = |\bar{\mathbf{u}}| 1_{L^\Phi} + \|\tilde{\mathbf{u}}\|_{L^\Phi}$$

We know that Hölder's inequality implies that  $L_d^\infty \hookrightarrow L_d^\Phi$ , that is, there exists  $C > 0$  such that for any  $\tilde{\mathbf{u}} \in L_d^\infty$  we have

$$\|\tilde{\mathbf{u}}\|_{L^\Phi} \leq C \|\tilde{\mathbf{u}}\|_{L^\infty}$$

and, applying Sobolev's inequality to the previous formula, we get

$$\|\tilde{\mathbf{u}}\|_{L^\Phi} \leq C \|\dot{\mathbf{u}}\|_{L^\Phi}$$

**La desigualdad anterior sería del tipo Wirtinger's que no tenemos enunciada en ningún lado.**

Therefore,

$$\|\mathbf{u}\|_{L^\Phi} \leq C(|\bar{\mathbf{u}}| + \|\dot{\mathbf{u}}\|_{L^\Phi}) \tag{19}$$

As  $\|\mathbf{u}\|_{W^1 L^\Phi} = \|\mathbf{u}\|_{L^\Phi} + \|\dot{\mathbf{u}}\|_{L^\Phi}$ , then

$$\|\mathbf{u}\|_{W^1 L^\Phi} \leq C(|\bar{\mathbf{u}}| + \|\dot{\mathbf{u}}\|_{L^\Phi})$$

and by hypothesis  $\|\mathbf{u}\|_{W^1 L^\Phi} \rightarrow \infty$ , then  $|\bar{\mathbf{u}}| + \|\dot{\mathbf{u}}\|_{L^\Phi} \rightarrow \infty$ .  $\square$

Given a continuous function  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$ , we define the composition operator  $\mathbf{a} : \mathcal{M}_d \rightarrow \mathcal{M}_d$  by  $\mathbf{a}(\mathbf{u})(t) = a(|\mathbf{u}(t)|)$ . We will often use the following elementary consequence of Lemma 2.1.

**Corollary 2.3.** *If  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$  then  $\mathbf{a} : W^1 L_d^\Phi \rightarrow L_1^\infty([0, T])$  is bounded. More concretely, there exists a non decreasing function  $A : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\|\mathbf{a}(\mathbf{u})\|_{L^\infty([0, T])} \leq A(\|\mathbf{u}\|_{W^1 L^\Phi})$ .*

*Proof.* Let  $\alpha \in C(\mathbb{R}^+, \mathbb{R}^+)$  be a non decreasing majorant of  $a$ , for example  $\alpha(s) := \sup_{0 \leq t \leq s} a(t)$ . If  $\mathbf{u} \in W^1 L_d^\Phi$  then, by Lemma 2.1,

$$a(|\mathbf{u}(t)|) \leq \alpha(\|\mathbf{u}\|_{L^\infty}) \leq \alpha\left(\Phi^{-1}\left(\frac{1}{T}\right) \max\{1, T\} \|\mathbf{u}\|_{W^1 L^\Phi}\right) =: A(\|\mathbf{u}\|_{W^1 L^\Phi}).$$

□

**Definition 2.4.** *We say that a function  $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function if for fixed  $(\mathbf{x}, \mathbf{y})$  the map  $t \mapsto \mathcal{L}(t, \mathbf{x}, \mathbf{y})$  is measurable and for fixed  $t$  the map  $(\mathbf{x}, \mathbf{y}) \mapsto \mathcal{L}(t, \mathbf{x}, \mathbf{y})$  is continuously differentiable for almost everywhere  $t \in [0, T]$ .*

In [1] we proved the next result.

**Theorem 2.5.** *Let  $\mathcal{L}$  be a Carathéodory function satisfying (2), (3) and (4). Then the following statements hold:*

1. *The action integral given by (5) is finitely defined on  $\mathcal{E}_d^\Phi(\lambda) := W^1 L_d^\Phi \cap \{\mathbf{u} | \dot{\mathbf{u}} \in \Pi(E_d^\Phi, \lambda)\}$ .*
2. *The function  $I$  is Gâteaux differentiable on  $\mathcal{E}_d^\Phi(\lambda)$  and its derivative  $I'$  is demi-continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $[W^1 L_d^\Phi]^*$ . Moreover,  $I'$  is given by the following expression*

$$\langle I'(\mathbf{u}), \mathbf{v} \rangle = \int_0^T \{D_{\mathbf{x}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} + D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \dot{\mathbf{v}}\} dt. \quad (20)$$

3. *If  $\Psi \in \Delta_2$  then  $I'$  is continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $[W^1 L_d^\Phi]^*$  when both spaces are equipped with the strong topology.*

In [1] we derived the Euler-Lagrange equations associated to critical points of action integrals on the subspace of  $T$ -periodic functions. We denote by  $W^1 L_T^\Phi$  the subspace of  $W^1 L_d^\Phi$  containing all  $T$ -periodic functions. As usual, when  $Y$  is a subspace of the Banach space  $X$ , we denote by  $Y^\perp$  the *annihilator subspace* of  $X^*$ , i.e. the subspace that consists of all bounded linear functions which are identically zero on  $Y$ .

We recall that a function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is called *strictly convex* if  $f\left(\frac{\mathbf{x}+\mathbf{y}}{2}\right) < \frac{1}{2}(f(\mathbf{x}) + f(\mathbf{y}))$  for  $\mathbf{x} \neq \mathbf{y}$ . It is well known that if  $f$  is a strictly convex and differentiable function, then  $D_{\mathbf{x}} f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a one-to-one map (see, e.g. [24, Thm. 12.17]).

**Theorem 2.6.** *Let  $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$  be a  $T$ -periodic function. The following statements are equivalent:*

1.  $I'(\mathbf{u}) \in (W^1 L_T^\Phi)^\perp$ .
2.  $D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t))$  is an absolutely continuous function and  $\mathbf{u}$  solves the following boundary value problem

$$\begin{cases} \frac{d}{dt} D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) = D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) & \text{a.e. } t \in (0, T) \\ \mathbf{u}(0) - \mathbf{u}(T) = D_{\mathbf{y}}\mathcal{L}(0, \mathbf{u}(0), \dot{\mathbf{u}}(0)) - D_{\mathbf{y}}\mathcal{L}(T, \mathbf{u}(T), \dot{\mathbf{u}}(T)) = 0. \end{cases} \quad (21)$$

Moreover if  $D_{\mathbf{y}}\mathcal{L}(t, x, y)$  is  $T$ -periodic with respect to the variable  $t$  and strictly convex with respect to  $\mathbf{y}$ , then  $D_{\mathbf{y}}\mathcal{L}(0, \mathbf{u}(0), \dot{\mathbf{u}}(0)) - D_{\mathbf{y}}\mathcal{L}(T, \mathbf{u}(T), \dot{\mathbf{u}}(T)) = 0$  is equivalent to  $\dot{\mathbf{u}}(0) = \dot{\mathbf{u}}(T)$ .

### 3 Coercivity discussion

DECIR ALGO DE LOS ÍNDICES ACÁ O EN LA INTRO...??? We recall a usual definition in the context of calculus of variations.

**Definition 3.1.** Let  $X$  be a Banach space and let  $D$  be an unbounded subset of  $X$ . Suppose that  $J : D \subset X \rightarrow \mathbb{R}$ . We say that  $J$  is coercive if  $J(u) \rightarrow +\infty$  when  $\|u\|_X \rightarrow +\infty$ .

It is well known that coercivity is a useful ingredient in the process of establishing existence of minima. Therefore, we are interested in finding conditions which ensure the coercivity of the action integral  $I$  acting on  $\mathcal{E}_d^\Phi(\lambda)$ . For this purpose, we need to introduce the following extra condition on the lagrangian function  $\mathcal{L}$

$$\mathcal{L}(t, \mathbf{x}, \mathbf{y}) \geq \alpha_0 \Phi\left(\frac{|\mathbf{y}|}{\Lambda}\right) + F(t, \mathbf{x}), \quad (22)$$

where  $\alpha_0, \Lambda > 0$  and  $F : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function, i.e.  $F(t, \mathbf{x})$  is measurable with respect to  $t$  for every fixed  $\mathbf{x} \in \mathbb{R}^d$  and it is continuous at  $\mathbf{x}$  for a.e.  $t \in [0, T]$ . We observe that, from (22) and (2), we have  $F(t, \mathbf{x}) \leq a(|\mathbf{x}|)b_0(t)$  with  $b_0(t) := b(t) + \Phi(f(t)) \in L_1^1([0, T])$ . In order to guarantee that integral  $\int_0^T F(t, \mathbf{u}) dt$  is finite for  $\mathbf{u} \in W^1 L^\Phi$ , we need to assume

$$|F(t, \mathbf{x})| \leq a(|\mathbf{x}|)b_0(t), \quad \text{for a.e. } t \in [0, T] \quad \text{and for every } \mathbf{x} \in \mathbb{R}^d. \quad (23)$$

As we shall see in Theorem 3.3, when  $\mathcal{L}$  satisfies (2), (3), (4), (22) and (23), the coercivity of the action integral  $I$  is related to the coercivity of the functional

$$J_{C, \nu}(\mathbf{u}) := \rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right) - C\|\mathbf{u}\|_{L^\Phi}^\nu, \quad (24)$$

for  $C, \nu > 0$ . If  $\Phi(x) = |x|^p/p$  then  $J_{C, \nu}$  is clearly coercive for  $\nu < p$ . For more general  $\Phi$  the situation is more interesting as it will be shown in the following lemma.

**Lemma 3.2.** Let  $\Phi$  and  $\Psi$  be complementary  $N$ -functions. Then:

1. If  $C\Lambda < 1$ , then  $J_{C, 1}$  is coercive.



2. If  $\Psi \in \Delta_2$  globally, then there exists a constant  $\alpha_\Phi > 1$  such that, for any  $0 < \mu < \alpha_\Phi$ ,

$$\lim_{\|\mathbf{u}\|_{L^\Phi} \rightarrow \infty} \frac{\rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right)}{\|\mathbf{u}\|_{L^\Phi}^\mu} = +\infty. \quad (25)$$

In particular, the functional  $J_{C,\mu}$  is coercive for every  $C > 0$  and  $0 < \mu < \alpha_\Phi$ . The constant  $\alpha_\Phi$  is one of the so-called Matuszewska-Orlicz indices (see [18, Ch. 11]).

3. If  $J_{C,1}$  is coercive with  $C\Lambda > 1$ , then  $\Psi \in \Delta_2$ .

*Proof.* By (9) we have

$$(1 - C\Lambda)\|\mathbf{u}\|_{L^\Phi} + C\Lambda\|\mathbf{u}\|_{L^\Phi} = \|\mathbf{u}\|_{L^\Phi} \leq \Lambda + \Lambda\rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right),$$

then

$$\frac{(1 - C\Lambda)}{\Lambda}\|\mathbf{u}\|_{L^\Phi} - 1 \leq \rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right) - C\|\mathbf{u}\|_{L^\Phi} = J_{C,1}(\mathbf{u}).$$

This inequality shows that  $J_{C,1}$  is coercive and therefore item 1 is proved.

In virtue of [2, Eq. (2.8)], the  $\Delta_2$ -condition on  $\Psi$ , [18, Thm. 11.7] and [18, Cor. 11.6], we obtain constants  $K > 0$  and  $\alpha_\Phi > 1$  such that

$$\Phi(rs) \geq Kr^\nu\Phi(s) \quad (26)$$

for any  $0 < \nu < \alpha_\Phi$ ,  $s \geq 0$  and  $r > 1$ .

Let  $1 < \mu < \nu < \alpha_\Phi$  and let  $r > \Lambda$  be a constant that will be specified later. Then, from (26) and (9), we get

$$\begin{aligned} \frac{\int_0^T \Phi\left(\frac{|\mathbf{u}|}{\Lambda}\right) dt}{\|\mathbf{u}\|_{L^\Phi}^\mu} &\geq K\left(\frac{r}{\Lambda}\right)^\nu \frac{\int_0^T \Phi(r^{-1}|\mathbf{u}|) dt}{\|\mathbf{u}\|_{L^\Phi}^\mu} \\ &\geq K\left(\frac{r}{\Lambda}\right)^\nu \frac{r^{-1}\|\mathbf{u}\|_{L^\Phi} - 1}{\|\mathbf{u}\|_{L^\Phi}^\mu}. \end{aligned}$$

We choose  $r = \|\mathbf{u}\|_{L^\Phi}/2$ . Since  $\|\mathbf{u}\|_{L^\Phi} \rightarrow +\infty$  we can assume  $\|\mathbf{u}\|_{L^\Phi} > 2\Lambda$ . Thus  $r > \Lambda$  and

$$\frac{\int_0^T \Phi\left(\frac{|\mathbf{u}|}{\Lambda}\right) dt}{\|\mathbf{u}\|_{L^\Phi}^\mu} \geq \frac{K}{2^\nu \Lambda^\nu} \|\mathbf{u}\|_{L^\Phi}^{\nu-\mu} \rightarrow +\infty \quad \text{as } \|\mathbf{u}\|_{L^\Phi} \rightarrow +\infty,$$

because  $\nu > \mu$ .

With the aim of proving item 3, we suppose that  $\Psi \notin \Delta_2$ . By [17, Thm. 4.1], there exists a sequence of real numbers  $r_n$  such that  $r_n \rightarrow \infty$  and

$$\lim_{n \rightarrow \infty} \frac{r_n \psi(r_n)}{\Psi(r_n)} = +\infty. \quad (27)$$

Now, we choose  $\mathbf{u}_n$  such that  $|\mathbf{u}_n| = \Lambda \psi(r_n) \chi_{[0, \frac{1}{\Psi(r_n)}]}$ . Then, by [17, Eq. (9.11)], we get

$$\|\mathbf{u}_n\|_{L^\Phi} = \Lambda \frac{\psi(r_n)}{\Psi(r_n)} \Psi^{-1}(\Psi(r_n)) = \Lambda \frac{r_n \psi(r_n)}{\Psi(r_n)} \rightarrow \infty, \quad \text{as } n \rightarrow \infty.$$

And, using Young's equality (see [17, Eq. (2.7)]), we have

$$\begin{aligned} J_{C,1}(\mathbf{u}_n) &= \int_0^T \Phi\left(\frac{|\mathbf{u}_n|}{\Lambda}\right) dt - C \|\mathbf{u}_n\|_{L^\Phi} \\ &= \frac{1}{\Psi(r_n)} [\Phi(\psi(r_n)) - C \Lambda r_n \psi(r_n)] \\ &= \frac{1}{\Psi(r_n)} [r_n \psi(r_n) - \Psi(r_n) - C \Lambda r_n \psi(r_n)] \\ &= \frac{(1 - C\Lambda) r_n \psi(r_n)}{\Psi(r_n)} - 1. \end{aligned}$$

From (27) and the condition  $C\Lambda > 1$ , we obtain  $J_{C,1}(\mathbf{u}_n) \rightarrow -\infty$ , which contradicts the coercivity of  $J_{C,1}$ .  $\square$

In [1] we established coercivity of action integrals under different assumptions as follows.

**Theorem 3.3.** *Let  $\mathcal{L}$  be a lagrangian function satisfying (2), (3), (4), (22) and (23). We assume the following conditions:*

1. *There exist a non negative function  $b_1 \in L^1_1$  and a constant  $\mu > 0$  such that for any  $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^d$  and a.e.  $t \in [0, T]$*

$$|F(t, \mathbf{x}_2) - F(t, \mathbf{x}_1)| \leq b_1(t)(1 + |\mathbf{x}_2 - \mathbf{x}_1|^\mu). \quad (28)$$

*We suppose that  $\mu < \alpha_\Phi$ , with  $\alpha_\Phi$  as in Lemma 3.2, in the case that  $\Psi \in \Delta_2$ ; and, we suppose  $\mu = 1$  if  $\Psi$  is an arbitrary  $N$ -function.*

- 2.

$$\int_0^T F(t, \mathbf{x}) dt \rightarrow \infty \quad \text{as } |\mathbf{x}| \rightarrow \infty. \quad (29)$$

3.  $\Psi \in \Delta_2$  or, alternatively,  $\alpha_0^{-1} T \Phi^{-1}(1/T) \|b_1\|_{L^1} \Lambda < 1$ .

*Then the action integral  $I$  is coercive.*

Based on [21] we say that  $F$  satisfies the condition (A) if  $F(t, \mathbf{x})$  is a Carathéodory function,  $F$  verifies (23) and  $F$  is continuously differentiable with respect to  $\mathbf{x}$ . Moreover, the next inequality holds

$$|D_{\mathbf{x}} F(t, \mathbf{x})| \leq a(|\mathbf{x}|) b_0(t), \quad \text{for a.e. } t \in [0, T] \text{ and for every } \mathbf{x} \in \mathbb{R}^d. \quad (30)$$

The following result was proved in [21, p. 18].

**Lemma 3.4.** Suppose that  $F$  satisfies condition (A) and (29),  $F(t, \cdot)$  is differentiable and convex a.e.  $t \in [0, T]$ . Then, there exists  $\mathbf{x}_0 \in \mathbb{R}^d$  such that

$$\int_0^T D_{\mathbf{x}} F(t, \mathbf{x}_0) dt = 0. \quad (31)$$

**Theorem 3.5.** Let  $\mathcal{L}$  be as in Theorem 3.3 and let  $F$  be as in Lemma 3.4. Moreover, assume that  $\Psi \in \Delta_2$  or, alternatively  $\alpha_0^{-1} T \Phi^{-1}(1/T) a(|\mathbf{x}_0|) \|b_0\|_{L^1} \Lambda < 1$ , with  $a$  and  $b_0$  as in (23) and  $\mathbf{x}_0 \in \mathbb{R}^d$  any point satisfying (31). Then  $I$  is coercive.

## 4 New result

The symbol  $C$  with various subscripts will stand for a constant, not necessarily the same at each occurrence, which can depend (unless otherwise stated) on the constants previously displayed such as  $\|b_1\|_{L^1}, \|b_2\|_{L^1}, T, \mu, \mu'$ .

Now, we have another result about coercivity of  $I$  assuming some conditions on the nonlinearity  $\nabla F$ .

**Theorem 4.1.** Let  $\mathcal{L}$  be a lagrangian function satisfying (2), (3), (4), (22) and (23). We assume the following conditions:

1.  $\Psi \in \Delta_2$ .
2. There exist non negative functions  $b_1, b_2 \in L_1^1$  and a constant  $1 < \mu < \alpha_\Psi$  such that for any  $\mathbf{x} \in \mathbb{R}^d$  and a.e.  $t \in [0, T]$

$$|\nabla F(t, \mathbf{x})| \leq b_1(t) |\mathbf{x}|^{\mu-1} + b_2(t). \quad (32)$$

3. There exists a real positive number  $\sigma$  such that  $\sigma > (\mu - 1)\beta_\Psi$  and

$$|\mathbf{x}|^\sigma = o\left(\int_0^T F(t, \mathbf{x}) dt\right) \text{ as } |\mathbf{x}| \rightarrow \infty. \quad (33)$$

Then the action integral  $I$  is coercive.

*Proof.* By the decomposition  $u = \bar{u} + \tilde{u}$ , Mean Value Theorem, Cauchy-Schwarz inequality and (32), we have

$$\begin{aligned} \left| \int_0^T F(t, \mathbf{u}) - F(t, \bar{\mathbf{u}}) dt \right| &= \left| \int_0^T \int_0^1 \nabla F(t, \bar{\mathbf{u}} + s\tilde{\mathbf{u}}(t)) \cdot \tilde{\mathbf{u}}(t) ds dt \right| \\ &\leq \int_0^T \int_0^1 b_1(t) |\bar{\mathbf{u}} + s\tilde{\mathbf{u}}(t)|^{\mu-1} |\tilde{\mathbf{u}}(t)| ds dt + \int_0^T \int_0^1 b_2(t) |\tilde{\mathbf{u}}(t)| ds dt \\ &= I_1 + I_2. \end{aligned} \quad (34)$$

On the one hand, by Hölder's inequality and Sobolev's inequality, we estimate  $I_2$  as follows

$$I_2 \leq \|b_2\|_{L^1} \|\tilde{\mathbf{u}}\|_{L^\infty} \leq C \|\dot{\mathbf{u}}\|_{L^\Phi}. \quad (35)$$

On the other hand, as  $\bar{\mathbf{u}} \in \mathbb{R}$  and  $s \in [0, 1]$ , we have

$$|\bar{\mathbf{u}} + s\tilde{\mathbf{u}}(t)|^{\mu-1} \leq C(|\bar{\mathbf{u}}|^{\mu-1} + \|\tilde{\mathbf{u}}\|_{L^\infty}^{\mu-1}). \quad (36)$$

Now, inequality (36), Hölder's inequality and Sobolev's inequality imply that

$$\begin{aligned} I_1 &\leq C \left( |\bar{\mathbf{u}}|^{\mu-1} \int_0^T b_1(t) |\tilde{\mathbf{u}}(t)| dt + \|\tilde{\mathbf{u}}\|_{L^\infty}^{\mu-1} \int_0^T b_1(t) |\tilde{\mathbf{u}}(t)| dt \right) \\ &\leq C (|\bar{\mathbf{u}}|^{\mu-1} \|b_1\|_{L^1} \|\tilde{\mathbf{u}}\|_{L^\infty} + \|b_1\|_{L^1} \|\tilde{\mathbf{u}}\|_{L^\infty}^\mu) \\ &\leq C (|\bar{\mathbf{u}}|^{\mu-1} \|\tilde{\mathbf{u}}\|_{L^\infty} + \|\dot{\mathbf{u}}\|_{L^\Phi}^\mu). \end{aligned} \quad (37)$$

Let  $\mu'$  be a positive constant such that  $1 < \mu \leq \mu' < \alpha_\Phi$ . Next, using Young's inequality with conjugate exponents  $\mu'$  and  $\frac{\mu'}{\mu'-1}$  and Sobolev's inequality, we get

$$|\bar{\mathbf{u}}|^{\mu-1} \|\tilde{\mathbf{u}}\|_{L^\infty} \leq \frac{(\mu'-1)}{\mu'} |\bar{\mathbf{u}}|^\sigma + \frac{1}{\mu'} \|\tilde{\mathbf{u}}\|_{L^\infty}^{\mu'} \quad (38)$$

where  $\sigma = \frac{(\mu-1)\mu'}{\mu'-1}$  is a positive constant such that  $\sigma > (\mu-1)b_\Psi$ .

From (37), (38) and (35), we have

$$I_1 + I_2 \leq C(|\bar{\mathbf{u}}|^\sigma + \|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + \|\dot{\mathbf{u}}\|_{L^\Phi}^\mu + \|\dot{\mathbf{u}}\|_{L^\Phi}). \quad (39)$$

In the subsequent estimates, we use the decomposition  $u = \bar{u} + \tilde{u}$ , (22), (34), (39) and we get

$$\begin{aligned} I(\mathbf{u}) &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T F(t, \mathbf{u}) dt \\ &= \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T [F(t, \mathbf{u}) - F(t, \bar{\mathbf{u}})] dt + \int_0^T F(t, \bar{\mathbf{u}}) dt \\ &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - C(\|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + \|\dot{\mathbf{u}}\|_{L^\Phi}^\mu + \|\dot{\mathbf{u}}\|_{L^\Phi}) \\ &\quad + \int_0^T F(t, \bar{\mathbf{u}}) dt - C|\bar{\mathbf{u}}|^\sigma. \end{aligned} \quad (40)$$

As  $1 < \mu \leq \mu'$ , we have  $\|\dot{\mathbf{u}}\|_{L^\Phi} \leq \|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + 1$  and  $\|\dot{\mathbf{u}}\|_{L^\Phi}^\mu \leq \|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + 1$ , then

$$-C(\|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + \|\dot{\mathbf{u}}\|_{L^\Phi}^\mu + \|\dot{\mathbf{u}}\|_{L^\Phi}) \geq -C(3\|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + 2). \quad (41)$$

In this way, from (40) and (41)

$$\begin{aligned} I(\mathbf{u}) &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - C\|\dot{\mathbf{u}}\|_{L^\Phi}^{\mu'} + \int_0^T F(t, \bar{\mathbf{u}}) dt - K|\bar{\mathbf{u}}|^\sigma - C \\ &= \alpha_0 J_{C, \mu'}(\dot{\mathbf{u}}) + \gamma(\bar{\mathbf{u}}) - C. \end{aligned}$$

Let  $\mathbf{u}_n$  be a sequence in  $\mathcal{E}_d^\Phi(\lambda)$  with  $\|\mathbf{u}_n\|_{W^1 L^\Phi} \rightarrow \infty$  and we have to prove that  $I(\mathbf{u}_n) \rightarrow \infty$ .

On the contrary, suppose that for a subsequence, still denoted by  $\mathbf{u}_n$ ,  $I(\mathbf{u}_n)$  is upper bounded, that is, there exists  $M > 0$  such that  $|I(\mathbf{u}_n)| \leq M$ . As  $\|\mathbf{u}_n\|_{W^1 L^\Phi} \rightarrow \infty$ , from Lemma 2.2, we have  $\|\bar{\mathbf{u}}_n\| + \|\dot{\mathbf{u}}_n\|_{L^\Phi} \rightarrow \infty$ . Then, there exists subsequence of the subsequence  $\{\mathbf{u}_n\}$ , still denoted by  $\mathbf{u}_n$ , which is not bounded. Then,  $\bar{\mathbf{u}}_n \rightarrow \infty$  or  $\|\dot{\mathbf{u}}_n\|_{L^\Phi} \rightarrow \infty$ . Now, as the functionals  $J_{C,\mu'}(\dot{\mathbf{u}})$  and  $\gamma(\bar{\mathbf{u}})$  are coercive, then  $J_{C,\mu'}(\dot{\mathbf{u}}_n) \rightarrow \infty$  or  $\gamma(\bar{\mathbf{u}}_n) \rightarrow \infty$ . By (23), the functional  $\gamma(\bar{\mathbf{u}}_n)$  is lower bounded and  $J_{C,\mu'}(\dot{\mathbf{u}}_n)$  is also lower bounded on a bounded set because the modular  $\rho_\Phi(\frac{u}{\Lambda})$  is always bigger than zero. Therefore,  $I(\mathbf{u}_n) \rightarrow \infty$  as  $\|\mathbf{u}_n\|_{W^1 L^\Phi} \rightarrow \infty$  which contradicts the initial assumption on the behavior of  $I(\mathbf{u}_n)$ .  $\square$

REVISAR LA PRUEBA ANTERIOR Y MEJORAR LA ESCRITURA!!!!

## 5 Limit case $\mu = \alpha_\Phi$

In [] coercivity was obtained even in the limit case  $\mu = 1$  and  $\mu = p$  assuming additional conditions on ... This was possible because in  $L^p$  spaces, the norm and the modular coincides, that is,  $\|\cdot\|_p^p = O(\int_0^T |\cdot|^p dt)$ . In Orlicz spaces,  $\|\cdot\|_{L^\Phi}^\mu$  can be upper controlled by a modular provided that  $\mu < \alpha_\Phi$  for any  $N$ -function  $\Phi$ . But, the limit case does not hold for any  $\Phi$ , i.e. in general  $\|\cdot\|_{L^\Phi}^{\alpha_\Phi} = O(\int_0^T \Phi(|u|) dt)$  is false as can be seen as follows.

Let  $\Phi, \Psi \in \Delta_2$ , then the next inequality  $\Phi(tu) \geq t^{\alpha_\Phi} \Phi(u)$  for any  $u > 0$  and for any  $t \geq 1$  is false.

$$\text{In fact, let } \Phi(u) = \begin{cases} \frac{p-1}{p} u^p & u \leq e \\ \frac{u^p}{\log u} - \frac{e^p}{p} & u > e \end{cases}$$

**Theorem 5.1.** *If  $p \geq \frac{1+\sqrt{2}}{2}$ , then  $\Phi$  is an  $N$ -function.*

*Proof.* Resumir la prueba....  $\square$

**Theorem 5.2.** *There exists a constant  $C > 0$  such that*

$$\Phi(tu) \leq ct^p \Phi(u) \quad t \geq 1, u > 0. \quad (42)$$

*For every  $\varepsilon > 0$  there exists a constant  $C = C(\varepsilon, p)$  such that*

$$\Phi(tu) \geq Ct^{p-\varepsilon} \Phi(u) \quad t \geq 1, u > 0. \quad (43)$$

*Proof.* Resumir la prueba  $\square$

**Remark 5.3.** *The inequality*

$$\Phi(tu) \geq Ct^p \Phi(u)$$

*is false for every  $C$  because for every  $u \geq e$  we have*

$$\lim_{t \rightarrow \infty} \frac{\Phi(tu)}{t^p \Phi(u)} = 0$$

**Theorem 5.4.**  $\alpha_\Phi = \beta_\Phi = p$

*Proof.* Resumir la prueba. □

Now, we are able to see that

$$\rho_\Phi(u) = \int_0^T \Phi(|u|) dx \geq C \|u\|_{L^\Phi}^{\alpha_\Phi} = C \|u\|_{L^\Phi}^p$$

is false.

If we take  $u \equiv t > 0$ , then  $\|u\|_{L^\Phi}^p = C_1 t^p$  where  $C_1 = \|1\|_{L^\Phi}$  and  $\int_0^T \Phi(|u|) dx = C_2 \Phi(t)$  with  $C_2 = T$ . Then, if  $\rho_\Phi(u) \geq C \|u\|_{L^\Phi}^p$  were true, then  $\Phi(t) \geq C t^p$  were also true but this last inequality is false.

## Acknowledgments

The authors are partially supported by a UNRC grant number 18/C417. The first author is partially supported by a UNSL grant number 22/F223.

## References

- [1] S. Acinas, L. Buri, G. Giubergia, F. Mazzone, and E. Schwindt. “Some existence results on periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting”. In: (2015).
- [2] S. Acinas, G. Giubergia, F. Mazzone, and E. Schwindt. “On Estimates for the Period of Solutions of Equations Involving the  $\phi$ -Laplace Operator”. In: *Journal of Abstract Differential Equations and Applications* 5.1 (2014), pp. 21–34.
- [3] R.A. Adams and J.J.F. Fournier. *Sobolev Spaces*. Elsevier Science, 2003.
- [4] C. Bereanu and J. Mawhin. “Existence and multiplicity results for some nonlinear problems with singular  $\phi$ -Laplacian”. In: *Journal of Differential Equations* 243 (2007).
- [5] C. Bereanu and J. Mawhin. “Nonhomogeneous boundary value problems for some nonlinear equations with singular  $\phi$ -Laplacian”. In: *Journal of Mathematical Analysis and Applications* 352 (2009).
- [6] C. Bereanu and J. Mawhin. “Periodic solutions of nonlinear perturbations of  $\phi$ -Laplacians with possibly bounded  $\phi$ ”. In: *Nonlinear Analysis: Theory, Methods & Applications* 68 (2008).
- [7] A. Cianchi. “A fully anisotropic Sobolev inequality”. In: *Pacific J. Math* 196.2 (2000), pp. 283–295.
- [8] A. Cianchi. “Local boundedness of minimizers of anisotropic functionals”. In: *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*. Vol. 17. 2. Elsevier. 2000, pp. 147–168.
- [9] A. Cianchi. “Some results in the theory of Orlicz spaces and applications to variational problems”. In: *Nonlinear analysis, function spaces and applications* 6 (1999), pp. 50–92.
- [10] A. Cianchi and N. Fusco. “Gradient regularity for minimizers under general growth conditions”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 1999.507 (1999), pp. 15–36.
- [11] N. Clavero. *Optimal Sobolev embeddings and Function Spaces*. 2011. URL: <http://www.maia.ub.edu/~soria/sobolev1.pdf>.
- [12] D.E. Edmunds, R. Kerman, and L. Pick. “Optimal Sobolev imbeddings involving rearrangement-invariant quasinorms”. In: *Journal of Functional Analysis* 170.2 (2000), pp. 307–355.
- [13] I. Ekeland and R. Témam. *Convex Analysis and Variational Problems*. Society for Industrial and Applied Mathematics, 1999.

- [14] M. García-Huidobro, V.K. Le, R. Manásevich, and K. Schmitt. “On principal eigenvalues for quasilinear elliptic differential operators: an Orlicz-Sobolev space setting”. In: *Nonlinear Differential Equations and Applications NoDEA* 6 (1999).
- [15] H. Hudzik and L. Maligranda. “Amemiya norm equals Orlicz norm in general”. In: *Indagationes Mathematicae* 11.4 (2000), pp. 573–585.
- [16] R. Kerman and L. Pick. “Optimal Sobolev imbeddings”. In: *Forum Mathematicum*. Vol. 18. 2006, pp. 535–570.
- [17] M. A. Krasnosel’skiĭ and Ja. B. Rutickiĭ. *Convex functions and Orlicz spaces*. Groningen: P. Noordhoff Ltd., 1961.
- [18] L. Maligranda. *Orlicz spaces and interpolation*. Vol. 5. Seminários de Matemática [Seminars in Mathematics]. Universidade Estadual de Campinas, Departamento de Matemática, Campinas, 1989.
- [19] R. Manásevich and J. Mawhin. “Boundary value problems for nonlinear perturbations of vector  $p$ -Laplacian-like operators”. In: *J. Korean Math. Soc* 37.5 (2000), pp. 665–685.
- [20] R. Manásevich and J. Mawhin. “Periodic Solutions for Nonlinear Systems with  $p$ -Laplacian-Like Operators”. In: *Journal of Differential Equations* 145.2 (1998), pp. 367–393.
- [21] J. Mawhin and M. Willem. *Critical point theory and Hamiltonian systems*. Vol. 74. Applied Mathematical Sciences. Springer-Verlag, New York, 1989.
- [22] M. M. Rao and Z. D. Ren. *Theory of Orlicz spaces*. M. Dekker, 1991.
- [23] M.M. Rao and Z.D. Ren. *Applications Of Orlicz Spaces*. Monographs and textbooks in pure and applied mathematics. Taylor & Francis, 2002.
- [24] R.T. Rockafellar and R.J.B. Wets. *Variational Analysis*. Springer, 2009.
- [25] G. Schappacher. “A Notion of Orlicz Spaces for Vector Valued Functions”. English. In: *Applications of Mathematics* 50.4 (2005), pp. 355–386.
- [26] M. Struwe. *Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, 2008.
- [27] C.-L. Tang. “Periodic solutions for nonautonomous second order systems with sublinear nonlinearity”. In: *Proceedings of the American Mathematical Society* (1998), pp. 3263–3270.
- [28] C.-L. Tang. “Periodic solutions of non-autonomous second-order systems with  $\gamma$ -quasisubadditive potential”. In: *Journal of Mathematical Analysis and Applications* 189.3 (1995), pp. 671–675.
- [29] C.-L. Tang and X.-P. Wu. “Periodic solutions for second order systems with not uniformly coercive potential”. In: *Journal of Mathematical Analysis and Applications* 259.2 (2001), pp. 386–397.
- [30] X. Tang and X. Zhang. “Periodic solutions for second-order Hamiltonian systems with a  $p$ -Laplacian”. In: *Annales UMCS, Mathematica* 64.1 (2010), pp. 93–113.



## References

---

- [31] Y. Tian and W. Ge. “Periodic solutions of non-autonomous second-order systems with a p-Laplacian”. In: *Nonlinear Analysis: Theory, Methods & Applications* 66.1 (2007), pp. 192–203.
- [32] X.-P. Wu and C.-L. Tang. “Periodic solutions of a class of non-autonomous second-order systems”. In: *Journal of Mathematical Analysis and Applications* 236.2 (1999), pp. 227–235.
- [33] B. Xu and C.-L. Tang. “Some existence results on periodic solutions of ordinary p-Laplacian systems”. In: *Journal of mathematical analysis and applications* 333.2 (2007), pp. 1228–1236.
- [34] Y.-W. Ye and C.-L. Tang. “Periodic solutions for some nonautonomous second order Hamiltonian systems”. In: *Journal of Mathematical Analysis and Applications* 344.1 (2008), pp. 462–471.
- [35] F. Zhao and X. Wu. “Periodic solutions for a class of non-autonomous second order systems”. In: *Journal of mathematical analysis and applications* 296.2 (2004), pp. 422–434.