

Periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting by the dual least action principle

Sonia Acinas *

Instituto de Matemática Aplicada San Luis (IMASL)
Universidad Nacional de San Luis and CONICET
Ejército de los Andes 950, (D5700HDW) San Luis, Argentina
Universidad Nacional de La Pampa
(L6300CLB) Santa Rosa, La Pampa, Argentina
sonia.acinas@gmail.com

Fernando D. Mazzone †

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales
Universidad Nacional de Río Cuarto
(5800) Río Cuarto, Córdoba, Argentina,
fmazzone@exa.unrc.edu.ar

Abstract

1 Introduction

This paper deals with system of equations of the type:

$$\begin{cases} \frac{d}{dt} D_y \mathcal{L}(t, u(t), u'(t)) = D_x \mathcal{L}(t, u(t), u'(t)) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = u'(0) - u'(T) = 0, \end{cases} \quad (1)$$

where $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, $d \geq 1$, is called the *Lagrange function* or *lagrangian* and the unknown function $u : [0, T] \rightarrow \mathbb{R}^d$ is absolutely continuous. In other words, we are interested in finding *periodic weak solutions* of *Euler-Lagrange system of ordinary*

*SECyT-UNRC, UNSL and CONICET

†SECyT-UNRC and CONICET

2010 AMS Subject Classification. Primary: . Secondary: .

Keywords and phrases. .

equations. This topic was deeply addressed for the *Lagrange function*

$$\mathcal{L}_{p,F}(t, x, y) = \frac{|y|^p}{p} + F(t, x), \quad (2)$$

for $1 < p < \infty$. For example, the classic book [9] deals mainly with problem (1), for the lagrangian $\mathcal{L}_{2,F}$, through various methods: direct, dual action, minimax, etc. The results in [9] were extended and improved in several articles, see [19, 17, 22, 18, 23] to cite some examples. Lagrange functions (2) for arbitrary $1 < p < \infty$ were considered in [21, 20] and in this case (1) is reduced to the p -laplacian system

$$\begin{cases} \frac{d}{dt}(u'(t)|u'|^{p-2}) = \nabla F(t, u(t)) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = u'(0) - u'(T) = 0. \end{cases} \quad (3)$$

In this context, it is customary to call F a *potential function*, and it is assumed that $F(t, x)$ is differentiable with respect to x for a.e. $t \in [0, T]$ and the following conditions are verified:

(C) F and its gradient ∇F , with respect to $x \in \mathbb{R}^d$, are Carathéodory functions, i.e. they are measurable functions with respect to $t \in [0, T]$, for every $x \in \mathbb{R}^d$, and they are continuous functions with respect to $x \in \mathbb{R}^d$ for a.e. $t \in [0, T]$.

(A) For a.e. $t \in [0, T]$, it holds that

$$|F(t, x)| + |\nabla F(t, x)| \leq a(|x|)b(t). \quad (4)$$

In this inequality we assume that the function $a : [0, +\infty) \rightarrow [0, +\infty)$ is continuous and nondecreasing and $0 \leq b \in L^1([0, T], \mathbb{R})$.

In [1] it was treated the case of a lagrangian \mathcal{L} which is lower bounded by a Lagrange function

$$\mathcal{L}_{\Phi,F}(t, x, y) = \Phi(|y|) + F(t, x), \quad (5)$$

where Φ is an N -function (see section 2 for the definition of this concept). In the paper [1] it was assumed a condition of *bounded oscillation* on F (see xxxxx below). In this paper we apply the dual method ([9, Ch. 3]) to obtain solutions of (1).

2 Preliminaries

In this section, we give a short introduction to Orlicz and Orlicz-Sobolev spaces of vector valued functions and other brief introduction to superposition operators between these spaces. References for these topics are [15, 16, 3] and [13, 10, 12, 11].

Hereafter we denote by \mathbb{R}^+ the set of all non negative real numbers. A function $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is called an N -function if Φ is convex and it also satisfies that

$$\lim_{t \rightarrow +\infty} \frac{\Phi(t)}{t} = +\infty \quad \text{and} \quad \lim_{t \rightarrow 0} \frac{\Phi(t)}{t} = 0.$$

In addition, in this paper for the sake of simplicity we assume that Φ is differentiable and we call φ the derivative of Φ . On these assumptions, $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a homeomorphism whose inverse will be denoted by ψ . We denote by Ψ the primitive of ψ that satisfies $\Psi(0) = 0$. Then, Ψ is an N -function which is called the *complementary function* of Φ .

We recall that an N -function $\Phi(u)$ has *principal part* $f(u)$ if $\Phi(u) = f(u)$ for large values of the argument (see [6, p. 16] and [6, Sec. 7] for properties of principal part).

There exist several orders and equivalence relations between N -functions (see [14, Sec. 2.2]). Following [14, Def. 1, pp. 15-16] we say that the N -function Φ_2 is *stronger* than the N -function Φ_1 , in symbols $\Phi_1 < \Phi_2$, if there exist $a > 0$ and $x_0 \geq 0$ such that

$$\Phi_1(x) \leq \Phi_2(ax), \quad x \geq x_0. \quad (6)$$

The N -functions Φ_1 and Φ_2 are *equivalent* ($\Phi_1 \sim \Phi_2$) when $\Phi_1 < \Phi_2$ and $\Phi_2 < \Phi_1$. We say that Φ_2 is *essentially stronger* than Φ_1 ($\Phi_1 \ll \Phi_2$) if and only if for every $a > 0$ there exists $x_0 = x_0(a) \geq 0$ such that (6) holds. Finally, we say that Φ_2 is *completely stronger* than Φ_1 ($\Phi_1 \prec \Phi_2$) if and only if for every $a > 0$ there exist $K = K(a) > 0$ and $x_0 = x_0(a) \geq 0$ such that

$$\Phi_1(x) \leq K\Phi_2(ax), \quad x \geq x_0. \quad (7)$$

We also say that a non decreasing function $\eta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfies the Δ_2^∞ -condition, denoted by $\eta \in \Delta_2^\infty$, if there exist constants $K > 0$ and $x_0 \geq 0$ such that

$$\eta(2x) \leq K\eta(x), \quad (8)$$

for every $x \geq x_0$. We note that $\eta \in \Delta_2^\infty$ if and only if $\eta \prec \eta$. If $x_0 = 0$, the function $\eta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is said to satisfy the Δ_2 -condition ($\eta \in \Delta_2$). If there exists $x_0 > 0$ such that inequality (8) holds for $x \leq x_0$, we will say that Φ satisfies the Δ_2^0 -condition ($\Phi \in \Delta_2^0$).

We denote by α_η and β_η the so called *Matuszewska-Orlicz indices* of the function η , which are defined next. Given an increasing, unbounded, continuous function $\eta : [0, +\infty) \rightarrow [0, +\infty)$ such that $\eta(0) = 0$, we define

$$\alpha_\eta := \lim_{t \rightarrow 0^+} \frac{\log \left(\sup_{u>0} \frac{\eta(tu)}{\eta(u)} \right)}{\log(t)}, \quad \beta_\eta := \lim_{t \rightarrow +\infty} \frac{\log \left(\sup_{u>0} \frac{\eta(tu)}{\eta(u)} \right)}{\log(t)}. \quad (9)$$

It is known that the previous limits exist and $0 \leq \alpha_\eta \leq \beta_\eta \leq +\infty$ (see [8, p. 84]). The relation $\beta_\eta < +\infty$ holds true if and only if $\eta \in \Delta_2$ ([8, Thm. 11.7]). If (Φ, Ψ) is a complementary pair of N -functions then

$$\frac{1}{\alpha_\Phi} + \frac{1}{\beta_\Psi} = 1, \quad (10)$$

(see [8, Cor. 11.6]). Therefore $1 \leq \alpha_\Phi \leq \beta_\Phi \leq \infty$.

If η is an increasing function that satisfies the Δ_2 -condition, then η is controlled by above and below by power functions ([4, Sec. 1], [?, Eq. (2.3)-(2.4)] and [8, Thm.

11.13]). More concretely, for every $\epsilon > 0$ there exists a constant $K = K(\eta, \epsilon)$ such that, for every $t, u \geq 0$,

$$K^{-1} \min \{t^{\beta_\eta + \epsilon}, t^{\alpha_\eta - \epsilon}\} \eta(u) \leq \eta(tu) \leq K \max \{t^{\beta_\eta + \epsilon}, t^{\alpha_\eta - \epsilon}\} \eta(u). \quad (11)$$

Let d be a positive integer. We denote by $\mathcal{M} := \mathcal{M}([0, T], \mathbb{R}^d)$ the set of all measurable functions defined on $[0, T]$ with values on \mathbb{R}^d and we write $u = (u_1, \dots, u_d)$ for $u \in \mathcal{M}$. For the set of functions \mathcal{M} , as for other similar sets, we will omit the reference to codomain \mathbb{R}^d when $d = 1$.

Given an N -function Φ we define the *modular function* $\rho_\Phi : \mathcal{M} \rightarrow \mathbb{R}^+ \cup \{+\infty\}$ by

$$\rho_\Phi(u) := \int_0^T \Phi(|u|) dt.$$

Here $|\cdot|$ is the euclidean norm of \mathbb{R}^d . Now, we introduce the *Orlicz class* $C^\Phi = C^\Phi([0, T], \mathbb{R}^d)$ by setting

$$C^\Phi := \{u \in \mathcal{M} | \rho_\Phi(u) < \infty\}. \quad (12)$$

The *Orlicz space* $L^\Phi = L^\Phi([0, T], \mathbb{R}^d)$ is the linear hull of C^Φ ; equivalently,

$$L^\Phi := \{u \in \mathcal{M} | \exists \lambda > 0 : \rho_\Phi(\lambda u) < \infty\}. \quad (13)$$

The Orlicz space L^Φ equipped with the *Orlicz norm*

$$\|u\|_{L^\Phi} := \sup \left\{ \int_0^T u \cdot v dt | \rho_\Psi(v) \leq 1 \right\},$$

is a Banach space. By $u \cdot v$ we denote the usual dot product in \mathbb{R}^d between u and v .

The following inequality holds for any $u \in L^\Phi$

$$\|u\|_{L^\Phi} \leq \frac{1}{k} \{1 + \rho_\Phi(ku)\}, \quad \text{for every } k > 0. \quad (14)$$

In fact, $\|u\|_{L^\Phi}$ is the infimum for $k > 0$ of the right hand side in above expression (see [6, Thm. 10.5] and [5]).

The subspace $E^\Phi = E^\Phi([0, T], \mathbb{R}^d)$ is defined as the closure in L^Φ of the subspace $L^\infty([0, T], \mathbb{R}^d)$ of all \mathbb{R}^d -valued essentially bounded functions. It is shown that E^Φ is the only one maximal subspace contained in the Orlicz class C^Φ , i.e. $u \in E^\Phi$ if and only if $\rho_\Phi(\lambda u) < \infty$ for any $\lambda > 0$. The equality $L^\Phi = E^\Phi$ is true if and only if $\Phi \in \Delta_2^\infty$.

A generalized version of *Hölder's inequality* holds in Orlicz spaces (see [6, Thm. 9.3]). Namely, if $u \in L^\Phi$ and $v \in L^\Psi$ then $u \cdot v \in L^1$ and

$$\int_0^T v \cdot u dt \leq \|u\|_{L^\Phi} \|v\|_{L^\Psi}. \quad (15)$$

Like in [6], we will consider the subset $\Pi(E_d^\Phi, r)$ of L_d^Φ given by

$$\Pi(E_d^\Phi, r) := \{u \in L_d^\Phi | d(u, E_d^\Phi) < r\}.$$

This set is related to the Orlicz class C_d^Φ by means of inclusions, namely,

$$\Pi(E_d^\Phi, r) \subset rC_d^\Phi \subset \overline{\Pi(E_d^\Phi, r)} \quad (16)$$

for any positive r . If $\Phi \in \Delta_2$, then the sets L_d^Φ , E_d^Φ , $\Pi(E_d^\Phi, r)$ and C_d^Φ are equal.

Let $\mathcal{E}_d^{\Phi_i}(\lambda) := W^1 L_d^{\Phi_i} \cap \{u | u \in \Pi(E_d^{\Phi_i}, \lambda)\}$.

If X and Y are Banach spaces such that $Y \subset X^*$, we denote by $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{R}$ the bilinear pairing map given by $\langle x^*, x \rangle = x^*(x)$. Hölder's inequality shows that $L^\Psi \subset [L^\Phi]^*$, where the pairing $\langle v, u \rangle$ is defined by

$$\langle v, u \rangle = \int_0^T v \cdot u \, dt, \quad (17)$$

with $u \in L^\Phi$ and $v \in L^\Psi$. Unless $\Phi \in \Delta_2^\infty$, the relation $L^\Psi = [L^\Phi]^*$ will not be satisfied. In general, it is true that $[E^\Phi]^* = L^\Psi$.

We define the *Sobolev-Orlicz space* $W^1 L^\Phi$ (see [2]) by

$$W^1 L^\Phi := \{u | u \text{ is absolutely continuous on } [0, T] \text{ and } u' \in L^\Phi\}.$$

$W^1 L^\Phi$ is a Banach space when equipped with the norm

$$\|u\|_{W^1 L^\Phi} = \|u\|_{L^\Phi} + \|u'\|_{L^\Phi}. \quad (18)$$

And, we introduce the following subspaces of $W^1 L^\Phi$

$$\begin{aligned} W^1 E^\Phi &= \{u \in W^1 L^\Phi | u' \in E^\Phi\}, \\ W^1 E_T^\Phi &= \{u \in W^1 E^\Phi | u(0) = u(T)\}. \end{aligned} \quad (19)$$

We will use repeatedly the decomposition $u = \bar{u} + \tilde{u}$ for a function $u \in L^1([0, T])$ where $\bar{u} = \frac{1}{T} \int_0^T u(t) \, dt$ and $\tilde{u} = u - \bar{u}$.

As usual, if $(X, \|\cdot\|_X)$ is a Banach space and $(Y, \|\cdot\|_Y)$ is a subspace of X , we write $Y \hookrightarrow X$ and we say that Y is *embedded* in X when the restricted identity map $i_Y : Y \rightarrow X$ is bounded. That is, there exists $C > 0$ such that for any $y \in Y$ we have $\|y\|_X \leq C\|y\|_Y$. With this notation, Hölder's inequality states that $L^\Psi \hookrightarrow [L^\Phi]^*$; and, it is easy to see that for every N -function Φ we have that $L^\infty \hookrightarrow L^\Phi \hookrightarrow L^1$.

Recall that a function $w : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is called a *modulus of continuity* if w is a continuous increasing function which satisfies $w(0) = 0$. For example, it can be easily shown that $w(s) = s\Phi^{-1}(1/s)$ is a modulus of continuity for every N -function Φ . It is said that $u : [0, T] \rightarrow \mathbb{R}^d$ has modulus of continuity w when there exists a constant $C > 0$ such that

$$|u(t) - u(s)| \leq Cw(|t - s|). \quad (20)$$

We denote by $C^w([0, T], \mathbb{R}^d)$ the space of w -Hölder continuous functions that satisfy (20) for some $C > 0$. This is a Banach space with norm

$$\|u\|_{C^w([0, T], \mathbb{R}^d)} := \|u\|_{L^\infty} + \sup_{t \neq s} \frac{|u(t) - u(s)|}{w(|t - s|)}.$$

The following simple embedding lemma, whose proof can be found in [1], will be used systematically.

Lemma 2.1. *Let $w(s) := s\Phi^{-1}(1/s)$. Then, the following statements hold:*

1. $W^1 L^\Phi \hookrightarrow C^w([0, T], \mathbb{R}^d)$ and for every $u \in W^1 L^\Phi$

$$|u(t) - u(s)| \leq \|u'\|_{L^\Phi} w(|t - s|) \quad (\text{Morrey's inequality}), \quad (21)$$

$$\|u\|_{L^\infty} \leq \Phi^{-1}\left(\frac{1}{T}\right) \max\{1, T\} \|u\|_{W^1 L^\Phi} \quad (\text{Sobolev's inequality}). \quad (22)$$

2. For every $u \in W^1 L^\Phi$ we have $\tilde{u} \in L_d^\infty$ and

$$\|\tilde{u}\|_{L^\infty} \leq T\Phi^{-1}\left(\frac{1}{T}\right) \|u'\|_{L^\Phi} \quad (\text{Sobolev-Wirtinger's inequality}). \quad (23)$$

3 Once upon a time...

Vamos escribiendo lo que queremos...(de acuerdo a mis apuntes y sin ver las hojitas de la semana pasada)

For $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ we denote by \mathfrak{f} the Nemytskii (o superposition) operator defined for functions $u : [0, T] \rightarrow \mathbb{R}^d$ by

$$\mathfrak{f}u(t) = f(t, u(t))$$

Referencias y alguna propiedad interesante medibles en medibles? [7, 6]

Theorem 3.1. *Let $\Phi_1, \Phi_2, \dots, \Phi_n$ be N -functions. Assume that M is another N -functions that satisfy the Δ_2 -condition. We write $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ with $x_i \in \mathbb{R}^d$, $y_i \in \mathbb{R}^d$. Let $f(t, x_1, \dots, x_n, y_1, \dots, y_n)$ be a function Chatratheodory? with $f : [0, T] \times (\mathbb{R}^d)^n \times (\mathbb{R}^d)^n \rightarrow \mathbb{R}^{d'}$.*

Suppose that $a : (\mathbb{R}^d)^n \rightarrow [0, +\infty)$ is a bounded function on bounded sets and $b \in L^M([0, T])$, for a.e. $t \in [0, T]$ such that

$$|f| \leq a(x)[b(t) + \sum_{i=1}^n M^{-1}(\Phi_i(|y_i|))], \quad (24)$$

then

$$\mathfrak{f} : \left(\prod_{i=1}^n L^\infty([0, T], \mathbb{R}^d) \right) \times \left(\prod_{i=1}^n \Pi(E^{\Phi_i}([0, T], \mathbb{R}^d), \lambda = 1) \right) \rightarrow L^M.$$

Proof. If $(u, v) \in \left(\prod_{i=1}^n L^\infty([0, T], \mathbb{R}^d) \right) \times \left(\prod_{i=1}^n \Pi(E^{\Phi_i}, \lambda = 1) \right)$. By [6, Thm. 17.6] (y otras cosas), we get

$$|\mathfrak{f}u(t)| = |f(t, u(t), v(t))| \leq M_a[b_j(t) + \sum_{i=1}^n M_j^{-1}(\Phi_i(|v_i(t)|))] \in L_1^{M_j}.$$

□

We define the space X by $X = \{v = (v_1, v_2) : v_1 \in W^1 L_T^{\Phi_1}, v_2 \in W^1 L_T^{\Phi_2}\}$ and $X^* = \{v = (v_1, v_2) : v_1 \in (W^1 L_T^{\Phi_1})^*, v_2 \in (W^1 L_T^{\Phi_2})^*\}$ where $(W^1 L_T^{\Phi_i})^*$ stands for the conjugate space of $W^1 L_T^{\Phi_i}$ for $i = 1, 2$.

Corollary 3.2. *We will consider the Lagrange function $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$, $(t, x_1, x_2, y_1, y_2) \rightarrow \mathcal{L}(t, x_1, x_2, y_1, y_2)$ which is measurable in t for each $(x_1, x_2, y_1, y_2) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ and continuously differentiable in (x_1, x_2, y_1, y_2) for almost every $t \in [0, T]$.*

Let $x = (x_1, x_2)$, $y = (y_1, y_2)$ with $x_i \in \mathbb{R}^d$ and $y_i \in \mathbb{R}^d$ and let

$$I(x) = \int_0^T \mathcal{L}(t, x, y) dt \quad (25)$$

If there exist $a \in C(\mathbb{R}^+, \mathbb{R}^+)$, $i = 1, 2$, $b \in L_1^1([0, T])$, $j = 1, \dots, d'$ for a.e. $t \in [0, T]$ and every $(x_1, x_2, y_1, y_2) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ satisfying the structure conditions

$$|\mathcal{L}(t, x, y)| + \sum_{i=1}^2 |D_{x_i} \mathcal{L}(t, x, y)| \leq a(|x|)(b(t) + \Phi_1(|y_1|) + \Phi_2(|y_2|)), \quad (26)$$

$$|D_{y_i} \mathcal{L}(t, x, y)| \leq a(|x|)(c_i(t) + \sum_{j=1}^n \Psi_i^{-1}(\Phi_j(|y_j|))) \quad i = 1, 2. \quad (27)$$

The nonlinear operator $(x_1, x_2) \mapsto D_x \mathcal{L}(t, x_1, y_1, y_2)$ is continuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda) \times \dots \times \mathcal{E}_d^{\Phi_n}(\lambda)$ with the strong topology into $L^1([0, T])$ with the strong topology on both sets.

The nonlinear operator $(x_1, x_2) \mapsto D_y \mathcal{L}(t, x_1, y_1, y_2)$ is continuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda) \times \dots \times \mathcal{E}_d^{\Phi_n}(\lambda)$ with the strong topology into X with the weak topology.*

The function I is Gâteaux differentiable on $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ and its derivative I' is demicontinuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ into X^ . Moreover, I' is given by the following expression*

$$\begin{aligned} \langle I'(x), w \rangle = \int_0^T & [(D_{x_1} \mathcal{L}(t, x_1(t), x_2(t), y_1(t), y_2(t)), w_1(t)) + \\ & (D_{x_2} \mathcal{L}(t, x_1(t), x_2(t), y_1(t), y_2(t)), w_2(t)) + \\ & (D_{y_1} \mathcal{L}(t, x_1(t), x_2(t), y_1(t), y_2(t)), w'_1(t)) + \\ & (D_{y_2} \mathcal{L}(t, x_1(t), x_2(t), y_1(t), y_2(t)), w'_2(t))] dt \end{aligned} \quad (28)$$

If $\Psi \in \Delta_2$ then I' is continuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ into X^ when both spaces are equipped with the strong topology.*

We denote by $\mathfrak{A}(a, b, c, \lambda, f, \Phi)$ the set of all Lagrange functions satisfying (??), (??) and (??).

Proof. OJO!!!! Es algo que teníamos del trabajo anterior!!! con algunas adaptaciones a 2 variables sin controlar y a lo bruto!!!!

Let $u \in \mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$.

Step 1. The non linear operator $(x_1, x_2) \mapsto (D_{x_1} \mathcal{L}(t, x_1, x_2, y_1, y_2), D_{x_2} \mathcal{L}(t, x_1, x_2, y_1, y_2))$ is continuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ into $L_d^1([0, T]) \times L_d^1([0, T])$ with the strong topology on both sets.

If $u \in \mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$, from (??) and (??), we obtain Let $\{x_n = (x_{1n}, x_{2n})\}_{n \in \mathbb{N}}$ be a sequence of functions in $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ and let $x = (x_1, x_2) \in \mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ such that $x_n \rightarrow x$ in X . From $x_{in} \rightarrow x_i$ in L^{Φ_i} , there exists a subsequence x_{in_k} such that $x_{in_k} \rightarrow x_i$ a.e.; and, as $x_{in} \rightarrow x_i \in \mathcal{E}_d^{\Phi}(\lambda)$, by Lemma ??, there exist a subsequence of x_{in_k} (again denoted x_{in_k}) and a function $h_i \in \Pi(E_1^{\Phi}, \lambda)$ such that $x_{in_k} \rightarrow u_i$ a.e. and $|x_{in_k}| \leq h_i$ a.e. Since $x_{in_k}, k = 1, 2, \dots$, is a strong convergent sequence in $W^1 L_d^{\Phi_i}$, it is a bounded sequence in $W^1 L_d^{\Phi_i}$. According to Lemma 2.1 and Corollary ??, there exist $M_i > 0$ such that $\|a(x_{in_k})\|_{L^\infty} \leq M_i, k = 1, 2, \dots$. From the previous facts and (??), we get

$$|D_{x_i} \mathcal{L}(\cdot, x_{1n_k}, x_{2n_k}, y_{1n_k}, y_{2n_k})| \leq M_i(b + \Phi_i(|h_i|)) \in L_1^1 \quad i = 1, 2.$$

On the other hand, by the continuous differentiability of \mathcal{L} , we have

$$D_{x_i} \mathcal{L}(t, x_{in_k}(t), y_{in_k}(t)) \rightarrow D_{x_i} \mathcal{L}(t, x_i(t), y_i(t)) \quad \text{for a.e. } t \in [0, T].$$

Applying the Dominated Convergence Theorem we conclude the proof of step 1.

Step 2. The non linear operator $(x_1, x_2) \mapsto (D_{y_1} \mathcal{L}(t, x_1, y_1), D_{y_2} \mathcal{L}(t, x_2, y_2))$ is continuous from $\mathcal{E}_d^{\Phi_1}(\lambda) \times \mathcal{E}_d^{\Phi_2}(\lambda)$ with the strong topology into X with the weak topology.*

Note that (??), (??) and the imbeddings $W^1 L_d^{\Phi} \hookrightarrow L_d^\infty$ and $L_d^\Psi \hookrightarrow [L_d^{\Phi}]^*$ imply that the second member of (28) defines an element in $[W^1 L_d^{\Phi}]^*$.

Let $(x_{1n}, x_{2n}) \in \mathcal{E}_d^{\Phi}(\lambda)$ such that $(x_{1n}, x_{2n}) \rightarrow (x_1, x_2)$ in the norm of X . We must prove that $D_{y_i} \mathcal{L}(\cdot, x_{1n}, x_{2n}) \xrightarrow{w^*} D_{y_i} \mathcal{L}(\cdot, x_1, x_2, y_1, y_2)$ para $i = 1, 2$. On the contrary, there exist $v = (v_1, v_2) \in L^{\Phi_1} \times L^{\Phi_2}$, $\epsilon > 0$ and a subsequence of $\{x_n\}$ (denoted $\{x_n\}$ for simplicity) such that

$$|\langle D_{y_i} \mathcal{L}(\cdot, x_{1n}, x_{2n}, y_{1n}, y_{2n}), v \rangle - \langle D_{y_i} \mathcal{L}(\cdot, x_1, x_2, y_1, y_2), v \rangle| \geq \epsilon. \quad (29)$$

We have $x_n \rightarrow x$ in X and $y_n \rightarrow y$ in X . By Lemma ??, there exist a subsequence x_{n_k} and a function $h \in \Pi(E_1^{\Phi_1}, \lambda) \times \Pi(E_1^{\Phi_2}, \lambda)$ such that $x_{n_k} \rightarrow x$ a.e., $y_{n_k} \rightarrow y$ a.e. and $|y_{n_k}| \leq h$ a.e. As in the previous step, since x_n is a convergent sequence, the Corollary ?? implies that $a(|y_n(t)|)$ is uniformly bounded by a certain constant $M > 0$. Therefore, with x_{n_k} instead of x , inequality (??) becomes

$$|D_{y_i} \mathcal{L}(\cdot, x_{n_k}, y_{n_k})| \leq M_i(c_i + \varphi_i(h_i) + \Psi_i^{-1}(\Phi_j(|y_j|))) \in L_1^{\Psi_i}. \quad (30)$$

Consequently, as $v \in L_d^{\Phi}$ and employing Hölder's inequality, we obtain that

$$\sup_k |D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k}) \cdot v| \in L_1^1.$$

Finally, from the Lebesgue Dominated Convergence Theorem, we deduce

$$\int_0^T D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k}) \cdot v \, dt \rightarrow \int_0^T D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot v \, dt \quad (31)$$

which contradicts the inequality (29). This completes the proof of step 2.

Step 3. We will prove (28). The proof follows similar lines as [9, Thm. 1.4]. For $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$ and $\mathbf{0} \neq \mathbf{v} \in W^1 L_d^\Phi$, we define the function

$$H(s, t) := \mathcal{L}(t, \mathbf{u}(t) + s\mathbf{v}(t), \dot{\mathbf{u}}(t) + s\dot{\mathbf{v}}(t)).$$

From [6, Lemma 10.1] (or [15, Thm. 5.5]) we obtain that if $|\mathbf{u}| \leq |\mathbf{v}|$ then $d(\mathbf{u}, E_d^\Phi) \leq d(\mathbf{v}, E_d^\Phi)$. Therefore, for $|s| \leq s_0 := (\lambda - d(\dot{\mathbf{u}}, E_d^\Phi)) / \|\mathbf{v}\|_{W^1 L^\Phi}$ we have

$$d(\dot{\mathbf{u}} + s\dot{\mathbf{v}}, E_d^\Phi) \leq d(|\dot{\mathbf{u}}| + s|\dot{\mathbf{v}}|, E_1^\Phi) \leq d(|\dot{\mathbf{u}}|, E_1^\Phi) + s\|\dot{\mathbf{v}}\|_{L^\Phi} < \lambda.$$

Thus $\dot{\mathbf{u}} + s\dot{\mathbf{v}} \in \Pi(E_d^\Phi, \lambda)$ and $|\dot{\mathbf{u}}| + s|\dot{\mathbf{v}}| \in \Pi(E_1^\Phi, \lambda)$. These facts imply, in virtue of Theorem ?? item ??, that $I(\mathbf{u} + s\mathbf{v})$ is well defined and finite for $|s| \leq s_0$. And, using Corollary ??, we also see that

$$\|a(|\mathbf{u} + s\mathbf{v}|)\|_{L^\infty} \leq A(\|\mathbf{u} + s\mathbf{v}\|_{W^1 L^\Phi}) \leq A(\|\mathbf{u}\|_{W^1 L^\Phi} + s_0\|\mathbf{v}\|_{W^1 L^\Phi}) =: M$$

Now, applying Chain Rule, (??), (??) the monotonicity of φ and Φ , the fact that $\mathbf{v} \in L_d^\infty$ and $\dot{\mathbf{v}} \in L_d^\Phi$ and Hölder's inequality, we get

$$\begin{aligned} |D_s H(s, t)| &= |D_x \mathcal{L}(t, \mathbf{u} + s\mathbf{v}, \dot{\mathbf{u}} + s\dot{\mathbf{v}}) \cdot \mathbf{v} + D_y \mathcal{L}(t, \mathbf{u} + s\mathbf{v}, \dot{\mathbf{u}} + s\dot{\mathbf{v}}) \cdot \dot{\mathbf{v}}| \\ &\leq M \left[\left(b(t) + \Phi \left(\frac{|\dot{\mathbf{u}}| + s_0|\dot{\mathbf{v}}|}{\lambda} + f(t) \right) \right) |\mathbf{v}| \right. \\ &\quad \left. + \left(c(t) + \varphi \left(\frac{|\dot{\mathbf{u}}| + s_0|\dot{\mathbf{v}}|}{\lambda} + f(t) \right) \right) |\dot{\mathbf{v}}| \right] \in L_1^1. \end{aligned} \quad (32)$$

Consequently, I has a directional derivative and

$$\langle I'(\mathbf{u}), \mathbf{v} \rangle = \frac{d}{ds} I(\mathbf{u} + s\mathbf{v}) \Big|_{s=0} = \int_0^T \{ D_x \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} + D_y \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \dot{\mathbf{v}} \} dt.$$

Moreover, from (??), (??), Lemma 2.1 and the previous formula, we obtain

$$|\langle I'(\mathbf{u}), \mathbf{v} \rangle| \leq \|D_x \mathcal{L}\|_{L^1} \|\mathbf{v}\|_{L^\infty} + \|D_y \mathcal{L}\|_{L^\Psi} \|\dot{\mathbf{v}}\|_{L^\Phi} \leq C \|\mathbf{v}\|_{W^1 L^\Phi}$$

with a appropriate constant C . This completes the proof of the Gâteaux differentiability of I .

Step 4. The operator $I' : \mathcal{E}_d^\Phi(\lambda) \rightarrow [W^1 L_d^\Phi]^*$ is demicontinuous. This is a consequence of the continuity of the mappings $\mathbf{u} \mapsto D_x \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$ and $\mathbf{u} \mapsto D_y \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$. Indeed, if $\mathbf{u}_n, \mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$ with $\mathbf{u}_n \rightarrow \mathbf{u}$ in the norm of $W^1 L_d^\Phi$ and $\mathbf{v} \in W^1 L_d^\Phi$, then

$$\begin{aligned} \langle I'(\mathbf{u}_n), \mathbf{v} \rangle &= \int_0^T \{ D_x \mathcal{L}(t, \mathbf{u}_n, \dot{\mathbf{u}}_n) \cdot \mathbf{v} + D_y \mathcal{L}(t, \mathbf{u}_n, \dot{\mathbf{u}}_n) \cdot \dot{\mathbf{v}} \} dt \\ &\rightarrow \int_0^T \{ D_x \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} + D_y \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \dot{\mathbf{v}} \} dt \\ &= \langle I'(\mathbf{u}), \mathbf{v} \rangle. \end{aligned}$$

In order to prove item ??, it is necessary to see that the maps $\mathbf{u} \mapsto D_x \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$ and $\mathbf{u} \mapsto D_y \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$ are norm continuous from $\mathcal{E}_d^\Phi(\lambda)$ into L_d^1 and L_d^Ψ respectively. The

continuity of the first map has already been proved in step 1. Let $\mathbf{u}_n, \mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$ with $\|\mathbf{u}_n - \mathbf{u}\|_{W^1 L^\Phi} \rightarrow 0$. Therefore, there exist a subsequence $\mathbf{u}_{n_k} \in \mathcal{E}_d^\Phi(\lambda)$ and a function $h \in \Pi(E_1^\Phi, \lambda)$ such that (30) holds true. And, as $\Psi \in \Delta_2$ then the right hand side of (30) belongs to E_1^Ψ . Now, invoking Lemma ??, we prove that from any sequence \mathbf{u}_n which converges to \mathbf{u} in $W^1 L_d^\Phi$ we can extract a subsequence such that $D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k}) \rightarrow D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$ in the strong topology. The desired result is obtained by a standard argument.

The continuity of I' follows from the continuity of $D_x\mathcal{L}$ and $D_{\mathbf{y}}\mathcal{L}$ using the formula (28). \square

Acknowledgments

The authors are partially supported by a UNRC grant number 18/C417. The first author is partially supported by a UNSL grant number 22/F223.

References

- [1] S. Acinas, L. Buri, G. Giubergia, F. Mazzone, and E. Schwindt. Some existence results on periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting. *Nonlinear Analysis, TMA.*, 125:681 – 698, 2015.
- [2] R. Adams and J. Fournier. *Sobolev spaces*. Elsevier/Academic Press, Amsterdam, 2003.
- [3] W. Desch and R. Grimmer. On the well-posedness of constitutive laws involving dissipation potentials. *Trans. Amer. Math. Soc.*, (353):5095–5120, 2001.
- [4] Jan Gustavsson and Jaak Peetre. Interpolation of orlicz spaces. *Studia Mathematica*, 60(1):33–59, 1977.
- [5] H. Hudzik and L. Maligranda. Amemiya norm equals Orlicz norm in general. *Indag. Math. (N.S.)*, 11(4):573–585, 2000.
- [6] M. A. Krasnosel'skiĭ and Ja. B. Rutickiĭ. *Convex functions and Orlicz spaces*. P. Noordhoff Ltd., Groningen, 1961.
- [7] M.A. Krasnosel'skii, P.P. Zabreyko, E.I. Pustyl'nik, and P.E. Sobolevski. *Integral operators in spaces of summable functions*. Mechanics: Analysis. Springer Netherlands, 2011.
- [8] L. Maligranda. *Orlicz spaces and interpolation*, volume 5 of *Seminários de Matemática [Seminars in Mathematics]*. Universidade Estadual de Campinas, Departamento de Matemática, Campinas, 1989.
- [9] J. Mawhin and M. Willem. *Critical point theory and Hamiltonian systems*. Springer-Verlag, New York, 1989.

References

- [10] Nguen Hong Thai. The superposition operator in the Orlicz spaces of vector functions. *Dokl. Akad. Nauk BSSR*, 31:197â200, 1987.
- [11] Ryszard Pluciennik. Boundedness of the superposition operator in generalized Orlicz spaces of vector-valued functions. *Bull. Pol. Acad. Sci., Math.*, 33:531â540, 1985.
- [12] Ryszard Pluciennik. On some properties of the superposition operator in generalized Orlicz spaces of vector-valued functions. *Ann. Soc. Math. Pol., Ser. I, Commentat. Math.*, 25:321â337, 1985.
- [13] Ryszard Pluciennik. The superposition operator in Musielak-Orlicz spaces of vector-valued functions. Abstract analysis, Proc. 14th Winter Sch., Srní/Czech. 1986, Suppl. Rend. Circ. Mat. Palermo, II. Ser. 14, 411-417 (1987)., 1987.
- [14] M. M. Rao and Z. D. Ren. *Theory of Orlicz spaces*, volume 146. Marcel Dekker, Inc., New York, 1991.
- [15] G. Schappacher. A notion of Orlicz spaces for vector valued functions. *Appl. Math.*, 50(4):355–386, 2005.
- [16] M. S. Skaff. Vector valued orlicz spaces. ii. *Pacific J. Math.*, 28(2):413–430, 1969.
- [17] C.-L. Tang. Periodic solutions for nonautonomous second order systems with sublinear nonlinearity. *Proc. Amer. Math. Soc.*, 126(11):3263–3270, 1998.
- [18] C. L. Tang and X.-P. Wu. Periodic solutions for second order systems with not uniformly coercive potential. *J. Math. Anal. Appl.*, 259(2):386–397, 2001.
- [19] Chun-Lei Tang. Periodic solutions of non-autonomous second-order systems with γ -quasisubadditive potential. *Journal of Mathematical Analysis and Applications*, 189(3):671–675, 1995.
- [20] X. Tang and X. Zhang. Periodic solutions for second-order Hamiltonian systems with a p -Laplacian. *Ann. Univ. Mariae Curie-Skłodowska Sect. A*, 64(1):93–113, 2010.
- [21] Y. Tian and W. Ge. Periodic solutions of non-autonomous second-order systems with a p -Laplacian. *Nonlinear Anal.*, 66(1):192–203, 2007.
- [22] X.-P. Wu and C.-L. Tang. Periodic solutions of a class of non-autonomous second-order systems. *J. Math. Anal. Appl.*, 236(2):227–235, 1999.
- [23] F. Zhao and X. Wu. Periodic solutions for a class of non-autonomous second order systems. *J. Math. Anal. Appl.*, 296(2):422–434, 2004.