

# Periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting by the dual least action principle

Sonia Acinas \*

Dpto. de Matemática, Facultad de Ciencias Exactas y Naturales  
Universidad Nacional de La Pampa  
(L6300CLB) Santa Rosa, La Pampa, Argentina  
sonia.acinas@gmail.com

Fernando D. Mazzone †

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales  
Universidad Nacional de Río Cuarto  
(5800) Río Cuarto, Córdoba, Argentina,  
fmazzone@exa.unrc.edu.ar

## Abstract

## 1 Introduction

This paper deals with system of equations of the type:

$$\begin{cases} \frac{d}{dt} D_y \mathcal{L}(t, u(t), u'(t)) = D_x \mathcal{L}(t, u(t), u'(t)) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = u'(0) - u'(T) = 0, \end{cases} \quad (P_1)$$

where  $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $d \geq 1$ , is called the *Lagrange function* or *lagrangian* and the unknown function  $u : [0, T] \rightarrow \mathbb{R}^d$  is absolutely continuous. In other words, we are interested in finding *periodic weak solutions* of *Euler-Lagrange system*. This topic was deeply addressed for the *Lagrange function*

$$\mathcal{L}_{p,F}(t, x, y) := \frac{|y|^p}{p} + F(t, x), \quad (1)$$

---

\*SECyT-UNRC and FCEyN-UNLPam

†SECyT-UNRC, FCEyN-UNLPam and CONICET

**2010 AMS Subject Classification.** Primary: . Secondary: .

**Keywords and phrases.** .

for  $1 < p < \infty$ . For example, the classic book [Mawhin and Willem, 1989] deals mainly with problem  $(P_1)$ , for the lagrangian  $\mathcal{L}_{2,F}$ , through various methods: direct, dual action, minimax, etc. The results in [Mawhin and Willem, 1989] were extended and improved in several articles, see [Tang, 1995, Tang, 1998, Wu and Tang, 1999, Tang and Wu, 2001, Zhao and Wu, 2004] to cite some examples. Lagrange functions (1) for arbitrary  $1 < p < \infty$  were considered in [Tian and Ge, 2007, Tang and Zhang, 2010] and in this case  $(P_1)$  is reduced to the  $p$ -laplacian system

$$\begin{cases} \frac{d}{dt}(u'(t)|u'|^{p-2}) = \nabla F(t, u(t)) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = u'(0) - u'(T) = 0. \end{cases} \quad (P_2)$$

In this context, it is customary to call  $F$  a *potential function*, and it is assumed that  $F(t, x)$  is differentiable with respect to  $x$  for a.e.  $t \in [0, T]$  and the following conditions are verified:

(C)  $F$  and its gradient  $\nabla F$ , with respect to  $x \in \mathbb{R}^d$ , are Carathéodory functions, i.e. they are measurable functions with respect to  $t \in [0, T]$ , for every  $x \in \mathbb{R}^d$ , and they are continuous functions with respect to  $x \in \mathbb{R}^d$  for a.e.  $t \in [0, T]$ .

(A) For a.e.  $t \in [0, T]$ , it holds that

$$|F(t, x)| + |\nabla F(t, x)| \leq a(|x|)b(t). \quad (2)$$

In this inequality we assume that the function  $a : [0, +\infty) \rightarrow [0, +\infty)$  is continuous and non decreasing and  $0 \leq b \in L^1([0, T], \mathbb{R})$ .

In [Acinas et al., 2015] it was treated the case of a lagrangian  $\mathcal{L}$  which is lower bounded by a Lagrange function

$$\mathcal{L}_{\Phi, F}(t, x, y) = \Phi(|y|) + F(t, x), \quad (3)$$

where  $\Phi$  is an  $N$ -function (see section 2 for the definition of this concept). In the paper [Acinas et al., 2015] it was assumed a condition of *bounded oscillation* on  $F$  (see xxxxx below). In this paper we apply the dual method ([Mawhin and Willem, 1989, Ch. 3]) to obtain solutions of  $(P_1)$ .

## 2 Anisotropic Orlicz and Orlicz-Sobolev spaces

In this section, we give a short introduction to Orlicz and Orlicz-Sobolev spaces of vector valued functions associated to anisotropic Young functions  $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}_+$ , i.e. functions such that  $\Phi(x)$  depends on the direction of  $x$ , unlike the radial case where  $\Phi(x) = \Phi(|x|)$ . References for these topics are [Schappacher, 2005, Skaff, 1969, Desch and Grimmer, 2001].

In the framework of anisotropic Sobolev-Orlicz spaces, we can study system of  $p$ -laplacian equations as the following example shows.

**Example 1.** Let  $1 < p_1, p_2 < \infty$ . We define  $\Phi_{p_1, p_2} : \mathbb{R}^n \rightarrow \mathbb{R}_+$  by

$$\Phi_{p_1, p_2}(y_1, y_2) := \frac{|y_1|}{p_1} + \frac{|y_2|}{p_2}.$$

where  $|\cdot|$  is the Euclidean norm on  $\mathbb{R}^n$ . Suppose the following Lagrange function

$$\mathcal{L}(t, x, y) = \Phi_{p_1, p_2}(y) + F(t, x).$$

Then the equations  $(P_1)$  becomes

$$\begin{cases} \frac{d}{dt}(|u'_1|^{p_1-2}u'_1) = F_{x_1}(t, u) & \text{a.e. } t \in (0, T) \\ \frac{d}{dt}(|u'_2|^{p_2-2}u'_2) = F_{x_2}(t, u) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = u'(0) - u'(T) = 0, \end{cases} \quad (P_3)$$

EN CONSTRUCCION!!!!...FALTA ORDENAR, REVISAR, AGREGAR O QUITAR....

Las F de arriba....qué hacemos????

We apply the method of Lagrange multipliers to solve the problem

$$G(r) = \min\{\Phi(x, y) : \frac{1}{2}\|(x, y)\|_2^2 = \frac{r^2}{2}\}$$

then  $F(x, y) = \Phi(x, y) - \lambda \frac{1}{2}\|(x, y)\|_2^2$ .

Now, the first order equations are

$$\begin{cases} |x|^{p_1-2}x + \lambda x & = 0 \\ |y|^{p_2-2}y + \lambda y & = 0 \\ |x|^2 + |y|^2 & = r^2 \end{cases}$$

The critics points are  $(x, 0, -|x|^{p_1-2})$  and  $(0, y, -|y|^{p_2-2})$  with  $|x| = r$  and  $|y| = r$  respectively.

In case of  $2 \leq p_1 \leq p_2$ , the critic values are  $\Phi(x, 0) = \frac{r^{p_1}}{p_1}$  and  $\Phi(0, y) = \frac{r^{p_2}}{p_2}$

Now, suppose that  $x \neq 0$  and  $y \neq 0$  then  $|x|^2 + |y|^2 = r^2$  and  $|y| = |x|^{\frac{p_1-2}{p_2-2}}$  and  $\lambda = -|x|^{p_1-2}$ .

If  $r$  is bigger enough then  $G(r) \sim r^{p_1}$ . And, if  $r$  is small enough we have  $G(r) \sim r^{p_2}$ ????

If  $p_1 \leq 2$  and  $p_2 \leq 2$  with one of them different to 2, the second order criterium [?, Thm. —] implies that the minimum appear in  $x = 0$  or  $y = 0$ .

These equations are known in the literature as  $(p_1, p_2)$ -Laplacian system, see [Yang and Chen, 2013, Pasca and Wang, 2016, Yang and Chen, 2012, Pasca, 2010, Paşca and Tang, 2010, Pasca and Tang, 2011].

On the other hand, anisotropic Orlicz-Sobolev spaces allow to simplify the writing, and they provide the natural frame of statements of the type [Tian and Ge, 2007, Lemma 3.1]. This type of question was what motivated us to use these spaces.

Hereafter we denote by  $\mathbb{R}^+$  the set of all non negative real numbers. A function  $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}_+$  is called an *Young's function* if  $\Phi$  is convex,  $\Phi(0) = 0$ ,  $\Phi(-x) = \Phi(x)$  and  $\Phi(x) \rightarrow +\infty$ , when  $|x| \rightarrow +\infty$ . Additionally, we assume that the Young's functions which we deal with, satisfy that  $\Phi(x) > 0$  when  $x \neq 0$ . Following [Schappacher, 2005] we say that  $\Phi$  is an  $N_\infty$ -function if

$$\lim_{|x| \rightarrow \infty} \frac{\Phi(x)}{|x|} = +\infty.$$

Given a Young's function  $\Phi$ , we define function  $A_\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  by

$$A_\Phi(s) = \min \{ \Phi(x) \mid \|x\| = s \}, \quad (4)$$

Let us establish some elementary properties of  $A_\Phi$  that we will use in this article.

**Proposition 2.1.** *The function  $A_\Phi$  has the following properties:*

1.  $A_\Phi$  is continuous,
2.  $A_\Phi(s)/s$  is increasing,
3.  $A_\Phi(|x|)$  is the greatest radial minorant of  $\Phi(x)$ ,
4.  $\Phi$  is  $N_\infty$  if and only if  $A_\Phi$  is.

*Proof.* It is well known that finite and convex functions defined in finite dimensional vectorial spaces are locally Lipschitz functions (see [Clarke, 2013]). This fact imply item 1 immediately.

In order to prove item 2, suppose  $0 < r < s$  and  $x \in \mathbb{R}^d$  with  $A_\Phi(s) = \Phi(x)$ . Then, from the definition of  $A_\Phi$  and the convexity of  $\Phi$ ,

$$\frac{A_\Phi(r)}{r} \leq \frac{\Phi\left(\frac{r}{s}x\right)}{r} \leq \frac{\Phi(x)}{s} = \frac{A_\Phi(s)}{s}.$$

Property in items 3 and 4 are obtained easily. □

We also say that  $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}^+$  satisfies the  $\Delta_2^\infty$ -condition, denoted by  $\Phi \in \Delta_2^\infty$ , if there exist constants  $K > 0$  and  $M \geq 0$  such that

$$\Phi(2x) \leq KH(x), \quad (5)$$

for every  $|x| \geq M$ .

If  $\Phi$  is a Young's function we define its *Fenchel conjugate*  $\Phi^* : \mathbb{R}^d \rightarrow \mathbb{R}^+$  by:

$$\Phi^*(y) = \sup_{x \in \mathbb{R}^d} x \cdot y - \Phi(x) \quad (6)$$

We denote by  $\mathcal{M} := \mathcal{M}([0, T], \mathbb{R}^d)$ , with  $d \geq 1$ , the set of all measurable functions (i.e. functions which are limits of simple functions) defined on  $[0, T]$  with values on  $\mathbb{R}^d$  and we write  $u = (u_1, \dots, u_d)$  for  $u \in \mathcal{M}$ . For the set of functions  $\mathcal{M}$ , as for other similar sets, we will omit the reference to codomain  $\mathbb{R}^d$  when  $d = 1$ .

Given an  $N$ -function  $\Phi$  we define the *modular function*  $\rho_\Phi : \mathcal{M} \rightarrow \mathbb{R}^+ \cup \{+\infty\}$  by

$$\rho_\Phi(u) := \int_0^T \Phi(u) dt.$$

Here  $|\cdot|$  is the euclidean norm of  $\mathbb{R}^d$ . Now, we introduce the *Orlicz class*  $C^\Phi = C^\Phi([0, T], \mathbb{R}^d)$  by setting

$$C^\Phi := \{u \in \mathcal{M} \mid \rho_\Phi(u) < \infty\}. \quad (7)$$

The Orlicz space  $L^\Phi = L^\Phi([0, T], \mathbb{R}^d)$  is the linear hull of  $C^\Phi$ ; equivalently,

$$L^\Phi := \{u \in \mathcal{M} \mid \exists \lambda > 0 : \rho_\Phi(\lambda u) < \infty\}. \quad (8)$$

The Orlicz space  $L^\Phi$  equipped with the *Luxemburg norm*

$$\|u\|_{L^\Phi} := \inf \left\{ \lambda \left| \rho_\Phi \left( \frac{v}{\lambda} \right) dt \leq 1 \right. \right\},$$

is a Banach space. By  $u \cdot v$  we denote the usual dot product in  $\mathbb{R}^d$  between  $u$  and  $v$ .

The subspace  $E^\Phi = E^\Phi([0, T], \mathbb{R}^d)$  is defined as the closure in  $L^\Phi$  of the subspace  $L^\infty([0, T], \mathbb{R}^d)$  of all  $\mathbb{R}^d$ -valued essentially bounded functions. It is shown that (see [Schappacher, 2005, Thm. 5.1])  $u \in E^\Phi$  if and only if  $\rho_\Phi(\lambda u) < \infty$  for any  $\lambda > 0$ . The equality  $L^\Phi = E^\Phi$  is true if and only if  $\Phi \in \Delta_2^\infty$  (see [Schappacher, 2005, Thm. 5.2]). Another alternative characterization of  $E^\Phi$ , which is particularly useful for us, is that  $u \in E^\Phi$  if and only if  $u$  has *absolutely continuous norm*, i.e. if  $E_n \subset [0, T]$ ,  $n = 1, 2, \dots$  then  $\|\chi_{E_n} u\| \rightarrow 0$  when  $|E_n| \rightarrow 0$ .

A generalized version of *Hölder's inequality* holds in Orlicz spaces (see [Skaff, 1969, Thm. 4.1]). Namely, if  $u \in L^\Phi$  and  $v \in L^{\Phi^*}$  then  $u \cdot v \in L^1$  and

$$\int_0^T v \cdot u \, dt \leq 2 \|u\|_{L^\Phi} \|v\|_{L^{\Phi^*}}. \quad (9)$$

Like in [Krasnosel'skiĭ and Rutickiĭ, 1961] we will consider the subset  $\Pi(E^\Phi, r)$  of  $L^\Phi$  given by

$$\Pi(E^\Phi, r) := \{u \in L^\Phi \mid d(u, E^\Phi) < r\}.$$

This set is related to the Orlicz class  $C^\Phi$  by means of inclusions, namely,

$$\Pi(E^\Phi, r) \subset rC^\Phi \subset \overline{\Pi(E^\Phi, r)} \quad (10)$$

for any positive  $r$  (see [Schappacher, 2005, Thm. 5.6]). If  $\Phi \in \Delta_2^\infty$ , then the sets  $L^\Phi$ ,  $E^\Phi$ ,  $\Pi(E^\Phi, r)$  and  $C^\Phi$  are equal.

Following to [Desch and Grimmer, 2001] we introduce the next definition.

**Definition 2.2.** Let  $u_n, u \in L^\Phi([0, T], \mathbb{R}^d)$ . We say that  $u_n$  converges monotonically to  $u$  if there exists  $\alpha_n \in L^\infty([0, T], \mathbb{R})$ ,  $n = 1, 2, \dots$ , such that  $0 \leq \alpha_n(t) \leq \alpha_{n+1}(t)$ ,  $\alpha_n(t) \rightarrow 1$  a.e., when  $n \rightarrow \infty$  and  $u_n(t) = \alpha_n(t)u(t)$ .

As usual, if  $(X, \|\cdot\|_X)$  is a normed space and  $(Y, \|\cdot\|_Y)$  is a linear subspace of  $X$ , we write  $Y \hookrightarrow X$  and we say that  $Y$  is *embedded* in  $X$  when there exists  $C > 0$  such that  $\|y\|_X \leq C\|y\|_Y$  for any  $y \in Y$ . With this notation, Hölder's inequality states that  $L^{\Phi^*} \hookrightarrow [L^\Phi]^*$ , where a function  $v \in L^{\Phi^*}$  is associated to  $\xi_v \in [L^\Phi]^*$  being

$$\xi_v(u) = \langle \xi_v, u \rangle = \int_0^T v \cdot u \, dt, \quad (11)$$

In [Desch and Grimmer, 2001, Thm 2.9] it was characterized a subspace of  $[L^\Phi]^*$  which can be identified with  $L^{\Phi^*}$ .

**Proposition 2.3.** *Let  $F \in [L^\Phi([0, T], \mathbb{R}^d)]^*$ . Then the following statements are equivalent*

1.  $\xi \in L^{\Phi^*}([0, T], \mathbb{R}^d)$
2.  $\xi$  satisfies the monotone convergence property, which is if  $u_n$  converges monotonically to  $u$  then  $\langle \xi, u_n \rangle \rightarrow \langle \xi, u \rangle$ .

If  $\Phi \in \Delta_2^\infty$  and  $\Phi$  is  $N_\infty$  then  $L^{\Phi^*}([0, T], \mathbb{R}^d) = [L^\Phi([0, T], \mathbb{R}^d)]^*$  (see [Desch and Grimmer, 2001, Thm. 2.9, Thm. 2.10]).

We define the Sobolev-Orlicz space  $W^1 L^\Phi$  by

$$W^1 L^\Phi([0, T], \mathbb{R}^d) := \{u \mid u \text{ is absolutely continuous on } [0, T] \text{ and } u' \in L^\Phi([0, T], \mathbb{R}^d)\}.$$

$W^1 L^\Phi([0, T], \mathbb{R}^d)$  is a Banach space when equipped with the norm

$$\|u\|_{W^1 L^\Phi} = \|u\|_{L^\Phi} + \|u'\|_{L^\Phi}. \quad (12)$$

And, we introduce the following subspaces of  $W^1 L^\Phi$

$$\begin{aligned} W^1 E^\Phi &= \{u \in W^1 L^\Phi \mid u' \in E^\Phi\}, \\ W^1 E_T^\Phi &= \{u \in W^1 E^\Phi \mid u(0) = u(T)\}. \end{aligned} \quad (13)$$

We will use repeatedly the decomposition  $u = \bar{u} + \tilde{u}$  for a function  $u \in L^1([0, T])$  where  $\bar{u} = \frac{1}{T} \int_0^T u(t) dt$  and  $\tilde{u} = u - \bar{u}$ .

The following lemma is an elementary generalization to anisotropic Sobolev-Orlicz spaces of known results of Sobolev spaces.

**Lemma 2.4.** *Let  $\Phi : \mathbb{R}^d \rightarrow [0, +\infty)$  be a Young's function and let  $u \in W^1 L^\Phi([0, T], \mathbb{R}^d)$ . Let  $A_\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be the function defined by (4). Then*

1. For every  $s, t \in [0, T]$ ,  $s \neq t$ ,

$$|u(t) - u(s)| \leq \|u'\|_{L^\Phi} |s - t| A_\Phi^{-1} \left( \frac{1}{|s - t|} \right) \quad (\text{Morrey's inequality})$$

$$\|u\|_{L^\infty} \leq A_\Phi^{-1} \left( \frac{1}{T} \right) \max\{1, T\} \|u\|_{W^1 L^\Phi} \quad (\text{Sobolev's inequality})$$

2. We have  $\tilde{u} \in L^\infty([0, T], \mathbb{R}^d)$  and

$$\|\tilde{u}\|_{L^\infty} \leq T A_\Phi^{-1} \left( \frac{1}{T} \right) \|u'\|_{L^\Phi} \quad (\text{Sobolev-Wirtinger's inequality})$$

3. If  $\Phi$  is  $N_\infty$  then the space  $W^1 L^\Phi([0, T], \mathbb{R}^d)$  is compactly embedded in the space of continuous functions  $C([0, T], \mathbb{R}^d)$ .

*Proof.* By the absolutely continuity of  $u$ , Jensen's inequality and the definition of the Luxemburg norm, we have

$$\begin{aligned} \Phi\left(\frac{u(t) - u(s)}{\|u'\|_{L^\Phi} |s - t|}\right) &\leq \Phi\left(\frac{1}{|s - t|} \int_s^t \frac{u'(r)}{\|u'\|_{L^\Phi}} dr\right) \\ &\leq \frac{1}{|s - t|} \int_s^t \Phi\left(\frac{u'(r)}{\|u'\|_{L^\Phi}}\right) dr \leq \frac{1}{|s - t|}. \end{aligned}$$

By Proposition 2.1(3) we have  $A_\Phi^{-1}\Phi(x) \geq |x|$ , therefore we get

$$\frac{|u(t) - u(s)|}{\|u'\|_{L^\Phi} |s - t|} \leq A_\Phi^{-1}\left(\frac{1}{|s - t|}\right),$$

then 1 holds.

Now, we use Morrey's inequality and Proposition 2.1 (2) and we have

$$\begin{aligned} |u(t) - \bar{u}| &= \left| \frac{1}{T} \int_0^T u(t) - u(s) ds \right| \\ &\leq \frac{1}{T} \int_0^T |u(t) - u(s)| ds \\ &\leq \|u'\|_{L^\Phi} T A_\Phi^{-1}\left(\frac{1}{T}\right) \end{aligned}$$

In order to prove the Sobolev's inequality, we note that, using Jensen's inequality and the definition of  $\|u\|_{L^\Phi}$ , we obtain

$$\Phi\left(\frac{\bar{u}}{\|u\|_{L^\Phi}}\right) \leq \frac{1}{T} \int_0^T \Phi\left(\frac{u(s)}{\|u\|_{L^\Phi}}\right) ds \leq \frac{1}{T}$$

Then by Proposition 2.1(3)

$$|\bar{u}| \leq A_\Phi^{-1}\left(\frac{1}{T}\right) \|u\|_{L^\Phi}.$$

Therefore, from this and (Sobolev-Wirtinger's inequality) we get

$$\begin{aligned} \|u\|_{L^\infty} &\leq |\bar{u}| + \|\tilde{u}\|_{L^\infty} \\ &\leq A_\Phi^{-1}\left(\frac{1}{T}\right) \|u\|_{L^\Phi} + T A_\Phi^{-1}\left(\frac{1}{T}\right) \|u'\|_{L^\Phi} \\ &\leq A_\Phi^{-1}\left(\frac{1}{T}\right) \max\{1, T\} \|u\|_{W^1 L^\Phi} \end{aligned}$$

In order to prove item 3, we take a bounded sequence  $u_n$  in  $W^1 L^\Phi([0, T], \mathbb{R}^d)$ . Since  $\Phi$  is  $N_\infty$ , from Proposition 2.1(4) we obtain  $s A_\Phi^{-1}(1/s) \rightarrow 0$  when  $s \rightarrow 0$ . Therefore (Morrey's inequality) implies that  $u_n$  are equicontinuous. Furthermore (Sobolev's inequality) implies that  $u_n$  is bounded in  $C([0, T], \mathbb{R}^d)$ . Therefore by the Arzela-Ascoli Theorem we obtain a subsequence  $n_k$  and  $u \in C([0, T], \mathbb{R}^d)$  with  $u_{n_k} \rightarrow u$  in  $C([0, T], \mathbb{R}^d)$ .  $\square$

**Lemma 2.5.** *Let  $\{u_n\}_{n \in \mathbb{N}}$  be a sequence of functions in  $\Pi(E^\Phi, 1)$  converging to  $u \in \Pi(E^\Phi, 1)$  in the  $L^\Phi$ -norm. Then, there exist a subsequence  $u_{n_k}$  and a real valued function  $h \in L^1([0, T], \mathbb{R})$  such that  $u_{n_k} \rightarrow u$  a.e. and  $\Phi(u_{n_k}) \leq h$  a.e.*

*Proof.* Since  $d(u, E^\Phi) < 1$  and  $u_n$  converges to  $u$ , there exists  $u_0 \in E^\Phi$ , a subsequence of  $u_n$  (again denoted  $u_n$ ) and  $0 < r < 1$  such that  $d(u_n, u_0) < r$ . Let  $\lambda_0 \in (r, 1)$ . By extracting more subsequences, if necessary, we can assume that  $u_n \rightarrow u$  a.e. and

$$\lambda_n := \|u_{n+1} - u_n\|_{L^\Phi} < \frac{1 - \lambda_0}{2^n}, \quad \text{for } n \geq 1.$$

We can assume  $\lambda_n > 0$  for every  $n = 0, \dots$

Let  $\lambda := 1 - \sum_{n=0}^{\infty} \lambda_n$  and define  $h : [0, T] \rightarrow \mathbb{R}$  by

$$h(x) = \lambda \Phi\left(\frac{u_0}{\lambda}\right) + \sum_{n=0}^{\infty} \lambda_n \Phi\left(\frac{u_{n+1} - u_n}{\lambda_n}\right). \quad (14)$$

Note that  $\sum_{n=0}^{\infty} \lambda_n + \lambda = 1$ , therefore for any  $n = 1, \dots$

$$\begin{aligned} \Phi(u_n) &= \Phi\left(\lambda \frac{u_0}{\lambda} + \sum_{j=0}^{n-1} \lambda_j \frac{u_{j+1} - u_j}{\lambda_j}\right) \\ &\leq \lambda \Phi\left(\frac{u_0}{\lambda}\right) + \sum_{j=0}^{n-1} \lambda_j \Phi\left(\frac{u_{j+1} - u_j}{\lambda_j}\right) \leq h \end{aligned}$$

Since  $v \in E^\Phi \subset C^\Phi$  and  $E^\Phi$  is a subspace we have that  $\Phi(u_0/\lambda) \in L^1([0, T], \mathbb{R})$ . On the other hand  $\|u_{n+1} - u_n\|_{L^\Phi} \leq \lambda_n$ , therefore

$$\int_0^T \Phi\left(\frac{u_{j+1} - u_j}{\lambda_j}\right) dt \leq 1.$$

Then  $h \in L^1([0, T], \mathbb{R})$ . □

### 3 Differentiability Gateaux of action integrals in anisotropic Orlicz spaces

In this section we give a brief introduction to superposition operators between anisotropic Orlicz Spaces. We apply these results to obtain Gateaux differentiability of action integrals associated to lagrangian functions defined in Sobolev-Orlicz spaces.

Henceforth we assume that  $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a *Carathéodory function*, i.e.

- (C)  $f$  is measurable with respect to  $t \in [0, T]$  for every  $x \in \mathbb{R}^d$ , and  $f$  is a continuous function with respect to  $x \in \mathbb{R}^d$  for a.e.  $t \in [0, T]$ .



**Definition 3.1.** For  $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  we denote by  $\mathbf{f}$  the Nemytskii (or superposition) operator defined for functions  $u : [0, T] \rightarrow \mathbb{R}^d$  by

$$\mathbf{f}u(t) = f(t, u(t))$$

In the following Theorem we enumerate some known properties for superposition operators defined in anisotropic Orlicz spaces of vectorial functions. For the proofs see [Krasnosel'skii et al., 2011] for scalar functions and [Pluciennik, 1987, Pluciennik, 1985b, Pluciennik, 1985a] for the generalization to  $\mathbb{R}^d$ -valued (moreover Banach spaces valued) functions in a anisotropic Orlicz Spaces (moreover modular anisotropic spaces).

**Theorem 3.2.** We assume that  $f$  satisfies condition ((C)) and that  $\Phi_1, \Phi_2 : \mathbb{R}^d \rightarrow [0, +\infty)$  are anisotropic Young functions. Then

1. Measurability. The operator  $\mathbf{f}$  maps measurable function into measurable functions
2. Extensibility. If the operator  $\mathbf{f}$  acts from the ball  $B_{L^{\Phi_1}}(r) := \{u \in L^{\Phi_1} \mid \|u\|_{L^{\Phi_1}} < r\}$  into the space  $L^{\Phi_2}$  or the space  $E^{\Phi_2}$  then  $\mathbf{f}$  can be extended to  $\Pi(E^{\Phi_1}, r)$  into space  $L^{\Phi_2}$  or  $E^{\Phi_2}$  respectively.
3. Continuity. If the operator  $\mathbf{f}$  acts from  $\Pi(E^{\Phi_1}, r)$  into space  $E^{\Phi_2}$ , then  $\mathbf{f}$  is continuous.

## Acknowledgments

The authors are partially supported by a UNRC grant number 18/C417. The first author is partially supported by a UNSL grant number 22/F223.

## References

- [Acinas et al., 2015] Acinas, S., Buri, L., Giubergia, G., Mazzone, F., and Schwindt, E. (2015). Some existence results on periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting. *Nonlinear Analysis, TMA.*, 125:681 – 698.
- [Clarke, 2013] Clarke, F. (2013). *Functional Analysis, Calculus of Variations and Optimal Control*. Graduate Texts in Mathematics.
- [Desch and Grimmer, 2001] Desch, W. and Grimmer, R. (2001). On the well-posedness of constitutive laws involving dissipation potentials. *Trans. Amer. Math. Soc.*, (353):5095–5120.
- [Krasnosel'skii et al., 2011] Krasnosel'skii, M., Zabreyko, P., Pustynnik, E., and Sobolevski, P. (2011). *Integral operators in spaces of summable functions*. Mechanics: Analysis. Springer Netherlands.
- [Krasnosel'skiĭ and Rutickiĭ, 1961] Krasnosel'skiĭ, M. A. and Rutickiĭ, J. B. (1961). *Convex functions and Orlicz spaces*. P. Noordhoff Ltd., Groningen.

- [Mawhin and Willem, 1989] Mawhin, J. and Willem, M. (1989). *Critical point theory and Hamiltonian systems*. Springer-Verlag, New York.
- [Pasca, 2010] Pasca, D. (2010). Periodic solutions of a class of nonautonomous second order differential systems with  $(q, p)$ -laplacian. *Bulletin of the Belgian Mathematical Society-Simon Stevin*, 17(5):841–851.
- [Paşca and Tang, 2010] Paşca, D. and Tang, C.-L. (2010). Some existence results on periodic solutions of nonautonomous second-order differential systems with  $(q, p)$ -laplacian. *Applied Mathematics Letters*, 23(3):246–251.
- [Pasca and Tang, 2011] Pasca, D. and Tang, C.-L. (2011). Some existence results on periodic solutions of ordinary  $(q, p)$ -laplacian systems. *Journal of applied mathematics & informatics*, 29(1.2):39–48.
- [Pasca and Wang, 2016] Pasca, D. and Wang, Z. (2016). On periodic solutions of nonautonomous second order hamiltonian systems with  $(q, p)$ -laplacian. *Electronic Journal of Qualitative Theory of Differential Equations*, 2016(106):1–9.
- [Płuciennik, 1985a] Płuciennik, R. (1985a). Boundedness of the superposition operator in generalized Orlicz spaces of vector-valued functions. *Bull. Pol. Acad. Sci., Math.*, 33:531–540.
- [Płuciennik, 1985b] Płuciennik, R. (1985b). On some properties of the superposition operator in generalized Orlicz spaces of vector-valued functions. *Ann. Soc. Math. Pol., Ser. I, Commentat. Math.*, 25:321–337.
- [Płuciennik, 1987] Płuciennik, R. (1987). The superposition operator in Musielak-Orlicz spaces of vector-valued functions. Abstract analysis, Proc. 14th Winter Sch., Srní/Czech. 1986, Suppl. Rend. Circ. Mat. Palermo, II. Ser. 14, 411–417 (1987).
- [Schappacher, 2005] Schappacher, G. (2005). A notion of Orlicz spaces for vector valued functions. *Appl. Math.*, 50(4):355–386.
- [Skaff, 1969] Skaff, M. S. (1969). Vector valued orlicz spaces. ii. *Pacific J. Math.*, 28(2):413–430.
- [Tang, 1995] Tang, C.-L. (1995). Periodic solutions of non-autonomous second-order systems with  $\gamma$ -quasisubadditive potential. *Journal of Mathematical Analysis and Applications*, 189(3):671–675.
- [Tang, 1998] Tang, C.-L. (1998). Periodic solutions for nonautonomous second order systems with sublinear nonlinearity. *Proc. Amer. Math. Soc.*, 126(11):3263–3270.
- [Tang and Wu, 2001] Tang, C. L. and Wu, X.-P. (2001). Periodic solutions for second order systems with not uniformly coercive potential. *J. Math. Anal. Appl.*, 259(2):386–397.
- [Tang and Zhang, 2010] Tang, X. and Zhang, X. (2010). Periodic solutions for second-order Hamiltonian systems with a  $p$ -Laplacian. *Ann. Univ. Mariae Curie-Skłodowska Sect. A*, 64(1):93–113.

## References

---

- [Tian and Ge, 2007] Tian, Y. and Ge, W. (2007). Periodic solutions of non-autonomous second-order systems with a  $p$ -Laplacian. *Nonlinear Anal.*, 66(1):192–203.
- [Wu and Tang, 1999] Wu, X.-P. and Tang, C.-L. (1999). Periodic solutions of a class of non-autonomous second-order systems. *J. Math. Anal. Appl.*, 236(2):227–235.
- [Yang and Chen, 2012] Yang, X. and Chen, H. (2012). Periodic solutions for a non-linear  $(q, p)$ -laplacian dynamical system with impulsive effects. *Journal of Applied Mathematics and Computing*, 40(1-2):607–625.
- [Yang and Chen, 2013] Yang, X. and Chen, H. (2013). Existence of periodic solutions for sublinear second order dynamical system with  $(q, p)$ -laplacian. *Mathematica Slovaca*, 63(4):799–816.
- [Zhao and Wu, 2004] Zhao, F. and Wu, X. (2004). Periodic solutions for a class of non-autonomous second order systems. *J. Math. Anal. Appl.*, 296(2):422–434.