

# Some existence results on periodic solutions of Euler-Lagrange equations in an Orlicz-Sobolev space setting

Sonia Acinas \*

Instituto de Matemática Aplicada San Luis (CONICET-UNSL)  
(5700) San Luis, Argentina  
Universidad Nacional de La Pampa  
(6300) Santa Rosa, La Pampa, Argentina  
sonia.acinas@gmail.com

Leopoldo Buri †

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales  
Universidad Nacional de Río Cuarto  
(5800) Río Cuarto, Córdoba, Argentina,  
lburi@exa.unrc.edu.ar

Graciela Giubergia ‡

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales  
Universidad Nacional de Río Cuarto  
(5800) Río Cuarto, Córdoba, Argentina,  
ggiubergia@exa.unrc.edu.ar

Fernando D. Mazzone §

Dpto. de Matemática, Facultad de Ciencias Exactas, Físico-Químicas y Naturales  
Universidad Nacional de Río Cuarto  
(5800) Río Cuarto, Córdoba, Argentina,  
fmazzone@exa.unrc.edu.ar

Erica L. Schwindt ¶

Université d'Orléans, Laboratoire MAPMO, CNRS, UMR 7349,  
Fédération Denis Poisson, FR 2964,  
Bâtiment de Mathématiques, BP 6759, 45067 Orléans Cedex 2, France,  
leris98@gmail.com

### Abstract

In this paper we consider the problem of finding periodic solutions of certain Euler-Lagrange equations. We employ the direct method of the calculus of variations, that is we obtain solutions minimizing certain functional  $I$ . We give conditions which ensure that  $I$  is finitely defined and differentiable on certain subsets of Orlicz-Sobolev spaces  $W^1 L^\Phi$  associated to an  $N$ -function  $\Phi$ . We show that, in some sense, it is necessary for the coercitivity that the complementary function of  $\Phi$  satisfy the  $\Delta_2$ -condition. We conclude by discussing conditions for existence of minima for  $I$ .

## 1 Introduction

This paper is concerned with the existence of periodic solutions of the problem

$$\begin{cases} \frac{d}{dt} D_y \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) = D_x \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) & \text{a.e. } t \in (0, T) \\ \mathbf{u}(0) - \mathbf{u}(T) = \dot{\mathbf{u}}(0) - \dot{\mathbf{u}}(T) = 0 \end{cases} \quad (1)$$

where  $T > 0$ ,  $\mathbf{u} : [0, T] \rightarrow \mathbb{R}^d$  is absolutely continuous and the *Lagrangian*  $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function satisfying the conditions

$$|\mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( b(t) + \Phi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right), \quad (2)$$

$$|D_x \mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( b(t) + \Phi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right), \quad (3)$$

$$|D_y \mathcal{L}(t, \mathbf{x}, \mathbf{y})| \leq a(|\mathbf{x}|) \left( c(t) + \varphi \left( \frac{|\mathbf{y}|}{\lambda} + f(t) \right) \right). \quad (4)$$

In these inequalities we assume that  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$ ,  $\lambda > 0$ ,  $\Phi$  is an  $N$ -function (see section Preliminaries for definitions),  $\varphi$  is the right continuous derivative of  $\Phi$  and the non negative functions  $b$ ,  $c$  and  $f$  belong to certain Banach spaces that will be introduced later.

It is well known that problem (1) comes from a variational one, that is, a solution of (1) is a critical point of the *action integral*

$$I(\mathbf{u}) = \int_0^T \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) dt. \quad (5)$$

Variational problems and hamiltonian systems have been studied extensively. Classic references of these subjects are [25, 30, 15]. Problems like (1) have maintained the interest of researchers as the recent literature on the topic testifies. For lagrangian functions of the type  $\mathcal{L}(t, \mathbf{x}, \mathbf{y}) = \frac{|\mathbf{y}|^2}{2} + F(t, \mathbf{x})$  many solvability conditions have

---

\*SECyT-UNRC, UNSL and CONICET

†SECyT-UNRC

‡SECyT-UNRC and CONICET

§SECyT-UNRC and CONICET

¶ANR. AVENTURES - ANR-12-BLAN-BS01-0001-01

**2010 AMS Subject Classification.** Primary: . Secondary: .

**Keywords and phrases.** .

been given expanding the results in [25]. In [32] the function  $F$  was split up into two potentials, one of them with a property of subadditivity and the other with a bounded gradient. In [31] it was required a certain sublinearity condition on the gradient of the potential  $F$ ; and, in [36] it was considered a potential  $F$  given by a sum of a subconvex function and a subquadratic one. In [33] the uniform coercivity of  $\int_0^T F(t, \mathbf{x}) dt$  was replaced by local coercivity of  $F$  in some positive measure subset of  $[0, T]$ . In [39], the authors took a similar potential to that in [36] getting new solvability conditions and they also studied the case in which the two potentials do not have any convexity.

The Lagrangian  $\mathcal{L}(t, \mathbf{x}, \mathbf{y}) = \frac{|\mathbf{y}|^p}{p} + F(t, \mathbf{x})$  for  $p > 1$  was treated in more recent papers. By using the dual least action principle, in [35] it was performed the extension of some results given in [25]; and, in [34] the authors improved the work done in [36]. On the other hand, by the minimax methods in critical point theory some existence theorems were obtained. In [37] it was employed a subquadratic potential  $F$  in Rabinowitz's sense and in [38]  $F$  was taken as in [31].

Another source of problems, close to our proposal, is the one in which a *p-laplacian-like* operator is involved. Assuming that the function  $\varphi$  is a homeomorphism from  $\mathbb{R}^d$  into itself, it is considered the differential operator  $\mathbf{u} \mapsto (\varphi(\mathbf{u}'))'$ . In [5, 6, 7, 23, 24], using the Leray-Schauder degree theory, some existence results of solutions of equations like  $(\varphi(\mathbf{u}'))' = \mathbf{f}(t, \mathbf{u}(t), \mathbf{u}'(t))$  were obtained under different boundary conditions (periodic, Dirichlet, von Neumann) and where  $\mathbf{f}$  is not necessarily a gradient. We point out that our approach differs from that of previous articles because we tackle the direct method of the calculus of variations.

In the Orlicz-Sobolev space setting, in [17] a constrained minimization problem associated to the existence of eigenvalues for certain differential operators involving  $N$ -functions was studied. Slightly away from the problems to be treated in this paper, we can mention [9, 11] where A. Cianchi dealt with the regularity of minimizers of action integrals defined on several variable functions.

In this article we consider lagrangian functions defined on Orlicz-Sobolev spaces  $W^1 L^\Phi$  (see [3, 21, 26, 27]) and we use the direct method of calculus of variations. The exposition is organized as follows. In Section 2 we enumerate results related to Orlicz spaces, Orlicz-Sobolev spaces and composition operators. Almost all results in this section are essentially known. Conditions (2), (3) and (4) are the means to ensure that  $I$  is finitely defined on a non trivial subset of  $W^1 L_d^\Phi$  and  $I$  is Gâteaux differentiable in this subspace. We develop these issues in Theorem 3.2 of Section 3. In Section 4 we prove that critical points of (5) are solution of (1). Conditions for the coercitivity of action integrals are discussed in Section 5. Finally, our main theorem about existence of solutions of (1) is introduced and proved in Section 6.

We put an emphasis on that we take care of using  $\Delta_2$ -condition. In some results where we have used it, we show that  $\Delta_2$ -condition is necessary in a certain sense (see, for example, Lemma 5.2).

## 2 Preliminaries

For reader convenience, we give a short introduction to Orlicz and Orlicz-Sobolev spaces of vector valued functions and a list of results that we will use throughout the

article. Classic references for Orlicz spaces of real valued functions are [3, 21, 26]. For Orlicz spaces of vector valued functions, see [29] and the references therein.

Hereafter we denote by  $\mathbb{R}^+$  the set of all non negative real numbers. A function  $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is called an *N-function* if  $\Phi$  is given by

$$\Phi(t) = \int_0^t \varphi(\tau) d\tau, \quad \text{for } t \geq 0,$$

where  $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a right continuous nondecreasing function satisfying  $\varphi(0) = 0$ ,  $\varphi(t) > 0$  for  $t > 0$  and  $\lim_{t \rightarrow \infty} \varphi(t) = +\infty$ .

Given a function  $\varphi$  as above, we consider the so-called right inverse function  $\psi$  of  $\varphi$  which is defined by  $\psi(s) = \sup_{\varphi(t) \leq s} t$ . The function  $\psi$  satisfies the same properties as the function  $\varphi$ , therefore we have an *N-function*  $\Psi$  such that  $\Psi' = \psi$ . The function  $\Psi$  is called the *complementary function* of  $\Phi$ .

We say that  $\Phi$  satisfies the  $\Delta_2$ -condition, denoted by  $\Phi \in \Delta_2$ , if there exist constants  $K > 0$  and  $t_0 \geq 0$  such that  $\Phi(2t) \leq K\Phi(t)$  for every  $t \geq t_0$ . If  $t_0 = 0$ , we say that  $\Phi$  satisfies the  $\Delta_2$ -condition globally ( $\Phi \in \Delta_2$  globally).

Let  $d$  be a positive integer. We denote by  $\mathcal{M}_d := \mathcal{M}_d([0, T])$  the set of all measurable functions defined on  $[0, T]$  with values on  $\mathbb{R}^d$  and we write  $\mathbf{u} = (u_1, \dots, u_d)$  for  $\mathbf{u} \in \mathcal{M}_d$ . In this paper we adopt the convention that bold symbols denote points in  $\mathbb{R}^d$ .

Given an *N-function*  $\Phi$  we define the *modular function*  $\rho_\Phi : \mathcal{M}_d \rightarrow \mathbb{R}^+ \cup \{+\infty\}$  by

$$\rho_\Phi(\mathbf{u}) := \int_0^T \Phi(|\mathbf{u}|) dt.$$

Here  $|\cdot|$  is the euclidean norm of  $\mathbb{R}^d$ . The *Orlicz class*  $C_d^\Phi = C_d^\Phi([0, T])$  is given by

$$C_d^\Phi := \{\mathbf{u} \in \mathcal{M}_d | \rho_\Phi(\mathbf{u}) < \infty\}. \quad (6)$$

The *Orlicz space*  $L_d^\Phi = L_d^\Phi([0, T])$  is the linear hull of  $C_d^\Phi$ ; equivalently,

$$L_d^\Phi := \{\mathbf{u} \in \mathcal{M}_d | \exists \lambda > 0 : \rho_\Phi(\lambda \mathbf{u}) < \infty\}. \quad (7)$$

The Orlicz space  $L_d^\Phi$  equipped with the *Orlicz norm*

$$\|\mathbf{u}\|_{L^\Phi} := \sup \left\{ \int_0^T \mathbf{u} \cdot \mathbf{v} dt | \rho_\Psi(\mathbf{v}) \leq 1 \right\},$$

is a Banach space. By  $\mathbf{u} \cdot \mathbf{v}$  we denote the usual dot product in  $\mathbb{R}^d$  between  $\mathbf{u}$  and  $\mathbf{v}$ . The following alternative expression for the norm, known as *Amemiya norm*, will be useful (see [21, Thm. 10.5] and [18]). For every  $\mathbf{u} \in L^\Phi$ ,

$$\|\mathbf{u}\|_{L^\Phi} = \inf_{k>0} \frac{1}{k} \{1 + \rho_\Phi(k\mathbf{u})\}. \quad (8)$$

The subspace  $E_d^\Phi = E_d^\Phi([0, T])$  is defined as the closure in  $L_d^\Phi$  of the subspace  $L_d^\infty$  of all  $\mathbb{R}^d$ -valued essentially bounded functions. It is shown that  $E_d^\Phi$  is the only one maximal subspace contained in the Orlicz class  $C_d^\Phi$ , i.e.  $\mathbf{u} \in E_d^\Phi$  if and only if  $\rho_\Phi(\lambda \mathbf{u}) < \infty$  for any  $\lambda > 0$ .

A generalized version of *Hölder's inequality* holds in Orlicz spaces. Namely, if  $\mathbf{u} \in L_d^\Phi$  and  $\mathbf{v} \in L_d^\Psi$  then  $\mathbf{u} \cdot \mathbf{v} \in L_1^1$  and

$$\int_0^T \mathbf{v} \cdot \mathbf{u} \, dt \leq \|\mathbf{u}\|_{L^\Phi} \|\mathbf{v}\|_{L^\Psi}. \quad (9)$$

If  $X$  and  $Y$  are Banach spaces, with  $Y \subset X^*$  we denote by  $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{R}$  the bilinear pairing map given by  $\langle x^*, x \rangle = x^*(x)$ . Hölder's inequality shows that  $L_d^\Psi \subset [L_d^\Phi]^*$ , where the pairing  $\langle \mathbf{v}, \mathbf{u} \rangle$  is defined by

$$\langle \mathbf{v}, \mathbf{u} \rangle = \int_0^T \mathbf{v} \cdot \mathbf{u} \, dt \quad (10)$$

with  $\mathbf{u} \in L_d^\Phi$  and  $\mathbf{v} \in L_d^\Psi$ . Unless  $\Phi \in \Delta_2$ , the relation  $L_d^\Psi = [L_d^\Phi]^*$  will not hold. In general, it is true that  $[L_d^\Phi]^* = L_d^\Psi$ .

Like in [21], we will consider the subset  $\Pi(E_d^\Phi, r)$  of  $L_d^\Phi$  given by

$$\Pi(E_d^\Phi, r) := \{\mathbf{u} \in L_d^\Phi \mid d(\mathbf{u}, E_d^\Phi) < r\}.$$

This set is related to the Orlicz class  $C_d^\Phi$  by means of inclusions, namely,

$$\Pi(E_d^\Phi, r) \subset rC_d^\Phi \subset \overline{\Pi(E_d^\Phi, r)} \quad (11)$$

for any positive  $r$ . If  $\Phi \in \Delta_2$ , then the sets  $L_d^\Phi$ ,  $E_d^\Phi$ ,  $\Pi(E_d^\Phi, r)$  and  $C_d^\Phi$  are equal.

We define the *Sobolev-Orlicz space*  $W^1 L_d^\Phi$  (see [3]) by

$$W^1 L_d^\Phi := \{\mathbf{u} \mid \mathbf{u} \text{ is absolutely continuous and } \mathbf{u}, \dot{\mathbf{u}} \in L_d^\Phi\}.$$

$W^1 L_d^\Phi$  is a Banach space when equipped with the norm

$$\|\mathbf{u}\|_{W^1 L^\Phi} = \|\mathbf{u}\|_{L^\Phi} + \|\dot{\mathbf{u}}\|_{L^\Phi}.$$

For a function  $\mathbf{u} \in L_d^1([0, T])$ , we write  $\mathbf{u} = \bar{\mathbf{u}} + \tilde{\mathbf{u}}$  where  $\bar{\mathbf{u}} = \frac{1}{T} \int_0^T \mathbf{u}(t) \, dt$  and  $\tilde{\mathbf{u}} = \mathbf{u} - \bar{\mathbf{u}}$ .

As usual, if  $(X, \|\cdot\|_X)$  is a Banach space and  $(Y, \|\cdot\|_Y)$  is a subspace of  $X$ , we write  $Y \hookrightarrow X$  and we say that  $Y$  is *embedded* in  $X$  when the restricted identity map  $i_Y : Y \rightarrow X$  is bounded. That is, there exists  $C > 0$  such that for any  $y \in Y$  we have  $\|y\|_X \leq C\|y\|_Y$ . With this notation, Hölder's inequality states that  $L_d^\Psi \hookrightarrow [L_d^\Phi]^*$ .

An important aspect of the theory of Sobolev spaces is related to embedding theorems. There is an extensive literature on this question in the Orlicz-Sobolev space setting, see for example [8, 10, 12, 14, 20]. The following simple lemma is essentially known and we will use it systematically. For the sake of completeness, we include a brief proof of it.

**Lemma 2.1.** *Let  $\mathbf{u} \in W^1 L_d^\Phi$ . Then  $\mathbf{u} \in L_d^\infty([0, T])$  and*

$$|\mathbf{u}(t) - \mathbf{u}(s)| \leq \|\dot{\mathbf{u}}\|_{L^\Phi} |t - s| \Phi^{-1} \left( \frac{1}{|t - s|} \right), \quad (12)$$

$$\|\tilde{\mathbf{u}}\|_{L^\infty} \leq T \Phi^{-1} \left( \frac{1}{T} \right) \|\dot{\mathbf{u}}\|_{L^\Phi} \quad (\text{Wirtinger's inequality}), \quad (13)$$

$$\|\mathbf{u}\|_{L^\infty} \leq \Phi^{-1} \left( \frac{1}{T} \right) \max\{1, T\} \|\mathbf{u}\|_{W^1 L^\Phi} \quad (\text{Sobolev's inequality}). \quad (14)$$

*Remark 1.* As usual, a function  $w : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is called a *modulus of continuity* if  $w$  is a continuous increasing function which satisfies  $w(0) = 0$ . We say that  $\mathbf{u} : [0, T] \rightarrow \mathbb{R}^d$  has modulus of continuity  $w$  when there exists a constant  $C > 0$  such that

$$|\mathbf{u}(t) - \mathbf{u}(s)| \leq Cw(|t - s|). \quad (15)$$

The inequality (12) establishes that  $w(s) := s\Phi^{-1}(1/s)$  is a modulus of continuity for all functions  $\mathbf{u} \in W^1L^\Phi$ . Since  $\Phi$  is an  $N$ -function then  $w(0) = 0$ ; and, from the concavity of  $\Phi^{-1}$ , we have that  $w$  is increasing.

*Proof.* For  $0 \leq s \leq t \leq T$ , we get

$$\begin{aligned} |\mathbf{u}(t) - \mathbf{u}(s)| &\leq \int_s^t |\dot{\mathbf{u}}(\tau)| \, d\tau \\ &\leq \|\chi_{[s,t]}\|_{L^\Psi} \|\dot{\mathbf{u}}\|_{L^\Phi} \\ &= \|\dot{\mathbf{u}}\|_{L^\Phi} (t-s) \Phi^{-1}\left(\frac{1}{t-s}\right), \end{aligned} \quad (16)$$

using Hölder's inequality and [21, Eq. (9.11)]. This completes the proof of (12).

**Hasta acá llegamos.**

Since  $u_i$  is continuous, from Mean Value Theorem for integrals, there exists  $s_i \in [0, T]$  such that  $u_i(s_i) = \bar{u}_i$ . Using this  $s_i$  value in (12) with  $u_i$  instead of  $\mathbf{u}$  and taking into account that  $s\Phi^{-1}(1/s)$  is increasing, we obtain Wirtinger's inequality for each  $u_i$ . The inequality (13) follows easily from the corresponding result for each component of  $\mathbf{u}$ .

On the other hand, again by Hölder's inequality and [21, Eq. (9.11)], we have

$$|\bar{\mathbf{u}}| = \frac{1}{T} \int_0^T |\mathbf{u}(s)| \, ds \leq \Phi^{-1}\left(\frac{1}{T}\right) \|\mathbf{u}\|_{L^\Phi}. \quad (17)$$

From (13), (17) and the fact that  $\mathbf{u} = \bar{\mathbf{u}} + \tilde{\mathbf{u}}$ , we obtain (14).  $\square$

*Remark 2.* Inequalities (12) and (13) proves that the embedding  $W^1L_d^\Phi \hookrightarrow C^w([0, T], \mathbb{R}^d)$  holds, where  $C^w([0, T], \mathbb{R}^d)$  denotes the space of generalized  $w$ -Hölder continuous functions. This is the space of all functions satisfying (15) for some  $C > 0$  and  $w(s) = s\Phi^{-1}(1/s)$  and it is a Banach space with norm

$$\|\mathbf{u}\|_{C^w([0, T], \mathbb{R}^d)} := \|\mathbf{u}\|_{L^\infty} + \sup_{t \neq s} \frac{|\mathbf{u}(t) - \mathbf{u}(s)|}{w(|t - s|)}.$$

As it is well known, Arzela-Ascoli Theorem implies that the embedding  $C^w([0, T], \mathbb{R}^d) \hookrightarrow C([0, T], \mathbb{R}^d)$  is compact (see [13, Ch. 5] for the case  $w(s) = |s|^\alpha$ ,  $0 < \alpha \leq 1$  and if  $w$  is arbitrary, the proof follows with some modifications). Therefore we have the following result.

**Corollary 2.2.** *Every bounded sequence  $\{\mathbf{u}_n\}$  in  $W^1L_d^\Phi$  has an uniformly convergent subsequence.*

Given a continuous function  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$ , we define the composition operator  $\mathbf{a} : \mathcal{M}_d \rightarrow \mathcal{M}_d$  by  $\mathbf{a}(\mathbf{u})(t) = a(|\mathbf{u}(t)|)$ . We will often use the following elementary consequence of the previous lemma.

**Corollary 2.3.** *If  $a \in C(\mathbb{R}^+, \mathbb{R}^+)$  then  $\mathbf{a} : W^1 L_d^\Phi \rightarrow L_1^\infty([0, T])$  is bounded. More concretely, there exists a non decreasing function  $A : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\|\mathbf{a}(\mathbf{u})\|_{L^\infty([0, T])} \leq A(\|\mathbf{u}\|_{W^1 L^\Phi})$ .*

*Proof.* Let  $\alpha \in C(\mathbb{R}^+, \mathbb{R}^+)$  be a non-decreasing majorant of  $a$ , for example  $\alpha(s) := \sup_{0 \leq t \leq s} a(t)$ . If  $\mathbf{u} \in W^1 L_d^\Phi$  then, by Lemma 2.1,

$$a(|\mathbf{u}(t)|) \leq \alpha(\|\mathbf{u}\|_{L^\infty}) \leq \alpha\left(\Phi^{-1}\left(\frac{1}{T}\right) \max\{1, T\} \|\mathbf{u}\|_{W^1 L^\Phi}\right) =: A(\|\mathbf{u}\|_{W^1 L^\Phi}).$$

□

The following lemma is an immediate consequence of principles related to operators of Nemitskii type, see [21, §17].

**Lemma 2.4.** *The composition operator  $\varphi$  acts from  $\Pi(E_d^\Phi, 1)$  into  $C_1^\Psi$ .*

*Proof.* As a consequence of [21, Lemma 9.1] we have that  $\varphi(B_{L^\Phi}(0, 1)) \subset C_1^\Psi$ , where  $B_X(\mathbf{u}_0, r)$  is the open ball with center  $\mathbf{u}_0$  and radius  $r > 0$  in the space  $X$ . Therefore, applying [21, Lemma 17.1], we deduce that  $\varphi$  acts from  $\Pi(E_d^\Phi, 1)$  into  $C_1^\Psi$ . □

We also need the following technical lemma.

**Lemma 2.5.** *Let  $\lambda > 0$  and let  $\{\mathbf{u}_n\}_{n \in \mathbb{N}}$  be a sequence of functions in  $\Pi(E_d^\Phi, \lambda)$  converging to  $\mathbf{u} \in \Pi(E_d^\Phi, \lambda)$  in the  $L^\Phi$ -norm. Then, there exist a subsequence  $\mathbf{u}_{n_k}$  and a real valued function  $h \in \Pi(E_1^\Phi([0, T]), \lambda)$  such that  $\mathbf{u}_{n_k} \rightarrow \mathbf{u}$  a.e. and  $|\mathbf{u}_{n_k}| \leq h$  a.e.*

*Proof.* Let  $r := d(\mathbf{u}, E_d^\Phi)$ ,  $r < \lambda$ . As  $\mathbf{u}_n$  converges to  $\mathbf{u}$ , there exists a subsequence  $(n_k)$  such that

$$\|\mathbf{u}_{n_k} - \mathbf{u}\|_{L^\Phi} < \frac{\lambda - r}{2} \quad \text{and} \quad \|\mathbf{u}_{n_k} - \mathbf{u}_{n_{k+1}}\|_{L^\Phi} < 2^{-(k+1)}(\lambda - r).$$

Let  $h : [0, T] \rightarrow \mathbb{R}$  defined by

$$h(x) = |\mathbf{u}_{n_1}(x)| + \sum_{k=2}^{\infty} |\mathbf{u}_{n_k}(x) - \mathbf{u}_{n_{k-1}}(x)|. \quad (18)$$

As a consequence of [21, Lemma 10.1] we have that  $d(\mathbf{v}, E_d^\Phi) = d(|\mathbf{v}|, E_1^\Phi)$  for any  $\mathbf{v} \in L_d^\Phi$ . Now

$$d(|\mathbf{u}_{n_1}|, E_1^\Phi) = d(\mathbf{u}_{n_1}, E_d^\Phi) \leq d(\mathbf{u}_{n_1}, \mathbf{u}) + d(\mathbf{u}, E_d^\Phi) < \frac{\lambda + r}{2}.$$

Then

$$d(h, E_1^\Phi) \leq d(h, |\mathbf{u}_{n_1}|) + d(|\mathbf{u}_{n_1}|, E_1^\Phi) < \lambda.$$

Therefore,  $h \in \Pi(E_1^\Phi, \lambda)$  and  $|h| < \infty$  a.e. We conclude that the series  $\mathbf{u}_{n_1}(x) + \sum_{k=2}^\infty (\mathbf{u}_{n_k}(x) - \mathbf{u}_{n_{k-1}}(x))$  is absolutely convergent a.e. and this fact implies that  $\mathbf{u}_{n_k} \rightarrow \mathbf{u}$  a.e. The inequality  $|\mathbf{u}_{n_k}| \leq h$  follows straightforwardly from the definition of  $h$ .  $\square$

A common obstacle with Orlicz spaces, that distinguishes it from  $L^p$  spaces, is that a sequence  $\mathbf{u}_n \in L_d^\Phi$  which is uniformly bounded by  $h \in L_1^\Phi$  and a.e. convergent to  $\mathbf{u}$  is not necessarily norm convergent. Fortunately, the subspace  $E_d^\Phi$  has that property.

**Lemma 2.6.** *Suppose that  $\mathbf{u}_n \in L_d^\Phi$  is a sequence such that  $\mathbf{u}_n \rightarrow \mathbf{u}$  a.e. and assume that there exist  $h \in E_1^\Phi$  with  $|\mathbf{u}_n| \leq h$  a.e. then  $\|\mathbf{u}_n - \mathbf{u}\|_{L^\Phi} \rightarrow 0$ .*

*Proof.* [26, p. 84] and [21, Thm. 10.3].  $\square$

We recall some useful concepts.

**Definition 2.7.** *Given a function  $I : U \rightarrow \mathbb{R}$  where  $U$  is an open set of a Banach space  $X$ , we say that  $I$  has a Gâteaux derivative at  $\mathbf{u} \in U$  if there exists  $\mathbf{u}^* \in X^*$  such that for every  $\mathbf{v} \in X$*

$$\lim_{s \rightarrow 0} \frac{I(\mathbf{u} + s\mathbf{v}) - I(\mathbf{u})}{s} = \langle \mathbf{u}^*, \mathbf{v} \rangle.$$

*See [4] for details.*

**Definition 2.8.** *Let  $X$  be a Banach space and let  $D \subset X$ . A non linear operator  $T : D \rightarrow X^*$  is called demicontinuous if it is continuous when  $X$  is equipped with the strong topology and  $X^*$  with the weak\* topology (see [19]).*

### 3 Differentiability of action integrals in Orlicz spaces

**Definition 3.1.** *We say that a function  $\mathcal{L} : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function if for fixed  $(\mathbf{x}, \mathbf{y})$  the map  $t \mapsto \mathcal{L}(t, \mathbf{x}, \mathbf{y})$  is measurable and for fixed  $t$  the map  $(\mathbf{x}, \mathbf{y}) \mapsto \mathcal{L}(t, \mathbf{x}, \mathbf{y})$  is continuously differentiable for almost everywhere  $t \in [0, T]$ .*

Comentamos el Remark 3 porque pensamos que debería escribirse en el texto, como lo hicimos debajo

In the following comments we discuss the relevance of the function  $f$  in the inequalities (2), (3) and (4). These conditions are a direct generalization of the conditions [25, Eq (a), p. 10]. In particular, we are interested in seeing when for every  $f \in E_1^\Phi$  there exist  $b \in L_1^1$  o  $\tilde{b}$ , para que no coincida con la ecuacion (2) y (3) and a constant  $C > 0$  such that

$$\Phi(s + f(t)) \leq C\Phi(s) + b(t) \quad \text{for every } s > 0. \quad (19)$$

If (19) is true then in the equations (2) and (3) we can suppose  $f = 0$ . The same considerations should be done with  $\varphi(s + f(t))$ .



*Remark 3.* As a direct consequence of convexity, we can bound the term  $\Phi(s+f(t))$  by the expression  $\frac{1}{2}\Phi(2s) + b(t)$  where  $b(t) := \frac{1}{2}\Phi(2f(t)) \in L_1^1$  and  $f \in E_1^\Phi$ . Therefore, we can always assume  $f = 0$  in (2) and (3) at the price of making smaller the value of  $\lambda$ .

*Remark 4.* If  $\Phi \in \Delta_2$  then we can assume  $f = 0$  keeping the same value of  $\lambda$ . This is a consequence of the fact that a non decreasing function  $G : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  satisfying the  $\Delta_2$ -condition, is a quasi-subadditive function (see [1, Prop. 4.2]) i.e. there exists  $K > 0$  such that

$$G(s_1 + s_2) \leq K(G(s_1) + G(s_2)).$$

Moreover, if  $\Phi \in \Delta_2$  then  $\varphi \in \Delta_2$  (see [16, Eq. (2.15)]). Therefore if  $\Phi \in \Delta_2$  we have that

$$\Phi(s + f(t)) \leq K\Phi(s) + K\Phi(f(t)) = K\Phi(s) + b_1(t),$$

where  $b_1(t) = b(t) + K\Phi(f(t)) \in L_1^1([0, T])$ . A similar fact holds with  $\varphi$  instead of  $\Phi$ , namely

$$\varphi(s + f(t)) \leq c_1(t) + \varphi(s),$$

where  $c_1(t) := K\varphi(f(t)) \in L_1^\Psi$ , as a consequence of Lemma 2.4 and the  $\Delta_2$ -condition on  $\Phi$ .

*Remark 5.* If  $\Phi \notin \Delta_2$ , then (19) may or may not be true. For example, if we consider the  $N$ -function  $\Phi(s) = e^s - s - 1$  which does not satisfy the  $\Delta_2$ -condition, we have that  $E_1^\Phi = L_1^\infty$ . In fact, if  $f \in E_1^\Phi$  then, since  $1/2e^s \leq \Phi(s) + 1$  and  $pf \in C_1^\Phi$  for every  $p > 0$ , we get  $\int_0^T e^{pf(t)} dt < \infty$  for every  $p > 0$ . This implies that  $f \in L_1^\infty$ . Therefore,

$$\Phi(s + f(t)) \leq e^{s+f(t)} \leq 2\|e^f\|_{L^\infty} \Phi(s) + 2\|e^f\|_{L^\infty}.$$

On the other hand, we consider the  $N$ -function  $\Phi(s) = e^{s^2} - 1$ . Suppose that  $0 \leq f \in E_1^\Phi$  and  $b \in L_1^1$  satisfy (19), then

$$e^{s^2} e^{2sf(t)} \leq \Phi(s + f(t)) + 1 \leq Ce^{s^2} + b(t).$$

Dividing by  $e^{s^2}$  and taking the limit as  $s \rightarrow \infty$ , we obtain that  $f = 0$  a.e. In other words, if  $f \neq 0$  on a set with positive measure, the estimations in (2) and (3) are essentially bigger than a bound of the type  $a(|x|)(b(t) + \Phi(|y|/\lambda))$ .

**Theorem 3.2.** *Let  $\mathcal{L}$  be a Carathéodory function satisfying (2), (3) and (4). Then the following statements hold:*

1. *The action integral given by (5) is finitely defined in  $\mathcal{E}_d^\Phi(\lambda) := W^1 L_d^\Phi \cap \{u | \dot{u} \in \Pi(E_d^\Phi, \lambda)\}$ .*
2. *The function  $I$  is Gâteaux differentiable on  $\mathcal{E}_d^\Phi(\lambda)$  and its derivative  $I'$  is demi-continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $[W^1 L_d^\Phi]^*$ . Moreover,  $I'$  is given by the following expression*

$$\langle I'(u), v \rangle = \int_0^T \{D_x \mathcal{L}(t, u, \dot{u}) \cdot v + D_y \mathcal{L}(t, u, \dot{u}) \cdot \dot{v}\} dt. \quad (20)$$

3. If  $\Psi \in \Delta_2$  then  $I'$  is continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $[W^1 L_d^\Phi]^*$  when both spaces are equipped with the strong topology.

*Proof.* Since  $\lambda \Pi(E_d^\Phi, r) = \Pi(E_d^\Phi, \lambda r)$ , we have  $\dot{\mathbf{u}}/\lambda \in \Pi(E_d^\Phi, 1)$ . Thus, as  $f \in E_1^\Phi$  and attending to (11), we get

$$|\dot{\mathbf{u}}/\lambda + f| \in \Pi(E_1^\Phi, 1) \subset C_1^\Phi. \quad (21)$$

By Corollary 2.3, we get

$$|\mathcal{L}(\cdot, \mathbf{u}, \dot{\mathbf{u}})| \leq A(\|\mathbf{u}\|_{W^1 L^\Phi}) \left( b + \Phi \left( \frac{|\dot{\mathbf{u}}|}{\lambda} + f \right) \right) \in L_1^1.$$

This fact proves item 1.

We split the proof of item 2 in four steps.

*Step 1.* The non linear operator  $\mathbf{u} \mapsto D_{\mathbf{x}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$  is continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $L_d^1([0, T])$  with the strong topology on both sets.

If  $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$ , from (3) and (21), we obtain

$$|D_{\mathbf{x}} \mathcal{L}(\cdot, \mathbf{u}, \dot{\mathbf{u}})| \leq A(\|\mathbf{u}\|_{W^1 L^\Phi}) \left( b + \Phi \left( \frac{|\dot{\mathbf{u}}|}{\lambda} + f \right) \right) \in L_1^1. \quad (22)$$

Let  $\{\mathbf{u}_n\}_{n \in \mathbb{N}}$  be a sequence of functions in  $\mathcal{E}_d^\Phi(\lambda)$ , and  $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$  such that  $\mathbf{u}_n \rightarrow \mathbf{u}$  in  $W^1 L_d^\Phi$ . Then  $\mathbf{u}_n \rightarrow \mathbf{u}$  in  $L_d^\Phi$  and  $\dot{\mathbf{u}}_n \rightarrow \dot{\mathbf{u}}$  in  $L_d^\Phi$ . By Lemma 2.5 there exist a subsequence  $\mathbf{u}_{n_k}$  and a function  $h \in \Pi(E_1^\Phi, \lambda)$  such that  $\mathbf{u}_{n_k} \rightarrow \mathbf{u}$  a.e.,  $\dot{\mathbf{u}}_{n_k} \rightarrow \dot{\mathbf{u}}$  a.e. and  $|\dot{\mathbf{u}}_{n_k}| \leq h$  a.e. Since  $\mathbf{u}_{n_k}, k = 1, 2, \dots$ , is a strong convergent sequence in  $W^1 L_d^\Phi$ , it is a bounded sequence in  $W^1 L_d^\Phi$ . According to Lemma 2.1 and Corollary 2.3, there exists  $M > 0$  such that  $\|\mathbf{a}(\mathbf{u}_{n_k})\|_{L^\infty} \leq M, k = 1, 2, \dots$ . From the previous facts and (22), we get

$$|D_{\mathbf{x}} \mathcal{L}(\cdot, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k})| \leq M \left( b + \Phi \left( \frac{|h|}{\lambda} + f \right) \right) \in L_1^1.$$

On the other hand, by the Carathéodory condition, we have

$$D_{\mathbf{x}} \mathcal{L}(t, \mathbf{u}_{n_k}(t), \dot{\mathbf{u}}_{n_k}(t)) \rightarrow D_{\mathbf{x}} \mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t)) \quad \text{for a.e. } t \in [0, T].$$

Applying the Dominated Convergence Theorem we conclude the proof of step 1.

*Step 2.* The non linear operator  $\mathbf{u} \mapsto D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$  is continuous from  $\mathcal{E}_d^\Phi(\lambda)$  with the strong topology into  $[L_d^\Phi]^*$  with the weak\* topology.

Let  $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$ . From (21), Lemma 2.4 and Corollary 2.3, it follows that

$$\varphi \left( \frac{|\dot{\mathbf{u}}|}{\lambda} + f \right) \in C_1^\Psi \quad (23)$$

and  $\mathbf{a}(|\mathbf{u}|) \in L_1^\infty$ . Therefore, in virtue of (4) we get

$$|D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}, \dot{\mathbf{u}})| \leq A(\|\mathbf{u}\|_{W^1 L^\Phi}) \left( c + \varphi \left( \frac{|\dot{\mathbf{u}}|}{\lambda} + f \right) \right) \in L_1^\Psi. \quad (24)$$

We note that (22), (24), the imbeddings  $W^1 L_d^\Phi \hookrightarrow L_d^\infty$  and  $L_d^\Psi \hookrightarrow [L_d^\Phi]^*$  imply that the second member of (20) defines an element in  $[W^1 L_d^\Phi]^*$ .

Let  $\mathbf{u}_n, \mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$  such that  $\mathbf{u}_n \rightarrow \mathbf{u}$  in the norm of  $W^1 L_d^\Phi$ . We must prove that  $D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}_n, \dot{\mathbf{u}}_n) \xrightarrow{w^*} D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}, \dot{\mathbf{u}})$ . On the contrary, if there exist  $\mathbf{v} \in L_d^\Phi$ ,  $\epsilon > 0$  and a subsequence of  $\{\mathbf{u}_n\}$  (denoted  $\{\mathbf{u}_n\}$  for simplicity) such that

$$|\langle D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}_n, \dot{\mathbf{u}}_n), \mathbf{v} \rangle - \langle D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}, \dot{\mathbf{u}}), \mathbf{v} \rangle| \geq \epsilon. \quad (25)$$

We have  $\mathbf{u}_n \rightarrow \mathbf{u}$  in  $L_d^\Phi$  and  $\dot{\mathbf{u}}_n \rightarrow \dot{\mathbf{u}}$  in  $L_d^\Phi$ . By Lemma 2.5, there exist a subsequence  $\mathbf{u}_{n_k}$  and a function  $h \in \Pi(E_1^\Phi, \lambda)$  such that  $\mathbf{u}_{n_k} \rightarrow \mathbf{u}$  a.e.,  $\dot{\mathbf{u}}_{n_k} \rightarrow \dot{\mathbf{u}}$  a.e. and  $|\dot{\mathbf{u}}_{n_k}| \leq h$  a.e. As in the previous step, since  $\mathbf{u}_n$  is a convergent sequence, the Corollary 2.3 implies that  $a(|\mathbf{u}_n(t)|)$  is uniformly bounded by a certain constant  $M > 0$ . Therefore, with  $\mathbf{u}_n$  instead of  $\mathbf{u}$ , inequality (24) becomes

$$|D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}_n, \dot{\mathbf{u}}_n)| \leq M \left( c + \varphi \left( \frac{h}{\lambda} + f \right) \right) \in L_1^\Psi. \quad (26)$$

Consequently, as  $\mathbf{v} \in L_d^\Phi$  and employing Hölder's inequality, we obtain that  $\sup_n |D_{\mathbf{y}} \mathcal{L}(\cdot, \mathbf{u}_n, \dot{\mathbf{u}}_n) \cdot \mathbf{v}| \in L_1^1$ . Finally, from the Lebesgue Dominated Convergence Theorem, we deduce

$$\int_0^T D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k}) \cdot \mathbf{v} \, dt \rightarrow \int_0^T D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} \, dt \quad (27)$$

which contradicts the inequality (25). This completes the proof of step 2.

*Step 3.* We will prove (20). The proof follows similar lines that [25, Thm. 1.4]. For  $\mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$  and  $\mathbf{0} \neq \mathbf{v} \in W^1 L_d^\Phi$ , we define the function

$$H(s, t) := \mathcal{L}(t, \mathbf{u}(t) + s\mathbf{v}(t), \dot{\mathbf{u}}(t) + s\dot{\mathbf{v}}(t)).$$

From [21, Thm. 10.1] we obtain that if  $|\mathbf{u}| \leq |\mathbf{v}|$  then  $d(\mathbf{u}, E_d^\Phi) \leq d(\mathbf{v}, E_d^\Phi)$ . Therefore, for  $|s| \leq s_0 := (\lambda - d(\dot{\mathbf{u}}, E_d^\Phi)) / \|\mathbf{v}\|_{W^1 L^\Phi}$  we have

$$d(\dot{\mathbf{u}} + s\dot{\mathbf{v}}, E_d^\Phi) \leq d(|\dot{\mathbf{u}}| + s|\dot{\mathbf{v}}|, E_1^\Phi) \leq d(|\dot{\mathbf{u}}|, E_1^\Phi) + s\|\dot{\mathbf{v}}\|_{L^\Phi} < \lambda.$$

As a consequence  $\dot{\mathbf{u}} + s\dot{\mathbf{v}} \in \Pi(E_d^\Phi, \lambda)$  and  $|\dot{\mathbf{u}}| + s|\dot{\mathbf{v}}| \in \Pi(E_1^\Phi, \lambda)$ . These facts imply, in virtue of Theorem 3.2 item 1, that  $I(\mathbf{u} + s\mathbf{v})$  is well defined and finite for  $|s| \leq s_0$ . Using Corollary 2.3 we see that

$$\|a(|\mathbf{u} + s\mathbf{v}|)\|_{L^\infty} \leq A(\|\mathbf{u} + s\mathbf{v}\|_{W^1 L^\Phi}) \leq A(\|\mathbf{u}\|_{W^1 L^\Phi} + s_0\|\mathbf{v}\|_{W^1 L^\Phi}) =: M$$

Now, applying Chain Rule, (22), (24) the monotonicity of  $\varphi$  and  $\Phi$ , the fact that  $\mathbf{v} \in L_d^\infty$  and  $\dot{\mathbf{v}} \in L_d^\Phi$  and Hölder's inequality, we have

$$\begin{aligned} |D_s H(s, t)| &= |D_{\mathbf{x}} \mathcal{L}(t, \mathbf{u} + s\mathbf{v}, \dot{\mathbf{u}} + s\dot{\mathbf{v}}) \cdot \mathbf{v} + D_{\mathbf{y}} \mathcal{L}(t, \mathbf{u} + s\mathbf{v}, \dot{\mathbf{u}} + s\dot{\mathbf{v}}) \cdot \dot{\mathbf{v}}| \\ &\leq M \left[ \left( b(t) + \Phi \left( \frac{|\dot{\mathbf{u}}| + s_0|\dot{\mathbf{v}}|}{\lambda} + f(t) \right) \right) |\mathbf{v}| \right. \\ &\quad \left. + \left( c(t) + \varphi \left( \frac{|\dot{\mathbf{u}}| + s_0|\dot{\mathbf{v}}|}{\lambda} + f(t) \right) \right) |\dot{\mathbf{v}}| \right] \in L_1^1. \end{aligned} \quad (28)$$

Consequently,  $I$  has a directional derivative and

$$\langle I'(\mathbf{u}), \mathbf{v} \rangle = \frac{d}{ds} I(\mathbf{u} + s\mathbf{v})|_{s=0} = \int_0^T \{D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} + D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \dot{\mathbf{v}}\} dt.$$

Moreover, from (22), (24), Lemma 2.1 and previous formula, we have

$$|\langle I'(\mathbf{u}), \mathbf{v} \rangle| \leq \|D_{\mathbf{x}}\mathcal{L}\|_{L^1} \|\mathbf{v}\|_{L^\infty} + \|D_{\mathbf{y}}\mathcal{L}\|_{L^\Psi} \|\dot{\mathbf{v}}\|_{L^\Phi} \leq C \|\mathbf{v}\|_{W^1 L^\Phi}$$

with a appropriate constant  $C$ . This completes the proof of the Gâteaux differentiability of  $I$ .

**Step 4. (agregamos este paso)** The operator  $I' : \mathcal{E}_d^\Phi(\lambda) \rightarrow [W^1 L_d^\Phi]^*$  is demicontinuous. This is a consequence of the continuity of the mappings  $\mathbf{u} \mapsto D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$  and  $\mathbf{u} \mapsto D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$ . Indeed, if  $\mathbf{u}_n, \mathbf{u} \in \mathcal{E}_d^\Phi(\lambda)$  with  $\mathbf{u}_n \rightarrow \mathbf{u}$  in the norm of  $W^1 L_d^\Phi$  and  $\mathbf{v} \in W^1 L_d^\Phi$ , then

$$\begin{aligned} \langle I'(\mathbf{u}_n), \mathbf{v} \rangle &= \int_0^T \{D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}_n, \dot{\mathbf{u}}_n) \cdot \mathbf{v} + D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}_n, \dot{\mathbf{u}}_n) \cdot \dot{\mathbf{v}}\} dt \\ &\rightarrow \int_0^T \{D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \mathbf{v} + D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}}) \cdot \dot{\mathbf{v}}\} dt \\ &= \langle I'(\mathbf{u}), \mathbf{v} \rangle. \end{aligned}$$

In order to prove item 3, it is necessary to see that the maps  $\mathbf{u} \mapsto D_{\mathbf{x}}\mathcal{L}(\cdot, \mathbf{u}(\cdot), \dot{\mathbf{u}}(\cdot))$  and  $\mathbf{u} \mapsto D_{\mathbf{y}}\mathcal{L}(\cdot, \mathbf{u}(\cdot), \dot{\mathbf{u}}(\cdot))$  are norm continuous from  $\mathcal{E}_d^\Phi(\lambda)$  into  $L_d^1$  and  $L_d^\Psi$  respectively. **Debemos unificar la notacion respecto a la usada en el item 2  $\mathbf{u} \mapsto D_{\mathbf{x}}\mathcal{L}(\cdot, \mathbf{u}(\cdot), \dot{\mathbf{u}}(\cdot))$  o  $\mathbf{u} \mapsto D_{\mathbf{x}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$  ?** The continuity of the first map has already been proved in step 1. We will prove the continuity of the second map developing a similar argument to the one given in step 2 of item 2. We consider  $\mathbf{u}_n$  and  $\mathbf{u}$  in  $\mathcal{E}_d^\Phi(\lambda)$  with  $\|\mathbf{u}_n - \mathbf{u}\|_{W^1 L^\Phi} \rightarrow 0$ . By Lemma 2.5, there exist a subsequence  $\mathbf{u}_{n_k} \in \mathcal{E}_d^\Phi(\lambda)$  and a function  $h \in \Pi(E_1^\Phi, \lambda)$  such that  $\mathbf{u}_{n_k} \rightarrow \mathbf{u}$  a.e.,  $\dot{\mathbf{u}}_{n_k} \rightarrow \dot{\mathbf{u}}$  a.e. and  $|\dot{\mathbf{u}}_{n_k}| \leq h$  a.e. Then, since  $\mathcal{L}$  is a Carathéodory function we have  $D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}_{n_k}(t), \dot{\mathbf{u}}_{n_k}(t)) \rightarrow D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}(t), \dot{\mathbf{u}}(t))$  a.e.  $t \in [0, T]$ . By (4) and the fact that  $\Psi \in \Delta_2$ , we obtain

$$\begin{aligned} |D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}_{n_k}(t), \dot{\mathbf{u}}_{n_k}(t))| &\leq a(|\mathbf{u}_{n_k}(t)|) \left( c(t) + \varphi \left( \frac{|\dot{\mathbf{u}}_{n_k}(t)|}{\lambda} + f(t) \right) \right) \\ &\leq C \left( c(t) + \varphi \left( \frac{|h(t)|}{\lambda} + f(t) \right) \right) \in L_1^\Psi = E_1^\Psi. \end{aligned}$$

Therefore, invoking Lemma 2.6, we have proved that from any sequence  $\mathbf{u}_n$  which converges to  $\mathbf{u}$  in  $W^1 L_d^\Phi$  we can extract a subsequence such that  $D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}_{n_k}, \dot{\mathbf{u}}_{n_k}) \rightarrow D_{\mathbf{y}}\mathcal{L}(t, \mathbf{u}, \dot{\mathbf{u}})$  in the strong topology. The desired result is obtained by a standard argument.

The continuity of  $I'$  follows from the continuity of  $D_{\mathbf{x}}\mathcal{L}$  and  $D_{\mathbf{y}}\mathcal{L}$  by using the formula (20).  $\square$

## 4 Critical points and Euler-Lagrange equations

In this section we derive the Euler-Lagrange equations associated to critical points of action integrals. We denote by  $W^1 L_T^\Phi$  the subspace of  $W^1 L_d^\Phi$  containing all  $T$ -periodic functions. As usual, when  $Y$  is a subspace of the Banach space  $X$ , we denote by  $Y^\perp$  the *annihilator subspace* of  $X^*$ , i.e. the subspace that consists of all bounded linear functions which are identically zero on  $Y$ .

We recall that a function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is called *strictly convex* if  $f\left(\frac{x+y}{2}\right) < \frac{1}{2}(f(x) + f(y))$  for  $x \neq y$ . It is well known that if  $f$  is a strictly convex and differentiable function, then  $D_x f : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a one-to-one map (see, for instance [28, Thm. 12.17]).

**Theorem 4.1.** *Let  $u \in \mathcal{E}_d^\Phi(\lambda)$ . The following statements are equivalent:*

1.  $I'(u) \in (W^1 L_T^\Phi)^\perp$ .
2.  $D_y \mathcal{L}(t, u(t), \dot{u}(t))$  is an absolutely continuous function and  $u$  solves the following boundary value problem

$$\begin{cases} \frac{d}{dt} D_y \mathcal{L}(t, u(t), \dot{u}(t)) = D_x \mathcal{L}(t, u(t), \dot{u}(t)) & \text{a.e. } t \in (0, T) \\ u(0) - u(T) = D_y \mathcal{L}(0, u(0), \dot{u}(0)) - D_y \mathcal{L}(T, u(T), \dot{u}(T)) = 0. \end{cases} \quad (29)$$

Moreover if  $D_y \mathcal{L}(t, x, y)$  is  $T$ -periodic with respect to the variable  $t$  and strictly convex with respect to  $y$ , then  $D_y \mathcal{L}(0, u(0), \dot{u}(0)) - D_y \mathcal{L}(T, u(T), \dot{u}(T)) = 0$  is equivalent to  $\dot{u}(0) = \dot{u}(T)$ .

*Proof.* The condition  $I'(u) \in (W^1 L_T^\Phi)^\perp$  and (20) imply

$$\int_0^T D_y \mathcal{L}(t, u(t), \dot{u}(t)) \cdot \dot{v}(t) dt = - \int_0^T D_x \mathcal{L}(t, u(t), \dot{u}(t)) \cdot v(t) dt.$$

Using [25, pp. 6-7] we obtain that  $D_y \mathcal{L}(t, u(t), \dot{u}(t))$  is absolutely continuous and  $T$ -periodic, therefore it is differentiable a.e. on  $[0, T]$  and the first equality of (29) holds true. This completes the proof of 1 implies 2. The proof of 2 implies 1 follows easily from (20) and (29).

The last part of the theorem is a consequence of that  $D_y \mathcal{L}(T, u(T), \dot{u}(T)) = D_y \mathcal{L}(0, u(0), \dot{u}(0)) = D_y \mathcal{L}(T, u(T), \dot{u}(0))$  and the injectivity of  $D_y \mathcal{L}(T, u(T), \cdot)$ .  $\square$

## 5 Coercivity discussion

We recall the following usual definition in the context of calculus of variations.

**Definition 5.1.** *Let  $X$  be a Banach space and let  $D$  be an unbounded subset of  $X$ . Suppose  $J : D \subset X \rightarrow \mathbb{R}$ . We say that  $J$  is coercive if  $J(u) \rightarrow +\infty$  when  $\|u\|_X \rightarrow +\infty$ .*

It is well known that coercivity is a useful ingredient in order to establish existence of minima. Therefore, we are interested in finding conditions which ensure the coercivity of the action integral  $I$  acting on  $\mathcal{E}_d^\Phi(\lambda)$ . For this purpose, we need to introduce the following extra condition on lagrangian function  $\mathcal{L}$

$$\mathcal{L}(t, \mathbf{x}, \mathbf{y}) \geq \alpha_0 \Phi\left(\frac{|\mathbf{y}|}{\Lambda}\right) + F(t, \mathbf{x}), \quad (30)$$

where  $\alpha_0, \Lambda > 0$  and  $F : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a Carathéodory function, i.e.  $F(t, \mathbf{x})$  is measurable with respect to  $t$  for every fixed  $\mathbf{x} \in \mathbb{R}^d$  and it is continuous at  $\mathbf{x}$  for a.e.  $t \in [0, T]$ . We note that, in virtue of (30) and (2), we have  $F(t, \mathbf{x}) \leq a(|\mathbf{x}|)b_0(t)$  with  $b_0(t) := b(t) + \Phi(f(t)) \in L_1^1([0, T])$ . In order to ensure that integral  $\int_0^T F(t, \mathbf{u}) dt$  is finite for  $\mathbf{u} \in W^1 L^\Phi$ , we need to assume

$$|F(t, \mathbf{x})| \leq a(|\mathbf{x}|)b_0(t), \quad \text{for a.e. } t \in [0, T] \quad \text{and for every } \mathbf{x} \in \mathbb{R}^d. \quad (31)$$

As we shall see in Theorem 5.3, when  $\mathcal{L}$  satisfies (2), (3), (4), (30) and (31), the coercivity of the action integral  $I$  is related to the coercivity of the functional

$$J_{C,\nu}(\mathbf{u}) := \rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right) - C\|\mathbf{u}\|_{L^\Phi}^\nu, \quad (32)$$

for  $C, \nu > 0$ . If  $\Phi(x) = |x|^p/p$  then  $J_{C,\nu}$  is clearly coercive for  $\nu < p$ . For more general  $\Phi$  the situation is more interesting as it will be shown in the following lemma.

**Lemma 5.2.** *Let  $\Phi$  and  $\Psi$  be complementary  $N$ -functions. Then:*

1. *If  $C\Lambda < 1$  then  $J_{C,1}$  is coercive.*
2. *If  $\Psi \in \Delta_2$  globally, then there exists a constant  $\alpha_\Phi > 1$  such that, for any  $0 < \mu < \alpha_\Phi$ ,*

$$\lim_{\|\mathbf{u}\|_{L^\Phi} \rightarrow \infty} \frac{\rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right)}{\|\mathbf{u}\|_{L^\Phi}^\mu} = +\infty. \quad (33)$$

*In particular, the functional  $J_{C,\mu}$  is coercive for every  $C > 0$  and  $0 < \mu < \alpha_\Phi$ . The constant  $\alpha_\Phi$  is one of the so-called Matuszewska-Orlicz indices (see [22, Ch. 11]).*

3. *If  $J_{C,1}$  is coercive with  $C\Lambda > 1$ , then  $\Psi \in \Delta_2$ .*

*Proof.* By (8) we have

$$(1 - C\Lambda)\|\mathbf{u}\|_{L^\Phi} + C\Lambda\|\mathbf{u}\|_{L^\Phi} = \|\mathbf{u}\|_{L^\Phi} \leq \Lambda + \Lambda\rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right),$$

then

$$\frac{(1 - C\Lambda)}{\Lambda}\|\mathbf{u}\|_{L^\Phi} - 1 \leq \rho_\Phi\left(\frac{\mathbf{u}}{\Lambda}\right) - C\|\mathbf{u}\|_{L^\Phi} = J_{C,1}(\mathbf{u}).$$

This shows that  $J_{C,1}$  is coercive and therefore item 1 is proved.

In virtue of [2, Eq. (2.8)], the  $\Delta_2$ -condition on  $\Psi$ , [22, Thm. 11.7] and [22, Cor. 11.6], we obtain constants  $K > 0$  and  $\alpha_\Phi > 1$  such that

$$\Phi(rs) \geq Kr^\nu \Phi(s) \quad (34)$$

for any  $0 < \nu < \alpha_\Phi$ ,  $s \geq 0$  and  $r > 1$ .

Let  $1 < \mu < \alpha_\Phi$  and let  $r > \Lambda$  be a constant that will be specified later. Then, from (34) and (8), we get

$$\begin{aligned} \frac{\int_0^T \Phi\left(\frac{|u|}{\Lambda}\right) dt}{\|u\|_{L^\Phi}^\mu} &\geq K \left(\frac{r}{\Lambda}\right)^\nu \frac{\int_0^T \Phi(r^{-1}|u|) dt}{\|u\|_{L^\Phi}^\mu} \\ &\geq K \left(\frac{r}{\Lambda}\right)^\nu \frac{r^{-1}\|u\|_{L^\Phi} - 1}{\|u\|_{L^\Phi}^\mu}. \end{aligned}$$

We choose  $r = \|u\|_{L^\Phi}/2$ . Since  $\|u\|_{L^\Phi} \rightarrow +\infty$  we can assume  $\|u\|_{L^\Phi} > 2\Lambda$ . Thus  $r > \Lambda$  and

$$\frac{\int_0^T \Phi\left(\frac{|u|}{\Lambda}\right) dt}{\|u\|_{L^\Phi}^\mu} \geq \frac{K}{2^\nu \Lambda^\nu} \|u\|_{L^\Phi}^{\nu-\mu} \rightarrow +\infty \quad \text{as } \|u\|_{L^\Phi} \rightarrow +\infty,$$

because  $\nu > \mu$ .

In order to prove the last item, we assume that  $\Psi \notin \Delta_2$ . By [21, Thm. 4.1], there exists a sequence of real numbers  $r_n$  such that  $r_n \rightarrow \infty$  and

$$\lim_{n \rightarrow \infty} \frac{r_n \psi(r_n)}{\Psi(r_n)} = +\infty. \quad (35)$$

Now, we choose  $u_n$  such that  $|u_n| = \Lambda \psi(r_n) \chi_{[0, \frac{1}{\Psi(r_n)}]}$ . Then, by [21, Eq. (9.11)], we get

$$\|u_n\|_{L^\Phi} = \Lambda \frac{\psi(r_n)}{\Psi(r_n)} \Psi^{-1}(\Psi(r_n)) = \Lambda \frac{r_n \psi(r_n)}{\Psi(r_n)} \rightarrow \infty, \quad \text{as } n \rightarrow \infty.$$

Next, using Young's equality (see [21, Eq. (2.7)]), we have

$$\begin{aligned} J_{C,1}(u_n) &= \int_0^T \Phi\left(\frac{|u_n|}{\Lambda}\right) dt - C \|u_n\|_{L^\Phi} \\ &= \frac{1}{\Psi(r_n)} [\Phi(\psi(r_n)) - C \Lambda r_n \psi(r_n)] \\ &= \frac{1}{\Psi(r_n)} [r_n \psi(r_n) - \Psi(r_n) - C \Lambda r_n \psi(r_n)] \\ &= \frac{(1 - C \Lambda) r_n \psi(r_n)}{\Psi(r_n)} - 1. \end{aligned}$$

From (35) and the condition  $C\Lambda > 1$ , we obtain  $J_{C,1}(u_n) \rightarrow -\infty$ , which contradicts the coercivity of  $J_{C,1}$ .  $\square$

Next, we present two theorems that establish coercivity of action integrals.

**Theorem 5.3.** *Let  $\mathcal{L}$  be a Lagrangian function satisfying (2), (3), (4), (30) and (31). We assume the following conditions:*

1. *There exist a non negative function  $b_1 \in L^1_1$  and a constant  $\mu > 0$  such that for any  $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^d$  and a.e.  $t \in [0, T]$*

$$|F(t, \mathbf{x}_2) - F(t, \mathbf{x}_1)| \leq b_1(t)(1 + |\mathbf{x}_2 - \mathbf{x}_1|^\mu). \quad (36)$$

*We suppose that  $\mu < \alpha_\Phi$ , with  $\alpha_\Phi$  as in Lemma 5.2, in the case that  $\Psi \in \Delta_2$ ; and we suppose  $\mu = 1$  if  $\Psi$  is an arbitrary  $N$ -function.*

- 2.

$$\int_0^T F(t, \mathbf{x}) dt \rightarrow \infty \quad \text{as} \quad |\mathbf{x}| \rightarrow \infty. \quad (37)$$

3.  $\Psi \in \Delta_2$  or, alternatively,  $\alpha_0^{-1} T \Phi^{-1}(1/T) \|b_1\|_{L^1} \Lambda < 1$ .

*Then the action integral  $I$  is coercive.*

*Proof.* In the following estimates, we will use (30), the decomposition  $\mathbf{u} = \bar{\mathbf{u}} + \tilde{\mathbf{u}}$ , Hölder's inequality and Wirtinger's inequality. Namely,

$$\begin{aligned} I(\mathbf{u}) &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T F(t, \mathbf{u}) dt \\ &= \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T [F(t, \mathbf{u}) - F(t, \bar{\mathbf{u}})] dt + \int_0^T F(t, \bar{\mathbf{u}}) dt \\ &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - \int_0^T b_1(t)(1 + |\tilde{\mathbf{u}}(t)|^\mu) dt + \int_0^T F(t, \bar{\mathbf{u}}) dt \\ &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - \|b_1\|_{L^1} (1 + \|\tilde{\mathbf{u}}\|_{L^\infty}^\mu) + \int_0^T F(t, \bar{\mathbf{u}}) dt \\ &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - \|b_1\|_{L^1} \left( 1 + \left[ T \Phi^{-1} \left( \frac{1}{T} \right) \right]^\mu \|\dot{\mathbf{u}}\|_{L^\Phi}^\mu \right) \\ &\quad + \int_0^T F(t, \bar{\mathbf{u}}) dt \\ &= \alpha_0 J_{C, \mu}(\dot{\mathbf{u}}) - \|b_1\|_{L^1} + \int_0^T F(t, \bar{\mathbf{u}}) dt, \end{aligned} \quad (38)$$

where  $C = \alpha_0^{-1} [T \Phi^{-1}(1/T)]^\mu \|b_1\|_{L^1}$ . Suppose that  $\mathbf{u}_n$  is a sequence in  $\mathcal{E}_d^\Phi(\lambda)$  such that the sequence  $\bar{\mathbf{u}}_n$  is bounded in  $\mathbb{R}^d$  and  $\|\mathbf{u}_n\|_{W^1 L^\Phi} \rightarrow \infty$ . Then the Wirtinger's inequality implies that  $\|\dot{\mathbf{u}}_n\|_{L^\Phi} \rightarrow \infty$ . Therefore, one of the following statements holds true  $\|\dot{\mathbf{u}}_n\|_{L^\Phi} \rightarrow \infty$  or  $|\bar{\mathbf{u}}_n| \rightarrow \infty$ . On the other hand, (31) and (37) imply that the integral  $\int_0^T F(t, \bar{\mathbf{u}}_n) dt$  is lower bounded. These observations, the lower bound of  $I$  given by (38), assumption 3 in Theorem 5.3 and Lemma 5.2 imply the desired result.  $\square$



Following [25] we say that  $F$  satisfies the condition (A) if  $F(t, \mathbf{x})$  is a Carathéodory function,  $F$  verifies (31) and  $F$  is continuously differentiable with respect to  $\mathbf{x}$ . Moreover, the next inequality holds

$$|D_{\mathbf{x}}F(t, \mathbf{x})| \leq a(|\mathbf{x}|)b_0(t), \quad \text{for a.e. } t \in [0, T] \text{ and for every } \mathbf{x} \in \mathbb{R}^d. \quad (39)$$

The following result was proved in [25, p. 18].

**Lemma 5.4.** *Suppose that  $F$  satisfies condition (A) and (37),  $F(t, \cdot)$  is differentiable and convex a.e.  $t \in [0, T]$ . Then there exists  $\mathbf{x}_0 \in \mathbb{R}^d$  such that*

$$\int_0^T D_{\mathbf{x}}F(t, \mathbf{x}_0) dt = 0. \quad (40)$$

**Theorem 5.5.** *Let  $\mathcal{L}$  be as in Theorem 5.3 and let  $F$  be as in Lemma 5.4. In addition, assume that  $\Psi \in \Delta_2$  or, alternatively  $\alpha_0^{-1}T\Phi^{-1}(1/T)a(|\mathbf{x}_0|)\|b_0\|_{L^1}\Lambda < 1$ , with  $a$  and  $b_0$  as in (31) and  $\mathbf{x}_0 \in \mathbb{R}^d$  any point satisfying (40). Then  $I$  is coercive.*

*Proof.* Using (30), [25, Eq. (18), p.17], the decomposition  $\mathbf{u} = \bar{\mathbf{u}} + \tilde{\mathbf{u}}$ , (40), (9) and Wirtinger's inequality, we get

$$\begin{aligned} I(\mathbf{u}) &\geq \alpha_0 \rho_{\Phi} \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T F(t, \mathbf{x}_0) dt + \int_0^T D_{\mathbf{x}}F(t, \mathbf{x}_0) \cdot (\mathbf{u} - \mathbf{x}_0) dt \\ &= \alpha_0 \rho_{\Phi} \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T F(t, \mathbf{x}_0) dt + \int_0^T D_{\mathbf{x}}F(t, \mathbf{x}_0) \cdot \tilde{\mathbf{u}} dt \\ &\quad + \int_0^T D_{\mathbf{x}}F(t, \mathbf{x}_0) \cdot (\bar{\mathbf{u}} - \mathbf{x}_0) dt \\ &= \alpha_0 \rho_{\Phi} \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + \int_0^T F(t, \mathbf{x}_0) dt + \int_0^T D_{\mathbf{x}}F(t, \mathbf{x}_0) \cdot \tilde{\mathbf{u}} dt \\ &\geq \alpha_0 \rho_{\Phi} \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) - a(|\mathbf{x}_0|)\|b_0\|_{L^1} - a(|\mathbf{x}_0|)\|b_0\|_{L^1}T\Phi^{-1} \left( \frac{1}{T} \right) \|\dot{\mathbf{u}}\|_{L^{\Phi}} \\ &= \alpha_0 J_{C,1}(\dot{\mathbf{u}}) - a(|\mathbf{x}_0|)\|b_0\|_{L^1} \end{aligned} \quad (41)$$

with  $C := \alpha_0^{-1}a(|\mathbf{x}_0|)\|b_0\|_{L^1}T\Phi^{-1}(1/T)$ .

Let  $\alpha$  be as in Corollary 2.3, then it is a non decreasing majorant of  $a$ . Using that  $F(t, \bar{\mathbf{u}}/2) \leq (1/2)F(t, \mathbf{u}) + (1/2)F(t, -\tilde{\mathbf{u}})$  and taking into account that  $\Phi$  is a non negative function, inequality (31), Hölder's inequality, Corollary 2.3 and Wirtinger's

inequality, we obtain

$$\begin{aligned}
 I(\mathbf{u}) &\geq \alpha_0 \rho_\Phi \left( \frac{\dot{\mathbf{u}}}{\Lambda} \right) + 2 \int_0^T F(t, \bar{\mathbf{u}}/2) dt - \int_0^T F(t, -\tilde{\mathbf{u}}) dt \\
 &\geq 2 \int_0^T F(t, \bar{\mathbf{u}}/2) dt - \|b_0\|_{L^1} \|\mathbf{a}(\tilde{\mathbf{u}})\|_{L^\infty} \\
 &\geq 2 \int_0^T F(t, \bar{\mathbf{u}}/2) dt - \|b_0\|_{L^1} \alpha(\|\tilde{\mathbf{u}}\|_{L^\infty}) \\
 &\geq 2 \int_0^T F(t, \bar{\mathbf{u}}/2) dt - C \alpha(C \|\dot{\mathbf{u}}\|_{L^\Phi})
 \end{aligned} \tag{42}$$

with a certain constant  $C > 0$ .

Let  $\mathbf{u}_n$  be a sequence in  $W^1 L_d^\Phi$  such that  $\|\mathbf{u}_n\|_{W^1 L^\Phi} \rightarrow \infty$ . We need to consider two situations:

- i) If  $\|\dot{\mathbf{u}}_n\|_{L^\Phi} \rightarrow \infty$  then, from (41) and Lemma 5.2, we have  $I(\mathbf{u}_n) \rightarrow \infty$ .
- ii) If  $\|\dot{\mathbf{u}}_n\|_{L^\Phi}$  is bounded and  $\|\mathbf{u}_n\|_{L^\Phi} \rightarrow \infty$ , then we obtain  $\bar{\mathbf{u}}_n \rightarrow \infty$ , reasoning in a similar way to that developed in the proof of Theorem 5.3. This fact together with (42) finishes the proof.  $\square$

## 6 Main result

For the lower semicontinuity of  $I$  we only need to adapt some results of [15] to our problem. **Este parrafo introductivo deberia hacer mejor propaganda del resultado principal**

**Theorem 6.1.** *We suppose that  $\mathcal{L}(t, \mathbf{x}, \mathbf{y})$ ,  $F(t, \mathbf{x})$  are Carathéodory functions satisfying*

$$\mathcal{L}(t, \mathbf{x}, \mathbf{y}) \geq \Phi(|\mathbf{y}|) + F(t, \mathbf{x}), \tag{43}$$

where  $\Phi$  is an  $N$ -function. We also assume that the function  $F$  satisfies inequality (31) and  $\mathcal{L}(t, \mathbf{x}, \cdot)$  is convex in  $\mathbb{R}^d$  for each  $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^d$ . Let  $\{\mathbf{u}_n\} \subset W^1 L^\Phi$  be a sequence such that  $\mathbf{u}_n$  converges uniformly to a function  $\mathbf{u} \in W^1 L^\Phi$  and  $\dot{\mathbf{u}}_n$  converges in the weak topology of  $L_d^1$  to  $\dot{\mathbf{u}}$ . Then

$$I(\mathbf{u}) \leq \liminf_{n \rightarrow \infty} I(\mathbf{u}_n). \tag{44}$$

*Proof.* First we note that (43) and (31) imply that  $I$  is defined on  $W^1 L^\Phi$  taking values in the interval  $(-\infty, +\infty]$ .

Let  $\{\mathbf{u}_n\}$  be a sequence satisfying the assumptions of the theorem. We define the Carathéodory function  $\hat{\mathcal{L}} = \mathcal{L} - F$ . Let  $\hat{I}$  be the associated to  $\hat{\mathcal{L}}$  action integral. Using [15, Thm. 2.1, p. 243] with  $\hat{\mathcal{L}}$  instead  $f$ , we have that

$$\int_0^T \hat{\mathcal{L}}(t, \mathbf{u}, \dot{\mathbf{u}}) dt \leq \liminf_{n \rightarrow \infty} \int_0^T \hat{\mathcal{L}}(t, \mathbf{u}_n, \dot{\mathbf{u}}_n) dt. \tag{45}$$

From the uniform convergence of  $\mathbf{u}_n$  and the Carathéodory conditions for  $F$  we obtain that  $F(t, \mathbf{u}_n(t)) \rightarrow F(t, \mathbf{u}(t))$  a.e.  $t \in [0, T]$ . Since  $\mathbf{u}_n$  are uniformly bounded,

the inequality (31) imply that there exists  $g \in L^1_1([0, T])$  such that  $|F(t, \mathbf{u}_n(t))| \leq g(t)$ . From the Dominated Convergence Theorem we have that

$$\lim_{n \rightarrow \infty} \int_0^T F(t, \mathbf{u}_n(t)) dt = \int_0^T F(t, \mathbf{u}(t)) dt. \quad (46)$$

Taking account of (45) and (46) we obtain (44).  $\square$

**Theorem 6.2.** *We suppose  $\Phi$  and  $\Psi$  complementary  $N$ -functions. We assume that the Carathéodory function  $\mathcal{L}(t, \mathbf{x}, \mathbf{y})$  is strictly convex in  $\mathbf{y}$ ,  $D_{\mathbf{y}}\mathcal{L}$  is  $T$ -periodic with respect to  $T$  and that it satisfies (2), (3), (4), (30) and (31). In addition, we suppose that some of the following statements hold (we recall the definitions and properties of  $\alpha_0$ ,  $b_1$ ,  $\mathbf{x}_0$  and  $b_0$  from (30), (36), (40) and (39) respectively)*

1.  $\Psi \in \Delta_2$ , (36) and (37).
2. As item 1 but with  $\alpha_0^{-1}T\Phi^{-1}(1/T) \|b_1\|_{L^1}\Lambda < 1$  instead of  $\Psi \in \Delta_2$ .
3.  $\Psi \in \Delta_2$ ,  $F$  satisfies condition (A), (37) and  $F(t, \cdot)$  is convex a.e.  $t \in [0, T]$ .
4. As item 3 but  $\alpha_0^{-1}T\Phi^{-1}(1/T) a(|\mathbf{x}_0|)\|b_0\|_{L^1}\Lambda < 1$  instead of  $\Psi \in \Delta_2$ .

Then problem (1) has a solution.

*Proof.* Let  $\{\mathbf{u}_n\} \subset W^1L_T^\Phi$  be a minimizing sequence for the problem  $\min\{I(\mathbf{u}) | \mathbf{u} \in W^1L_T^\Phi\}$ . Since  $I(\mathbf{u}_n)$ ,  $n = 1, 2, \dots$  is bounded o es acotada solo por arriba?, el espacio sobre el que minimizamos no excluye la posibilidad de  $-\infty$ . De todas maneras solo con acotacion superior se tiene el mismo resultado, Theorem 5.3 (or Theorem 5.5 according to which of the items 1-4 hold true) implies that  $\{\mathbf{u}_n\}$  is norm bounded in  $W^1L_d^\Phi$ . Hence, in virtue of Corollary 2.2 we can assume that  $\mathbf{u}_n$  converges uniformly to a  $T$ -periodic continuous function  $\mathbf{u}$ . The space  $L_d^\Phi$  is a predual space, concretely  $L_d^\Phi = [E_d^\Psi]^*$ . Therefore, from [26, Cor. 5, p. 148] and since  $\dot{\mathbf{u}}_n$  is bounded in  $L_d^\Phi$ , there exists a subsequence (again denoted  $\dot{\mathbf{u}}_n$ ) such that  $\dot{\mathbf{u}}_n$  converges to a function  $\mathbf{v} \in L_d^\Phi$  in the weak\* topology. From this fact and the uniform convergence of  $\mathbf{u}_n$  to  $\mathbf{u}$  we obtain for every  $T$ -periodic function  $\xi \in C^\infty([0, T], \mathbb{R}^d) \subset E_d^\Psi$

$$\int_0^T \xi \cdot \mathbf{u} dt = \lim_{n \rightarrow \infty} \int_0^T \xi \cdot \mathbf{u}_n dt = - \lim_{n \rightarrow \infty} \int_0^T \xi \cdot \dot{\mathbf{u}}_n dt = - \int_0^T \xi \cdot \mathbf{v} dt.$$

Thus  $\mathbf{v} = \dot{\mathbf{u}}$  a.e.  $t \in [0, T]$  (see [25, p. 6]) and  $\mathbf{u} \in W^1L_T^\Phi$ .

Finally from the relations  $[L_d^1]^* = L_d^\infty \subset E_d^\Psi$  and  $L_d^\Phi \subset L_d^1$  we obtain that  $\dot{\mathbf{u}}_n$  converges to  $\dot{\mathbf{u}}$  in the weak topology of  $L_d^1$ . Consequently Theorem 6.1, applying with the  $N$ -function  $\alpha_0\Phi(|\cdot|/\Lambda)$ , implies

$$I(\mathbf{u}) \leq \liminf_{n \rightarrow \infty} I(\mathbf{u}_n) = \min_{\mathbf{u} \in W^1L_T^\Phi} I(\mathbf{u}).$$

Hence  $\mathbf{u}$  is a minimum, and therefore a critical point, of  $I$ . Invoking Theorem 4.1 we conclude the proof.  $\square$

## References

- [1] S. Acinas and S. Favier. “Maximal inequalities in Orlicz spaces”. In: *Int. J. Math. Anal. (Ruse)* 6.41-44 (2012), pp. 2179–2198.
- [2] S. Acinas, G. Giubergia, F. Mazzone, and E. Schwindt. “On Estimates for the Period of Solutions of Equations Involving the  $\phi$ -Laplace Operator”. In: *Journal of Abstract Differential Equations and Applications* 5.1 (2014), pp. 21–34.
- [3] R.A. Adams and J.J.F. Fournier. *Sobolev Spaces*. Elsevier Science, 2003.
- [4] A. Ambrosetti and G. Prodi. *A Primer of Nonlinear Analysis*. Cambridge University Press, 1995.
- [5] C. Bereanu and J. Mawhin. “Existence and multiplicity results for some nonlinear problems with singular  $\phi$ -Laplacian”. In: *Journal of Differential Equations* 243 (2007).
- [6] C. Bereanu and J. Mawhin. “Nonhomogeneous boundary value problems for some nonlinear equations with singular  $\phi$ -Laplacian”. In: *Journal of Mathematical Analysis and Applications* 352 (2009).
- [7] C. Bereanu and J. Mawhin. “Periodic solutions of nonlinear perturbations of  $\phi$ -Laplacians with possibly bounded  $\phi$ ”. In: *Nonlinear Analysis: Theory, Methods & Applications* 68 (2008).
- [8] A. Cianchi. “A fully anisotropic Sobolev inequality”. In: *Pacific J. Math* 196.2 (2000), pp. 283–295.
- [9] A. Cianchi. “Local boundedness of minimizers of anisotropic functionals”. In: *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*. Vol. 17. 2. Elsevier. 2000, pp. 147–168.
- [10] A. Cianchi. “Some results in the theory of Orlicz spaces and applications to variational problems”. In: *Nonlinear analysis, function spaces and applications* 6 (1999), pp. 50–92.
- [11] A. Cianchi and N. Fusco. “Gradient regularity for minimizers under general growth conditions”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 1999.507 (1999), pp. 15–36.
- [12] N. Clavero. *Optimal Sobolev embeddings and Function Spaces*. 2011. URL: <http://www.maia.ub.edu/~soria/sobolev1.pdf>.
- [13] B. Driver. *Analysis Tools with Applications*. 2003. URL: [http://www.math.ucsd.edu/~bdriver/231-02-03/Lecture\\_Notes/PDE-Anal-Book/analpde1.pdf](http://www.math.ucsd.edu/~bdriver/231-02-03/Lecture_Notes/PDE-Anal-Book/analpde1.pdf).
- [14] D.E. Edmunds, R. Kerman, and L. Pick. “Optimal Sobolev imbeddings involving rearrangement-invariant quasinorms”. In: *Journal of Functional Analysis* 170.2 (2000), pp. 307–355.
- [15] I. Ekeland and R. Témam. *Convex Analysis and Variational Problems*. Society for Industrial and Applied Mathematics, 1999.

## References

---

- [16] S. Favier and F. Zó. “Extension of the best approximation operator in Orlicz spaces and weak-type inequalities”. In: *Abstr. Appl. Anal.* 6.2 (2001), pp. 101–114.
- [17] M. García-Huidobro, V.K. Le, R. Manásevich, and K. Schmitt. “On principal eigenvalues for quasilinear elliptic differential operators: an Orlicz-Sobolev space setting”. In: *Nonlinear Differential Equations and Applications NoDEA* 6 (1999).
- [18] H. Hudzik and L. Maligranda. “Amemiya norm equals Orlicz norm in general”. In: *Indagationes Mathematicae* 11.4 (2000), pp. 573–585.
- [19] T. Kato. “Demicontinuity, hemicontinuity and monotonicity”. In: *Bulletin of the American Mathematical Society* 70.4 (1964), pp. 548–550.
- [20] R. Kerman and L. Pick. “Optimal Sobolev imbeddings”. In: *Forum Mathematicum*. Vol. 18. 2006, pp. 535–570.
- [21] M. A. Krasnosel’skiĭ and Ja. B. Rutickiĭ. *Convex functions and Orlicz spaces*. Translated from the first Russian edition by Leo F. Boron. Groningen: P. Noordhoff Ltd., 1961.
- [22] L. Maligranda. *Orlicz spaces and interpolation*. Vol. 5. Seminários de Matemática [Seminars in Mathematics]. Universidade Estadual de Campinas, Departamento de Matemática, Campinas, 1989.
- [23] R. Manásevich and J. Mawhin. “Boundary value problems for nonlinear perturbations of vector p-Laplacian-like operators”. In: *J. Korean Math. Soc* 37.5 (2000), pp. 665–685.
- [24] R. Manásevich and J. Mawhin. “Periodic Solutions for Nonlinear Systems with p-Laplacian-Like Operators”. In: *Journal of Differential Equations* 145.2 (1998), pp. 367–393.
- [25] J. Mawhin and M. Willem. *Critical point theory and Hamiltonian systems*. Vol. 74. Applied Mathematical Sciences. Springer-Verlag, New York, 1989, pp. xiv+277.
- [26] M. M. Rao and Z. D. Ren. *Theory of Orlicz spaces*. M. Dekker, 1991.
- [27] Z.D. Ren. *Applications Of Orlicz Spaces*. Monographs and textbooks in pure and applied mathematics. Taylor & Francis, 2002.
- [28] R.T. Rockafellar and R.J.B. Wets. *Variational Analysis*. Springer, 2009.
- [29] G. Schappacher. “A Notion of Orlicz Spaces for Vector Valued Functions”. English. In: *Applications of Mathematics* 50.4 (2005), pp. 355–386.
- [30] M. Struwe. *Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, 2008.
- [31] C.-L. Tang. “Periodic solutions for nonautonomous second order systems with sublinear nonlinearity”. In: *Proceedings of the American Mathematical Society* (1998), pp. 3263–3270.
- [32] C.-L. Tang. “Periodic solutions of non-autonomous second-order systems with  $\gamma$ -quasisubadditive potential”. In: *Journal of Mathematical Analysis and Applications* 189.3 (1995), pp. 671–675.

- [33] C.-L. Tang and X.-P. Wu. “Periodic solutions for second order systems with not uniformly coercive potential”. In: *Journal of Mathematical Analysis and Applications* 259.2 (2001), pp. 386–397.
- [34] X. Tang and X. Zhang. “Periodic solutions for second-order Hamiltonian systems with a p-Laplacian”. In: *Annales UMCS, Mathematica* 64.1 (2010), pp. 93–113.
- [35] Y. Tian and W. Ge. “Periodic solutions of non-autonomous second-order systems with a p-Laplacian”. In: *Nonlinear Analysis: Theory, Methods & Applications* 66.1 (2007), pp. 192–203.
- [36] X.-P. Wu and C.-L. Tang. “Periodic solutions of a class of non-autonomous second-order systems”. In: *Journal of Mathematical Analysis and Applications* 236.2 (1999), pp. 227–235.
- [37] B. Xu and C.-L. Tang. “Some existence results on periodic solutions of ordinary p-Laplacian systems”. In: *Journal of mathematical analysis and applications* 333.2 (2007), pp. 1228–1236.
- [38] Y.-W. Ye and C.-L. Tang. “Periodic solutions for some nonautonomous second order Hamiltonian systems”. In: *Journal of Mathematical Analysis and Applications* 344.1 (2008), pp. 462–471.
- [39] F. Zhao and X. Wu. “Periodic solutions for a class of non-autonomous second order systems”. In: *Journal of mathematical analysis and applications* 296.2 (2004), pp. 422–434.