

Review article

Prevalence of depressive symptoms in primary school students in China: A systematic review and meta-analysis



Dan-Dan Xu^{a,b,1}, Wen-Wang Rao^{a,1}, Xiao-Lan Cao^{c,1}, Si-Ying Wen^{c,1}, Feng-Rong An^{d,1},
Weng-Ian Che^{e,f}, Daniel T. Bressington^g, Teris Cheung^g, Gabor S. Ungvari^{h,i}, Yu-Tao Xiang^{a,j,*}

^a Unit of Psychiatry, Faculty of Health Sciences, University of Macau, Macao SAR, China

^b Department of Biology, Faculty of Sciences, Harbin University, Harbin, China

^c Shenzhen Key Laboratory for Psychological Healthcare & Shenzhen Institute of Mental Health, Shenzhen Kangning Hospital & Shenzhen Mental Health Center, Shenzhen, China

^d The National Clinical Research Center for Mental Disorders & Beijing Key Laboratory of Mental Disorders, Beijing Anding Hospital & the Advanced Innovation Center for Human Brain Protection, Capital Medical University, Beijing, China

^e Unit of Clinical Epidemiology, Department of Medicine, Solna, Karolinska Institutet, Stockholm, Sweden

^f Department of Public Health Science, Karolinska Institutet, Stockholm, Sweden

^g School of Nursing, Hong Kong Polytechnic University, Hong Kong SAR, China

^h Division of Psychiatry, School of Medicine, University of Western Australia, Perth, Australia

ⁱ University of Notre Dame Australia, Fremantle, Australia

^j Center for Cognition and Brain Sciences, University of Macau, Macao SAR, China

ARTICLE INFO

Keywords:

Depressive symptoms
Primary school students
Prevalence
Meta-analysis
China

ABSTRACT

Backgrounds: Depression in children and adolescents is usually under-recognized. The findings of epidemiological studies on depressive symptoms in primary school students are inconsistent across studies. This study reports a systematic review and meta-analysis on the prevalence of depressive symptoms in primary school students in China.

Methods: Literature search was performed in both international (PubMed, PsycINFO, EMBASE) and Chinese (China National Knowledge Internet, WANFANG Data and Chinese Biological Medical Literature) databases. The random-effects model was used to analyze data.

Results: Twenty-seven studies involving 42,374 subjects were included. The pooled prevalence of depressive symptoms in Chinese primary school students was 17.2% (95% CI: 14.3%–20.5%). Subgroup analyses found that the prevalence significantly varied between geographic regions, with western China reporting the highest prevalence. Meta-regression analyses found that year of survey and study quality were significantly associated with the prevalence of depressive symptoms.

Conclusions: Given the high prevalence of depressive symptoms and its negative health outcomes, preventive measures, regular screening and effective treatments need to be implemented for this population.

1. Introduction

Depression is the most common mental health problem among children and adolescents and is also one of the top ten causes of disability among teenagers globally (Global Burden of Disease 2015 Disease and Injury Incidence and Prevalence Collaborators, 2016). Depression in childhood and adolescence is associated with impaired psychosocial functions (Ayuso-Mateos et al., 2010; Judd and Akiskal, 2000) and increased risk of future depression in adulthood (Hankin et al., 1998; Naicker et al., 2013; Rutter et al., 2006).

Depression could exist in a continuum of severity, ranging from depressive symptoms, minor depression to major depressive disorder (MDD) (Angst et al., 2000; Judd and Akiskal, 2000). Similar to clinically diagnosed depression, depressive symptoms also have negative health impacts among children and adolescents, such as social problems (Verboom et al., 2014), poor academic performance (Verboom et al., 2014), substance use (Earnshaw et al., 2017), obesity (Johnson et al., 2018) and impaired cognitive functions (Calvete et al., 2013; Cole et al., 2011).

Depression prevalence usually increases substantially throughout

* Corresponding author: 3/F, Building E12, Faculty of Health Sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR, China.
E-mail address: xyutly@gmail.com (Y.-T. Xiang).

¹ These authors contributed equally to the work.

the period of adolescence (Hankin et al., 1998; Lewinsohn et al., 1999), with the peak in mid- to late-adolescence (Cyranowski et al., 2000; Patton and Viner, 2007). Despite this observed peak in prevalence, studies examining depression in mid- and late- adolescence, i.e., in primary and secondary schools (Tang et al., 2018) have concluded that depression in primary school students was often overlooked. Although the prevalence of clinical depression (i.e., minor and major depression) is low, e.g., 1% in children in New Zealand (Hankin et al., 1998) and 1.3% in children and adolescents in China (Xu et al., 2018), depressive symptoms are relatively common in school-aged children. Compared with their Western counterparts, school children in Asian countries are more likely to experience depressive symptoms, for instance, the prevalence of depressive symptoms was 17.3% in Korea (Kwak et al., 2008) and 16.9% in Singapore (Magiati et al., 2015), while it was 10.6% in Italy (Frigerio et al., 2001), 9.85% in Greece (Giannakopoulos et al., 2009), 10% in Sweden (Larsson and Melin, 1992) and 8.3% in the UK (Charman and Pervova, 1996). Possible explanations on the discrepancy between Asian and Western countries include daily pressures arising from greater emphasis on academic achievement and higher academic stress in Asian countries (Magiati et al., 2015; Tepper et al., 2008). The use of different research methodologies and assessment tools may also result in the discrepancy of depression prevalence. Depression can be assessed by clinical interviews, structured interviews, self-rating scales and interviewer-rated scales (Birmaher et al., 1996). In the past decades, several instruments have been developed to assess depression, such as the Center for Epidemiologic Studies Depression Scale (CES-D) (Radloff, 1977), Beck Depression Inventory (BDI) (Teri, 1982) and instruments specific for children and adolescents, e.g., Children's Depression Inventory (CDI) (Kovacs, 1992) and Depression Self-Rating Scale for Children (DSRSC) (Birleson, 1981; Birleson et al., 1987).

Depressive symptoms in school children are associated with certain socio-demographic factors, such as gender (Frigerio et al., 2001; Giannakopoulos et al., 2009; Wang et al., 2016), age (Denda et al., 2006; Kwak et al., 2008), family socioeconomic status (SES) (Frigerio et al., 2001; Giannakopoulos et al., 2009; Zhou et al., 2018b), satisfaction with body image (Kwak et al., 2008; Li et al., 2007), anxiety (Lee et al., 2013), academic stress (Wang et al., 2016), and relationships with peers and family (Kwak et al., 2008; Larsson and Melin, 1992; Lee et al., 2013).

In order to reduce the negative health outcomes and develop preventive measures, it is important to understand the epidemiology of depressive symptoms in school children. The epidemiology of affective problems including depressive symptoms is considerably influenced by socioeconomic contexts (Compton et al., 2006; Kleinman, 2004), thus epidemiological studies of depressive symptoms in different countries and areas with different socioeconomic backgrounds have questionable generalizability and need to be examined separately. Although some studies have examined the prevalence of depressive symptoms in primary school students in China, the findings are inconsistent. To date, no systematic review and meta-analysis of epidemiology of depressive symptoms in this population has been published, which gave us the impetus to conduct this meta-analysis to examine the depressive symptoms and associated factors in primary school students in China. Based on previous findings (Rao et al., 2019), we assumed that depressive symptoms among primary school students in China would be common, and the prevalence would be associated with certain demographic characteristics, such as gender and use of different instruments.

2. Methods

This study protocol was prospectively registered in the International Prospective Register of Systematic Reviews (PROSPERO; registration number: CRD42019124895).

2.1. Data sources and search strategies

Two researchers (DDX and WWR) independently searched both international (PubMed, PsycINFO and EMBASE) and Chinese (China National Knowledge Internet, WANFANG Data and Chinese Biological Medical Literature) databases from commencement to June 18, 2018, using the search terms including: adolescents, students, children, teen, teenagers, depression, depressive disorders, depressive symptoms, prevalence, epidemiology, survey, rate, percentage, China and Chinese. Additionally, the reference lists of included articles and reviews were checked for additional studies. The first or corresponding authors of included studies were contacted for additional information if necessary. Literature search and screening were conducted following the guideline of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and Meta-analysis of observational studies in Epidemiology reporting guideline (Moher et al., 2009; Stroup et al., 2000).

2.2. Study criteria

Five researchers (DDX, WWR, XLC, SYW and WIC) independently screened articles by titles and abstracts first and then read the full texts of potentially eligible articles. According to the PRISMA statement, the following inclusion criteria were developed according to the PICOS acronym: Participants (P) studies on primary school students; using validated instruments to assess depressive symptoms, such as CES-D, BDI, CDI and DSRSC; Intervention (I): not applicable; Comparison (C): not applicable; Outcomes (O): reporting prevalence of depressive symptoms or relevant information that could generate the prevalence of depressive symptoms; and Study design (S): cross-sectional or cohort studies (only baseline data were extracted) conducted in mainland China (China thereafter) published in English or Chinese. Studies were excluded if they were conducted in special populations (e.g., left-behind children, outpatients, inpatients, obese children or those with physical or other psychiatric disorders), or focused on major depressive disorder. Any disagreement was resolved by a discussion with a senior researcher (YTX).

2.3. Data extraction

The same group of researchers independently extracted the following data from eligible studies using a standardized data collection form: geographic location, year of survey, sampling methods, sample size, response rate, mean age, percentage of male gender, school grades, assessment instrument and cutoff, and the prevalence estimates of depressive symptoms. If the study was based on a multi-center design, the data in each study site was extracted for subgroup analyses. If one dataset was used in different papers, only the one with the largest sample size was included. Data were extracted and double-checked independently by two researchers (DDX and WWR).

2.4. Quality assessment

The same researchers independently assessed study quality using the quality assessment instrument for epidemiological studies (Boyle, 1998; Ibrahim et al., 2013; Loney et al., 1998) with the following eight items: (1) target population; (2) sampling; (3) the response rate; (4) non-respondents; (5) the representativeness; (6) standardized data collection methods; (7) validated criteria for depressive symptoms; (8) the prevalence estimates with confidence intervals. Any discrepancy was resolved by a discussion with a senior researcher (YTX).

2.5. Statistical analysis

Due to different sampling methods, demographic characteristics and instruments between studies, the pooled prevalence estimate of depressive symptoms was calculated as effect size (ES) using the Der

Simonian and Laird random-effects model (Borenstein et al., 2010). Heterogeneity across studies was estimated with Cochran's Q test and I^2 statistic, with I^2 of $\geq 50\%$ or Cochran's Q of $p < 0.05$ indicating significant heterogeneity (Higgins et al., 2003). Subgroup analyses were conducted to explore the sources of heterogeneity according to the following categorical variables: gender (male vs. female), geographic region (Eastern vs. Central vs. Western China, according to the National Bureau of Statistics of China), year of survey (2001–2010 vs. 2011–2016 by median spitting method), and screening instrument (CDI vs. CES-D vs. DSRSC). Meta-regression analyses were used to examine the associations of the year of survey (continuous variable) and study quality with the prevalence estimates of depressive symptoms. Sensitivity analyses were conducted to explore the outlying studies. The funnel plots, Begg's test and Egger's test were used to assess publication bias. Significance level was set at 0.05 (two-tailed). The meta-analyses were conducted using the Comprehensive Meta-Analysis software, Version 2 (CMA, Biostat Inc., Englewood, New Jersey, USA).

3. Results

3.1. Study characteristics

Of the 12,374 articles initially identified, 27 studies with 42,374

participants met the study entry criteria and were included (Fig. 1). Study characteristics are summarized in Table 1. Studies were conducted from 2001 to 2016. Apart from one household survey (Zhou et al., 2018b), the remaining studies were school-based surveys. The most widely used screening instrument for depressive symptoms was the CDI ($n = 19$), followed by the DSRSC ($n = 5$) and the CES-D ($n = 3$). Only two studies were longitudinal (Wang and Su, 2005; Zhao, 2014), while the remaining were cross-sectional.

3.2. Prevalence of depressive symptoms in Chinese primary school students

The pooled prevalence estimate of depressive symptoms in Chinese primary school students was 17.2% (95% CI: 14.3%–20.5%, $I^2 = 98.65\%$) (Fig. 2).

Subgroup analyses did not find significant associations of prevalence estimates of depressive symptoms with gender, year of survey and screening instruments (all p values > 0.05). There were different prevalence estimates of depressive symptoms between geographical regions ($Q = 10.416$, $p = 0.005$); the pooled prevalence estimate was highest in western China (22.1%, 95% CI: 18.5%–26.2%), followed by eastern (15.8%, 95%CI: 11.6%–21.2%) and central regions of China (13.7%, 95%CI: 10.6%–17.6%) (Table 2). Meta-regression analyses revealed significant association of depressive symptoms prevalence




Fig. 1. PRISMA flowchart of literature selection.

Table 1
Characteristics of studies included in the meta-analysis

First author, year	References	Province	Region	Survey year	SM	ESS	RS (%)	ESS ^b	Age range	Mean age (Mean ± SD)	Male (%)	Grade	Scale	Cutoff	Prevalence (%)	Quality score
Wu LJ, 2015	(Wu et al., 2015)	Beijing	E	NR	C	1472	NR	1472	NR	NR	50.6	3–5	CDI	>12	31.05	5
Xu J, 2008 ^a	(Xu et al., 2008)	Anhui	C	NR	SC	2071	99.7	2071	8–12	9.7 ± 1.0	52.9	3–5	CDI	>19	11.88	6
Xu J, 2008 ^b	(Xu et al., 2008)	Guangdong	E	NR	SC	1153	95.5	1153	8–14	9.7 ± 1.0	55.0	3–5	CDI	>19	8.24	6
Wu HN, 2017	(Wu et al., 2017)	Zhejiang	E	2014	SC	2000	NR	1020	NR	NR	NR	NA ^c	CDI	>19	10.78	4
Zhang XY, 2015	(Zhang and Zhou, 2015)	Jiangsu	E	NR	SR	2362	96.2	2362	NR	NR	53.34	3–5	CDI	>19	10.70	6
Zhao X, 2014	(Zhao, 2014)	Anhui	C	2009	C	816	77.9	404	NR	NR	NR	3–4	CDI	>19	13.11	6
Li C, 2013	(Li et al., 2013)	Chongqing	W	2010	MRC	3013	97.7	931	NR	NR	NR	4–6	CDI	≥19	17.80	6
Li YP, 2007	(Li et al., 2007)	Beijing	E	2005	C	3886	96.0	3886	9–10	NR	51.1	3–4	CDI	≥19	13.20	6
Liu SR, 2018	(Liu and Miao, 2018)	Xinjiang	W	2016	RSC	3610	97.9	1574	NR	NR	NR	4–6	CDI	≥19	32.27	6
Peng LI, 2018	(Peng et al., 2018)	Chongqing	W	2016	SC	3351	95.4	1607	NR	NR	NR	4–6	CDI	≥19	22.50	6
Wang L, 2016	(Wang et al., 2016)	Chongqing	W	2013	MSCR	10,357	79.5	5462	NR	NR	NR	2–6	CDI	≥19	22.32	7
Wang HZ, 2015	(Wang et al., 2015)	Henan	C	2010	MSC	3002	95.2	653	NR	NR	NR	5–6	CDI	≥19	9.65	6
Xie S, 2013	(Xie et al., 2016)	Hubei	C	2011	C	2888	97.9	692	NR	NR	NR	5–6	CDI	≥19	11.85	5
Yan M, 2012	(Yan, 2012)	Hubei	C	2010	SR	3007	NR	756	NR	NR	NR	5–6	CDI	≥19	14.29	4
Yang J, 2014	(Yang et al., 2014)	Guangdong	E	NR	SC	1676	93.1	1676	8–14	10.68 ± 1.26	49.2	3–6	CDI	≥19	14.08	5
Zhou Z, 2011	(Zhou and Wang, 2011)	Anhui	C	NR	SC	1884	95.2	939	NR	NR	NR	3–6	CDI	≥19	16.83	5
Zhu HQ, 2006	(Zhu et al., 2016)	Hainan	E	2015	SC	4866	97.4	2670	NR	NR	NR	4–6	CDI	≥19	31.80	6
Zhu JH, 2013	(Zhu et al., 2013)	Hubei	C	2010	SR	1975	89.8	743	NR	NR	NR	2–6	CDI	≥19	6.50	7
Li L, 2017	(Li, 2017)	Chongqing	W	NR	SC	1295	98.2	1295	7–13	10.19 ± 1.17	53.51	3–6	CDI	NA	16.53	5
Ding H, 2017	(Ding et al., 2017)	Hubei	C	2015	C	6406	NR	2628	NR	NR	NR	5–6	CES-D	≥16	9.78	5
Zhou Q, 2018	(Zhou et al., 2018b)	25 provinces	-	2012	MS	3056	NR	567	10–15	NR	NR	NA ^d	CES-D	≥18	29.30	5
Fan J, 2010	(Fan et al., 2010)	Shanghai	E	2009	RC	3628	98.5	3628	8–12	10.0 ± 1.0	51.70	3–5	CES-D	≥20	18.11	8
Gu JX, 2007	(Gu et al., 2007)	Hebei	E	2004	C	522	100	522	8–10	8.92 ± 1.13	51.70	2–5	DSRSC	≥15	18.01	6
Tang J, 2003	(Tang et al., 2003)	Hunan	C	NR	C	565	95.9	565	6–13	10.09 ± 1.53	51.30	2–6	DSRSC	≥15	17.17	5
Wang K, 2005	(Wang and Su, 2005)	Hunan	C	2001	C	209	NR	209	NR	NR	NR	2–4	DSRSC	≥15	40.67	4
Yong N, 2011	(Yong et al., 2011)	Chongqing	W	NR	SR	1417	93.0	1417	8–13	10 ± 1	51.5	4–6	DSRSC	≥15	23.15	6
Sun L, 2005	(Sun and Zhou, 2005)	Tianjin	E	NR	RC	539	95.7	454	NR	NR	NR	3–6	DSRSC	CF ^e	13.66	5
Qi YL, 2006	(Qi, 2006)	Jilin	NE	NR	C	1018	94.3	1018	NR	NR	58.3	4–6	BDI	NA	52.75	4

NR = Not reported; NA = Not available; SD = Standard deviation. ESS = Effective sample size. RS = Response rate. Region: C = Central; E = East; NE = Northeast; W = West. SM = Sampling method: C = Cluster sampling; M = Multistage sampling; R = Random sampling; S = Stratified sampling. Scale: CDI = Children's Depression Inventory; CES-D = Center for Epidemiologic Studies Depression Scale; DSRSC = Depression Self-Rating Scale for Children; BDI = Beck Depression Inventory.

^a Data reported about Anhui study site in Xu J, 2008;

^b Data reported about Guangdong study site in Xu J, 2008;

^c The samples were primary school students aged <14;

^d The samples were primary school students;

^e CF = Cutoff: 8–12 years: male ≥ 16, female ≥ 15; 13–16 years: male ≥ 14, female ≥ 16;

^p Primary school.




Fig. 2. Forest plot of prevalence of depressive symptoms in Chinese primary school students.

with the year of survey (slope = 0.056, $p < 0.001$) and study quality (slope = -0.068 , $p < 0.001$). Sensitivity analysis did not find outlying study that could significantly affect the primary results.

3.3. Quality assessment and publication bias

The quality assessment scores ranged from 4 to 8, with a mean score of 5.54 ± 0.94 . Three studies clearly described the non-respondents (Fan et al., 2010; Wang et al., 2016; Zhao, 2014) and four studies reported the prevalence estimates with 95% confidence intervals (Fan et al., 2010; Wang et al., 2016; Zhu et al., 2016; Zhu et al., 2013).

Table 2
Subgroup analyses on the prevalence of depression symptoms

Subgroups	Categories (Number of studies)	Proportion (%)	95% CI(%)	Events	Sample size	I ² (%) within subgroup	Q (p) across subgroups
Gender ^a	Male (12)	21.8	16.1–28.9	2149	10,415	98.20	1.224 (0.269)
	Female (12)	16.7	11.5–23.8	1456	9564	98.17	
Region	East (10)	15.8	11.6–21.2	3326	18,843	98.80	10.416 (0.005)
	Central (10)	13.7	10.6–17.6	1197	9660	95.48	
	West (6)	22.1	18.5–26.2	2797	12,286	95.86	
Survey year	2001–2010 (9)	15.3	12.1–19.2	1787	11,732	95.79	0.948 (0.33)
	2011–2016 (8)	18.5	13.6–24.7	3484	16,220	98.77	
Scale	CDI (19)	15.4	12.5–18.9	5740	31,366	98.48	3.247 (0.197)
	CES-D (3)	14.5	9.4–21.9	1011	6823	97.62	
	DSRSC (5)	21.3	15.5–28.7	666	3167	94.36	

Bold value: $p < 0.05$; CDI = Children's Depression Inventory; CES-D = Center for Epidemiologic Studies Depression Scale; DSRSC = Depression Self-Rating Scale for Children.

^a Twelve studies reported the depressive symptoms prevalence of both genders. The extracted data were divided into two arms for each study.




Fig. 3. Publication bias of studies included in the meta-analysis.

school students in China. There are around 103 million primary school students in China (Ministry of Education of the People's Republic of China, 2018). According to the current findings, this translates to approximately 17.7 million primary school students suffering from depressive symptoms.

The pooled prevalence in this meta-analysis is similar to the findings in Chinese children and adolescents aged <18 years (19.85%, 95% CI: 14.75%–24.96%) (Rao et al., 2019), but lower than that in Chinese secondary school students (24.3%, 95% CI, 21.3%–27.6%) (Tang et al., 2018). It is also similar to the corresponding figures in some Asian countries, such as 17.3% in the elementary students in Korea and 16.9% in children aged 8–12 years in Singapore (Kwak et al., 2008; Magiati et al., 2015). However, the pooled prevalence is higher than those reported in some Western countries, such as 10.6% in Italian elementary students, 9.85% in Greek children aged 8–12 years and 10% in Swedish school children aged 8–13 years (Frigerio et al., 2001; Giannakopoulos et al., 2009; Larsson and Melin, 1992). A comparative study found that Chinese adolescents aged 6–15 years were more likely to experience depressive symptoms than their counterparts in North America (Tepper et al., 2008). This discrepancy between Chinese and Western school children may be due to different academic stress and competition. Chinese school children usually face great academic stress and too much homework from primary schools. In addition, Chinese parents often have high expectations on children's academic performance and are more strict in supervising their children (Magiati et al., 2015). Some Chinese parents may also be less likely to identify their children's emotional problems and seek help from health professionals than their Western counterparts (Huh et al., 2019; Tepper et al., 2008).

Subgroup analyses found that primary school students in western China reported higher prevalence of depressive symptoms, compared to those in central and eastern China. Since the “China's Western Development Program” launched in 2000, western China has been undergoing dramatic socioeconomic transformation, leading to rapid urbanization and rural-urban migration (Lai, 2002) and new public health concerns (Mou et al., 2013; Ren et al., 2019), particularly in western China (Cai et al., 2017). Some studies found that migrant children and left-behind families are more likely to have mental health problems (Fellmeth et al., 2018; Lu et al., 2012). In addition, as the main labor-sending areas in China, many parents in the western areas seek jobs in coastal cities. They could not well care for and supervise their children, which may increase the risk of depressive symptoms among their children (Liang et al., 2017; Wang et al., 2019). Moreover, compared to other parts of China, the average family income in western area is lower (Lai, 2002). Poor family economic status could also lead to

depression among children (Eamon, 2002; Zhou et al., 2018b).

Meta-regression analysis found an increasing trend in depressive symptoms prevalence in Chinese primary school students in the past two decades, which is consistent with previous findings in other countries (Collishaw et al., 2004; Mojtabei et al., 2016; Sigfusdottir et al., 2008; Tick et al., 2007). The possible reasons may be related to the dramatic socioeconomic changes in China (Cai et al., 2017; Xin et al., 2012). Socioeconomic contexts have direct and indirect impacts on child psychosocial development (Schoon et al., 2002). In recent decades, rapid urbanization and internal migration (Mou et al., 2013; Zhou et al., 2018a) have occurred in China, which could lead to family problems, e.g., short parent-child interactions (Zhao et al., 2015), parental emotional problems (Liu and Wang, 2015; Ni et al., 2016), and increasing divorce rates (Dong et al., 2002; Liu et al., 1999; National Bureau of Statistics of China, 1998–2017). All these factors could increase the risk of depression in children. The prevalence of depressive symptoms was found to be lower in high quality studies, which may be due to the reduced study bias and lower false positive rate of depressive symptoms.

The findings of this meta-analysis should be interpreted with caution due to several limitations. First, except for one national survey with small sample size ($n = 567$) (Zhou et al., 2018b), the other studies covered only 15 out of the 34 provinces/municipalities/autonomous regions in China. Thus, the present finding could not be generalized to all primary school students in China. Second, the heterogeneity was still high, although subgroup analyses have been performed, as this is unavoidable in meta-analysis of epidemiological surveys (Patsopoulos et al., 2008). Third, due to different socio-economic backgrounds, studies conducted in Hong Kong, Macau and Taiwan were not included in this meta-analysis. Fourth, relevant factors associated with depressive symptoms, such as age, rural/urban areas, household income, parents' education level, family history of psychiatric disorders and relationships with peers and teachers, were not examined due to unavailable data in most studies.

In conclusion, given the high prevalence of depressive symptoms in Chinese primary school students, particularly in western China, and its negative health outcomes, more emphasis should be paid to improving the mental health in primary school students. Further longitudinal studies are warranted to explore the contributing factors of depressive symptoms in Chinese school children.

Funding

The study was supported by the National Science and Technology

Major Project for investigational new drug (2018ZX09201-014), and the University of Macau (MYRG2019-00066-FHS).

Contributors

Study Design: Dan-Dan Xu, Wen-Wang Rao, Yu-Tao Xiang. Analysis and interpretation of data: Dan-Dan Xu, Wen-Wang Rao, Xiao-Lan Cao, Si-Ying Wen, Feng-Rong An, Weng-lan Che. Drafting of the manuscript: Dan-Dan Xu, Daniel T. Bressington, Teris Cheung. Critical revision of the manuscript: Gabor S. Ungvari, Yu-Tao Xiang. Approval of the final version for publication: All the authors.

Declaration of Competing Interest

There is no conflict of interest related to the topic of this manuscript.

Acknowledgements

NA.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.jad.2020.02.034.

References

- Angst, J., Sellaro, R., Merikangas, K.R., 2000. Depressive spectrum diagnoses. *Compr. Psychiatry* 41, 39–47.
- Ayuso-Mateos, J.L., Nuevo, R., Verdes, E., Naidoo, N., Chatterji, S., 2010. From depressive symptoms to depressive disorders: the relevance of thresholds. *Br. J. Psychiatry* 196, 365–371.
- Birleson, P., 1981. The validity of depressive disorder in childhood and the development of a self-rating scale: a research report. *J. Child Psychol. Psychiatry* 22, 73–88.
- Birleson, P., Hudson, I., Buchanan, D.G., Wolff, S., 1987. Clinical evaluation of a self-rating scale for depressive disorder in childhood (Depression Self-Rating Scale). *J. Child Psychol. Psychiatry* 28, 43–60.
- Birmaher, B., Ryan, N.D., Williamson, D.E., Brent, D.A., Kaufman, J., 1996. Childhood and adolescent depression: a review of the past 10 years. Part II. *J. Am. Acad. Child Adolesc. Psychiatry* 35, 1575–1583.
- Borenstein, M., Hedges, L.V., Higgins, J.P., Rothstein, H.R., 2010. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res. Synth. Methods* 1, 97–111.
- Boyle, M.H., 1998. Guidelines for evaluating prevalence studies. *Evid. Based Ment. Health* 1, 37–39.
- Cai, J., Coyte, P.C., Zhao, H., 2017. Determinants of and socio-economic disparities in self-rated health in China. *Int. J. Equity Health* 16, 7.
- Calvete, E., Orue, I., Hankin, B.L., 2013. Transactional relationships among cognitive vulnerabilities, stressors, and depressive symptoms in adolescence. *J. Abnorm. Child Psych.* 41, 399–410.
- Charman, T., Pervova, I., 1996. Self-reported depressed mood in Russian and UK schoolchildren. A research note. *J. Child Psychol. Psych.* 37, 879–883.
- Cole, D.A., Jacquez, F.M., LaGrange, B., Pineda, A.Q., Truss, A.E., Weitlauf, A.S., Tilghman-Osborne, C., Felton, J., Garber, J., Dallaire, D.H., Ciesla, J.A., Maxwell, M.A., Dufton, L., 2011. A longitudinal study of cognitive risks for depressive symptoms in children and young adolescents. *J. Early Adolesc.* 31, 782–816.
- Collishaw, S., Maughan, B., Goodman, R., Pickles, A., 2004. Time trends in adolescent mental health. *J. Child Psychol. Psychiatry* 45, 1350–1362.
- Compton, W.M., Conway, K.P., Stinson, F.S., Grant, B.F., 2006. Changes in the prevalence of major depression and comorbid substance use disorders in the United States between 1991–1992 and 2001–2002. *A. J. Psychiatry* 163, 2141–2147.
- Cyranowski, J.M., Frank, E., Young, E., Shear, M.K., 2000. Adolescent onset of the gender difference in lifetime rates of major depression - a theoretical model. *Arch. Gen. Psychiat.* 57, 21–27.
- Denda, K., Kako, Y., Kitagawa, N., Koyama, T., 2006. Assessment of depressive symptoms in Japanese school children and adolescents using the Birleson Depression Self-Rating Scale. *Int. J. Psychiat. Med.* 36, 231–241.
- Ding, H., Han, J., Zhang, M., Wang, K., Gong, J., Yang, S., 2017. Moderating and mediating effects of resilience between childhood trauma and depressive symptoms in Chinese children. *J. Affect. Disord.* 211, 130–135.
- Dong, Q., Wang, Y., Ollendick, T.H., 2002. Consequences of divorce on the adjustment of children in China. *J. Clin. Child Adolesc. Psychol.* 31, 101–110.
- Eamon, M.K., 2002. Influences and mediators of the effect of poverty on young adolescent depressive symptoms. *J. Youth Adolesc.* 31, 231–242.
- Earnshaw, V.A., Elliott, M.N., Reisner, S.L., Mrug, S., Windle, M., Emery, S.T., Peskin, M.F., Schuster, M.A., 2017. Peer victimization, depressive symptoms, and substance use: a longitudinal analysis. *Pediatrics* 139. <https://doi.org/10.1542/peds.2016-3426>.
- Fan, J., Gu, H.L., Yang, H.L., Wu, L., Yi, J., Du, Y.S., 2010. Prevalence of depressive disorders among school children 8–12 years of age in the Pudong District of Shanghai (in Chinese). *Shanghai Arch. Psychiatry* 22, 335–338.
- Fellmeth, G., Rose-Clarke, K., Zhao, C., Busert, L.K., Zheng, Y., Massazza, A., Sonmez, H., Eder, B., Blewitt, A., Lertgrai, W., Orcutt, M., Ricci, K., Mohamed-Ahmed, O., Burns, R., Knipe, D., Hargreaves, S., Hesketh, T., Opondo, C., Devakumar, D., 2018. Health impacts of parental migration on left-behind children and adolescents: a systematic review and meta-analysis. *Lancet* 392, 2567–2582.
- Frigerio, A., Pesenti, S., Molteni, M., Snider, J., Battaglia, M., 2001. Depressive symptoms as measured by the CDI in a population of northern Italian children. *Eur. Psychiatry* 16, 33–37.
- Giannakopoulos, G., Kazantzi, M., Dimitrakaki, C., Tsiantis, J., Kolaitis, G., Tountas, Y., 2009. Screening for children's depression symptoms in Greece: the use of the Children's Depression Inventory in a nation-wide school-based sample. *Eur. Child Adolesc. Psychiatry* 18, 485–492.
- Global Burden of Disease 2015 Disease and Injury Incidence and Prevalence Collaborators, 2016. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388, 1545–1602.
- Gu, J.X., Guo, L.X., Wang, J.Y., 2007. Prevalence of depressive disorders among 522 children aged 8–10 years (in Chinese). *Mater. Child Health Care China* 21, 2949–2950.
- Ibrahim, A.K., Kelly, S.J., Adams, C.E., Glazebrook, C., 2013. A systematic review of studies of depression prevalence in university students. *J. Psychiatr. Res.* 47, 391–400.
- Hankin, B.L., Abramson, L.Y., Moffitt, T.E., Silva, P.A., McGee, R., Angell, K.E., 1998. Development of depression from preadolescence to young adulthood: emerging gender differences in a 10-year longitudinal study. *J. Abnorm. Psychol.* 107, 128–140.
- Higgins, J.P., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analyses. *BMJ* 327, 557–560.
- Huh, S.Y., Kim, S.G., Lee, J.S., Jung, W.Y., Choi, B.S., Kim, J.H., 2019. A study on the school violence experience of children with attention-deficit hyperactivity disorder in the context of bullying. *Asia Pac. Psychiatry* 11, e12353.
- Johnson, D., Dupuis, G., Piche, J., Clayborne, Z., Colman, I., 2018. Adult mental health outcomes of adolescent depression: a systematic review. *Depress Anxiety* 35, 700–716.
- Judd, L.L., Akiskal, H.S., 2000. Delineating the longitudinal structure of depressive illness: beyond clinical subtypes and duration thresholds. *Pharmacopsychiatry* 33, 3–7.
- Kleinman, A., 2004. Culture and depression. *N. Engl. J. Med.* 351, 951–953.
- Kovacs, M., 1992. Manual for the children's depression inventory (CDI). *Psychopharmacol. Bull.* 21, 995–998.
- Kwak, Y.S., Lee, C.I., Hong, S.C., Song, Y.J., Kim, I.C., Moon, S.H., Moon, J.H., Seok, E.M., Jang, Y.H., Park, M.J., Hong, J.Y., Kim, Y.B., Lee, S.H., Kim, H.J., Kim, M.D., 2008. Depressive symptoms in elementary school children in Jeju Island, Korea: prevalence and correlates. *Eur. Child Adolesc. Psychiatry* 17, 343–351.
- Lai, H.Y.H., 2002. China's western development program - its rationale, implementation, and prospects. *Mod. China* 28, 432–466.
- Larsson, B., Melin, L., 1992. Prevalence and short-term stability of depressive symptoms in schoolchildren. *Acta Psychiatr. Scand.* 85, 17–22.
- Lee, G., McCreary, L., Kim, M.J., Park, C.G., Jun, W.H., Yang, S., 2013. Depression in low-income elementary school children in South Korea: gender differences. *J. Sch. Nurs.* 29, 132–141.
- Lewinsohn, P.M., Rohde, P., Klein, D.N., Seeley, J.R., 1999. Natural course of adolescent major depressive disorder: I. Continuity into young adulthood. *J. Am. Acad. Child Adolesc. Psychiatry* 38, 56–63.
- Li, C., Wang, H., Cao, X.Y., Gou, M., Zhang, Z., 2013. Prevalence of depressive symptoms and its related factors among primary and middle school students in an urban-rural-intergrated area of Chongqing (in Chinese). *J. Hyg. Res.* 42, 783–788.
- Li, L., 2017. Prevalence of depressive symptoms among children with one parent working outside in Chongqing (in Chinese). *West. China Qual. Edu.* 3 (103), 157.
- Li, Y.P., Ma, G.S., Schouten, E.G., Hu, X.Q., Cui, Z.H., Wang, D., Kok, F.J., 2007. Report on childhood obesity in China (5) body weight, body dissatisfaction, and depression symptoms of Chinese children aged 9–10 years. *Biomed. Environ. Sci.* 20, 11–18.
- Liang, Y., Wang, L., Rui, G., 2017. Depression among left-behind children in China. *J. Health Psychol.* 22, 1897–1905.
- Liu, L., Wang, M., 2015. Parenting stress and children's problem behavior in China: the mediating role of parental psychological aggression. *J. Fam. Psychol.* 29, 20–28.
- Liu, S.R., Miao, R.J., 2018. Prevalence of depressive symptoms and its influencing factors among primary and middle school students in Shihezi in Xinjiang Province (in Chinese). *Occup. Health* 34, 1258–1261.
- Liu, X.C., Kurita, H., Guo, C.Q., Miyake, Y., Ze, J., Cao, H.L., 1999. Prevalence and risk factors of behavioral and emotional problems among Chinese children aged 6 through 11 years. *J. Am. Acad. Child Psych.* 38, 708–715.
- Loney, P.L., Chambers, L.W., Bennett, K.J., Roberts, J.G., Stratford, P.W., 1998. Critical appraisal of the health research literature: prevalence or incidence of a health problem. *Chronic Dis. Can.* 19, 170–176.
- Lu, Y., Hu, P., Treiman, D.J., 2012. Migration and depressive symptoms in migrant-sending areas: findings from the survey of internal migration and health in China. *Int. J. Public Health* 57, 691–698.
- Magiati, I., Ponniah, K., Ooi, Y.P., Chan, Y.H., Fung, D., Woo, B., 2015. Self-reported depression and anxiety symptoms in school-aged Singaporean children. *Asia Pac. Psychiatry* 7, 91–104.
- Ministry of Education of the People's Republic of China, 2018. Educational Statistics in

- 2018, in: Development & Planning (Ed.), China.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., Group, P., 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 6, e1000097.
- Mojtabai, R., Olfson, M., Han, B., 2016. National trends in the prevalence and treatment of depression in adolescents and young adults. *Pediatrics* 138. <https://doi.org/10.1542/peds.2016-1878>.
- Mou, J., Griffiths, S.M., Fong, H., Dawes, M.G., 2013. Health of China's rural-urban migrants and their families: a review of literature from 2000 to 2012. *Br. Med. Bull.* 106, 19–43.
- Naicker, K., Galambos, N.L., Zeng, Y., Senthilselvan, A., Colman, I., 2013. Social, demographic, and health outcomes in the 10 years following adolescent depression. *J. Adolesc. Health* 52, 533–538.
- National Bureau of Statistics of China, 1998–2017. Crude Divorce Rate, 1998–2017. <http://data.stats.gov.cn/>.
- Ni, M.Y., Jiang, C., Cheng, K.K., Zhang, W., Gilman, S.E., Lam, T.H., Leung, G.M., Schooling, C.M., 2016. Stress across the life course and depression in a rapidly developing population: the Guangzhou biobank cohort study. *Int. J. Geriatr. Psychiatry* 31, 629–637.
- Patsopoulos, N.A., Evangelou, E., Ioannidis, J.P., 2008. Sensitivity of between-study heterogeneity in meta-analysis: proposed metrics and empirical evaluation. *Int. J. Epidemiol.* 37, 1148–1157.
- Patton, G.C., Viner, R., 2007. Pubertal transitions in health. *Lancet* 369, 1130–1139.
- Peng, L.L., He, F., Yang, J.W., Ran, M., Wang, H., 2018. Relationships between pubertal timing and depressive symptoms among primary and junior middle school students in urban areas of Chongqing (in Chinese). *Chin. J. Sch. Health* 39, 215–218.
- Qi, Y.L., 2006. Investigation of depressive emotion and related risk factors in 1018 pupils in Jilin (in Chinese). *J. Beihua Univ. (Natural Science)* 7, 452–454.
- Radloff, L.S., 1977. The CES-D scale: a self-report depression scale for research in the general populations. *Appl. Psych. Meas.* 1, 385–401.
- Rao, W.-W., Xu, D.-D., Cao, X.-L., Wen, S.-Y., Che, W.-L., Ng, C.H., Ungvari, G.S., He, F., Xiang, Y.-T., 2019. Prevalence of depressive symptoms in children and adolescents in China: a meta-analysis of observational studies. *Psychiatry Res.* 272, 790–796.
- Ren, F., Yu, X., Dang, W., Niu, W., Zhou, T., Lin, Y., Wu, Z., Lin, L., Zhong, B., Chu, H., Zhou, J., Ding, H., Yuan, P., 2019. Depressive symptoms in Chinese assembly-line migrant workers: a case study in the shoe-making industry. *Asia Pac. Psychiatry* 11, e12332.
- Rutter, M., Kim-Cohen, J., Maughan, B., 2006. Continuities and discontinuities in psychopathology between childhood and adult life. *J. Child Psychol. Psychiatry* 47, 276–295.
- Schoon, I., Bynner, J., Joshi, H., Parsons, S., Wiggins, R.D., Sacker, A., 2002. The influence of context, timing, and duration of risk experiences for the passage from childhood to midadulthood. *Child Dev.* 73, 1486–1504.
- Sigfusdottir, I.D., Asgeirsdottir, B.B., Sigurdsson, J.F., Gudjonsson, G.H., 2008. Trends in depressive symptoms, anxiety symptoms and visits to healthcare specialists: a national study among Icelandic adolescents. *Scand. J. Public Health* 36, 361–368.
- Stroup, D.F., Berlin, J.A., Morton, S.C., Olkin, I., Williamson, G.D., Rennie, D., Moher, D., Becker, B.J., Sipe, T.A., Thacker, S.B., 2000. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group. *JAMA* 283, 2008–2012.
- Sun, L., Zhou, T.H., 2005. The survey of depressive symptoms in children and adolescents aged 8–15 years (in Chinese). *Chin. J. Behav. Med. Sci.* 14, 58.
- Tang, J., Su, L.Y., Zu, Y., Wang, K., Luo, X.R., 2003. Depressive symptoms in Chinese primary students (in Chinese). *Chin. J. Clin. Psychol.* 11, 246–266.
- Tang, X., Tang, S., Ren, Z., Wong, D.F.K., 2018. Prevalence of depressive symptoms among adolescents in secondary school in mainland China: a systematic review and meta-analysis. *J. Affect. Disord.* 245, 498–507.
- Tepper, P., Liu, X., Guo, C., Zhai, J., Liu, T., Li, C., 2008. Depressive symptoms in Chinese children and adolescents: parent, teacher, and self reports. *J. Affect. Disord.* 111, 291–298.
- Teri, L., 1982. The use of the beck depression inventory with adolescents. *J. Abnorm. Child Psych.* 10, 277–284.
- Tick, N.T., van der Ende, J., Verhulst, F.C., 2007. Twenty-year trends in emotional and behavioral problems in Dutch children in a changing society. *Acta Psychiatr. Scand.* 116, 473–482.
- Verboom, C.E., Sijtsma, J.J., Verhulst, F.C., Penninx, B.W., Ormel, J., 2014. Longitudinal associations between depressive problems, academic performance, and social functioning in adolescent boys and girls. *Dev. Psychol.* 50, 247–257.
- Wang, H.Z., Yang, M.L., Lou, X.M., Zhou, Z.H., Wang, R.J., Wang, X., 2015. Prevalence of depressive symptoms among primary and middle school students in Zhengzhou (in Chinese). *Chin. J. Public Health* 31, 1005–1007.
- Wang, K., Su, L.Y., 2005. Two-years follow-up study of depressive symptoms in children aged 8–10 years (in Chinese). *Chin. J. Child Health Care* 13, 369–371.
- Wang, L., Feng, Z., Yang, G., Yang, Y., Wang, K., Dai, Q., Zhao, M., Hu, C., Zhang, R., Liu, K., Guang, Y., Xia, F., 2016. Depressive symptoms among children and adolescents in western china: an epidemiological survey of prevalence and correlates. *Psychiatry Res.* 246, 267–274.
- Wang, Y.Y., Xiao, L., Rao, W.W., Chai, J.X., Zhang, S.F., Ng, C.H., Ungvari, G.S., Zhu, H., Xiang, Y.T., 2019. The prevalence of depressive symptoms in 'left-behind children' in China: a meta-analysis of comparative studies and epidemiological surveys. *J. Affect. Disord.* 244, 209–216.
- Wu, H.N., Shao, F., Lin, B., 2017. Risk factors related to depressive symptoms among adolescents and the prevention strategies (in Chinese). *Mod. Pract. Med.* 29, 670–672.
- Wu, L.J., Wei, D.M., Gao, A.Y., Li, Q., Li, X.H., Lin, S.T., Wang, H.J., 2015. Association between physical activity, screen-based media use and depressive symptoms among children in Beijing (in Chinese). *Chin. J. Sch. Health* 36, 326–329.
- Xie, S., Yu, X.M., Wang, Y.L., Yang, S.B., Liu, W.W., Ding, H.S., Hu, Y., Xu, Y.H., Han, J., 2016. Relationships between depression and stressful events in adolescence (in Chinese). *Chin. J. Public Health* 32, 1680–1683.
- Xin, Z., Niu, J., Chi, L., 2012. Birth cohort changes in Chinese adolescents' mental health. *Int. J. Psychol.* 47, 287–295.
- Xu, D.D., Rao, W.W., Cao, X.L., Wen, S.Y., Che, W.L., Ng, C.H., Ungvari, G.S., Du, Y., Zhang, L., Xiang, Y.T., 2018. Prevalence of major depressive disorder in children and adolescents in China: a systematic review and meta-analysis. *J. Affect. Disord.* 241, 592–598.
- Xu, J., Lin, D.N., Wang, J.J., Hu, W., Zhang, H.B., Xu, G., 2008. Comparison of influential factors for depressive symptoms among primary school students in Hefei and Shenzhen (in Chinese). *Chin. Mental Health J.* 22, 246–248 + 252.
- Yan, M., 2012. The relationships between depression and growth and development in adolescents aged 9–17 years in Wuhan (in Chinese/Thesis). Huazhong University of Science and Technology.
- Yang, J., Chen, K.Y., Chen, Y.H., Huang, C., Wang, Z.Y., Ye, X.Q., Li, X., Li, S., Liang, W.D., Qiu, Y.S., 2014. Prevalence of depressive symptoms and the influencing factors among primary school students in Shenzhen (in Chinese). *Med. Inf.* 27, 183–184.
- Yong, N., Wang, H., Hu, H., Meng, H.Q., Chen, P.H., Du, L., Qu, Y., Zou, Z.L., Wang, T., Zhao, W.J., Liu, H.X., Chen, X., Fu, Y.X., Luo, Q.H., Qiu, H.T., Qiu, T., 2011. Relationships between child-abuse, depressive symptoms and sleep quality in grade 4th–6th primary school students (in Chinese). *Chin. Mental Health J.* 25, 616–621.
- Zhang, X.Y., Zhou, W.J., 2015. Depressive symptoms and related behavioral factors in pupils in Yancheng (in Chinese). *Chin. J. Soc. Med.* 32, 192–194.
- Zhao, J., Liu, X., Wang, M., 2015. Parent-child cohesion, friend companionship and left-behind children's emotional adaptation in rural China. *Child Abuse Negl.* 48, 190–199.
- Zhao, X., 2014. Depressive symptoms and the related factors among children in rural areas of Anhui Province: a two-year longitudinal study (in Chinese/Thesis). Anhui Med. Univ.
- Zhou, M., Sun, X., Huang, L., Zhang, G., Kenny, K., Xue, H., Auden, E., Rozelle, S., 2018a. Parental migration and left-behind children's depressive symptoms: estimation based on a nationally-representative panel dataset. *Int. J. Environ. Res. Public Health* 15. <https://doi.org/10.3390/ijerph15061069>.
- Zhou, Q., Fan, L., Yin, Z., 2018b. Association between family socioeconomic status and depressive symptoms among Chinese adolescents: evidence from a national household survey. *Psychiatry Res.* 259, 81–88.
- Zhou, Z., Wang, J., 2011. Depressive symptoms and its associated factors among primary and middle school students in rural areas (in Chinese). *Acta Univ. Med. Anhui* 46, 976–978.
- Zhu, H.Q., Li, Q., Wang, L.Q., Zhang, R., Wang, J.H., 2016. Prevalence of depressive symptoms and its influencing factors among primary and middle school students in Haikou (in Chinese). *Chin. J. Sch. Health* 37, 1345–1347 + 1350.
- Zhu, J.H., Zhong, B.L., Xu, L.J., 2013. Depressive symptoms among primary and middle school students aged 7–17 years in Wuhan: an epidemiological survey (in Chinese). *Chin. J. Brain Dis. Rehabil.* 3, 35–39.