

# An introduction to fractional calculus

Fundamental ideas and numerics

Fabio Durastante

Università di Pisa

✉ [fabio.durastante@unipi.it](mailto:fabio.durastante@unipi.it)

🌐 [fdurastante.github.io](https://fdurastante.github.io)

October, 2022



# Nonlocal operators (Andreu-Vaillo et al. 2010)

---

Let  $\Omega \subset \mathbb{R}^n$  denote a **bounded** and **open** domain.

The **action** of a **nonlocal diffusion** operator  $\mathcal{L}$  on  $u(\mathbf{x}) : \Omega \rightarrow \mathbb{R}$  is defined as

$$\mathcal{L}u(\mathbf{x}) = 2 \int_{\mathbb{R}^n} (u(\mathbf{y}) - u(\mathbf{x}))\gamma(\mathbf{x}, \mathbf{y}) \, d\mathbf{y}, \quad \forall \mathbf{x} \in \Omega \subseteq \mathbb{R}^n.$$

- ⚙ the *volume*  $\Omega$  is non-zero,
- ⚙ the *kernel*  $\gamma(\mathbf{x}, \mathbf{y}) : \Omega \times \Omega \rightarrow \mathbb{R}$  is **nonnegative** and **symmetric**.

# Nonlocal operators (Andreu-Vaillo et al. 2010)

Let  $\Omega \subset \mathbb{R}^n$  denote a **bounded** and **open** domain.

The **action** of a **nonlocal diffusion** operator  $\mathcal{L}$  on  $u(\mathbf{x}) : \Omega \rightarrow \mathbb{R}$  is defined as

$$\mathcal{L}u(\mathbf{x}) = 2 \int_{\mathbb{R}^n} (u(\mathbf{y}) - u(\mathbf{x}))\gamma(\mathbf{x}, \mathbf{y}) \, d\mathbf{y}, \quad \forall \mathbf{x} \in \Omega \subseteq \mathbb{R}^n.$$

- ⚙ the *volume*  $\Omega$  is non-zero,
- ⚙ the *kernel*  $\gamma(\mathbf{x}, \mathbf{y}) : \Omega \times \Omega \rightarrow \mathbb{R}$  is **nonnegative** and **symmetric**.

The first **interesting equation** is the *nonlocal steady-state*

$$\begin{cases} -\mathcal{L}u = f, & \text{on } \Omega, \\ u = 0, & \text{on } \Omega_{\mathcal{I}}, \end{cases}$$

- 👁 the **equality constraint** should be defined in general on an *interaction volume*  $\Omega_{\mathcal{I}}$  that is **disjoint** from  $\Omega$ ; typically  $\Omega_{\mathcal{I}} = \mathbb{R}^n \setminus \Omega \equiv \Omega^c$ .

# Fractional Laplacian

---

We are interested in a particular *nonlocal* operator  $\mathcal{L}$  called the **Fractional Laplacian**.

# Fractional Laplacian

---

We are interested in a particular *nonlocal* operator  $\mathcal{L}$  called the **Fractional Laplacian**.

## Fractional Laplacian

The fractional Laplacian is the pseudo-differential operator with Fourier symbol  $\mathfrak{F}$  satisfying

$$(-\Delta)^\alpha u(\xi) = |\xi|^{2\alpha} \hat{u}(\xi), \quad 0 < \alpha \leq 1,$$

where  $\hat{u}$  denotes the *Fourier transform* of  $u$ .

# Fractional Laplacian

We are interested in a particular *nonlocal* operator  $\mathcal{L}$  called the **Fractional Laplacian**.

## Fractional Laplacian

The fractional Laplacian is the pseudo-differential operator with Fourier symbol  $\mathfrak{F}$  satisfying

$$(-\Delta)^\alpha u(\xi) = |\xi|^{2\alpha} \hat{u}(\xi), \quad 0 < \alpha \leq 1,$$

where  $\hat{u}$  denotes the *Fourier transform* of  $u$ .

## Fractional Laplacian: integral formulation

An equivalent characterization of the fractional Laplacian is given by

$$(-\Delta)^\alpha u = c_{n,\alpha} \int_{\mathbb{R}^n} \frac{u(\mathbf{x}) - u(\mathbf{y})}{|\mathbf{y} - \mathbf{x}|^{n+2\alpha}} d\mathbf{y}, \quad 0 < \alpha < 1, \quad c_{n,\alpha} = \alpha 2^{2\alpha} \frac{\Gamma((n+2)/2)}{\Gamma(1/2)\Gamma(1-\alpha)}.$$

# Fractional Laplacian

We are interested in a particular *nonlocal* operator  $\mathcal{L}$  called the **Fractional Laplacian**.

## Fractional Laplacian

The fractional Laplacian is the pseudo-differential operator with Fourier symbol  $\mathfrak{F}$  satisfying

$$(-\Delta)^\alpha u(\xi) = |\xi|^{2\alpha} \hat{u}(\xi), \quad 0 < \alpha \leq 1,$$

where  $\hat{u}$  denotes the *Fourier transform* of  $u$ .

## Fractional Laplacian: integral formulation

An equivalent characterization of the fractional Laplacian is given by

$$-\mathcal{L} = (-\Delta)^\alpha, \quad 0 < \alpha < 1, \quad \gamma(\mathbf{x}, \mathbf{y}) = \frac{c_{n,\alpha}}{2|\mathbf{y} - \mathbf{x}|^{n+2\alpha}} \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

# Fractional Laplacian

We are interested in a particular *nonlocal* operator  $\mathcal{L}$  called the **Fractional Laplacian**.

## Fractional Laplacian

The fractional Laplacian is the pseudo-differential operator with Fourier symbol  $\mathfrak{F}$  satisfying

$$(-\Delta)^\alpha u(\xi) = |\xi|^{2\alpha} \hat{u}(\xi), \quad 0 < \alpha \leq 1,$$

where  $\hat{u}$  denotes the *Fourier transform* of  $u$ .

## Fractional Laplacian: integral formulation

An equivalent characterization of the fractional Laplacian is given by

$$-\mathcal{L} = (-\Delta)^\alpha, \quad 0 < \alpha < 1, \quad \gamma(\mathbf{x}, \mathbf{y}) = \frac{c_{n,\alpha}}{2|\mathbf{y} - \mathbf{x}|^{n+2\alpha}} \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

🔧 We can play around with the definitions...



# Fractional Laplacian (10 equivalent definitions)

☞ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(a) Fourier definition:

$$\mathcal{F}(\mathcal{L}f)(\xi) = -|\xi|^\beta \mathcal{F}f(\xi)$$

(if  $\mathcal{X} = \mathbb{L}^p$ ,  $p \in [1, 2]$ );

(b) distributional definition:

$$\int_{\mathbb{R}^d} \mathcal{L}f(y) \varphi(y) dy = \int_{\mathbb{R}^d} f(x) L\varphi(x) dx$$

for all Schwartz functions  $\varphi$ , with  $\mathcal{L}\varphi$  defined, for example, as in (a);

# Fractional Laplacian (10 equivalent definitions)

☰ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(c) Bochner's<sup>1</sup> definition:

$$\mathcal{L}f = \frac{1}{|\Gamma(-\frac{\beta}{2})|} \int_0^\infty (e^{t\Delta} f - f) t^{-1-\beta/2} dt,$$

with the Bochner's integral of an  $\mathcal{X}$ -valued function;

---

<sup>1</sup>Bochner's integral extends the definition of Lebesgue integral to functions that take values in a Banach space, as the limit of integrals of simple functions.

# Fractional Laplacian (10 equivalent definitions)

☞ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(d) Balakrishnan's definition:

$$\mathcal{L}f = \frac{\sin \frac{\beta\pi}{2}}{\pi} \int_0^\infty \Delta(sI - \Delta)^{-1} f s^{\beta/2-1} ds,$$

(e) singular integral definition:

$$\mathcal{L}f = \lim_{r \rightarrow 0^+} \frac{2^\beta \Gamma(\frac{d+\beta}{2})}{\pi^{d/2} |\Gamma(-\frac{\beta}{2})|} \int_{\mathbb{R}^d \setminus B(x,r)} \frac{f(\cdot + z) - f(\cdot)}{|z|^{d+\beta}} dz,$$

with the limit in  $\mathcal{X}$ ;

# Fractional Laplacian (10 equivalent definitions)

☞ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(f) Dynkin's definition:

$$\mathcal{L}f = \lim_{r \rightarrow 0^+} \frac{2^\beta \Gamma(\frac{d+\beta}{2})}{\pi^{d/2} |\Gamma(-\frac{\beta}{2})|} \int_{\mathbb{R}^d \setminus \overline{B}(x,r)} \frac{f(\cdot + z) - f(\cdot)}{|z|^d (|z|^2 - r^2)^{\beta/2}} dz,$$

with the limit in  $\mathcal{X}$ ;

# Fractional Laplacian (10 equivalent definitions)

☰ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(g) quadratic form definition:  $\langle \mathcal{L}f, \varphi \rangle = \mathcal{E}(f, \varphi)$  for all  $\varphi$  in the Sobolev space  $H^{\beta/2}$ , where

$$\mathcal{E}(f, g) = \frac{2^\beta \Gamma(\frac{d+\beta}{2})}{2\pi^{d/2} |\Gamma(-\frac{\beta}{2})|} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(f(y) - f(x))(\overline{g(y)} - \overline{g(x)})}{|x - y|^{d+\beta}} dx dy$$

(if  $\mathcal{X} = \mathbb{L}^2$ );

# Fractional Laplacian (10 equivalent definitions)

☞ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(h) semigroup definition:

$$\mathcal{L}f = \lim_{t \rightarrow 0^+} \frac{P_t f - f}{t},$$

where  $P_t f = f * p_t$  and  $\mathcal{F}p_t(\xi) = e^{-t|\xi|^\beta}$ ;

(i) definition as the inverse of the Riesz potential:

$$\frac{\Gamma(\frac{d-\beta}{2})}{2^\beta \pi^{d/2} \Gamma(\frac{\beta}{2})} \int_{\mathbb{R}^d} \frac{\mathcal{L}f(\cdot + z)}{|z|^{d-\beta}} dz = -f(\cdot)$$

(if  $\beta < d$  and  $\mathcal{X} = \mathbb{L}^p$ ,  $p \in [1, \frac{d}{\beta})$ );

# Fractional Laplacian (10 equivalent definitions)

☰ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent:

(j) definition through harmonic extensions:

$$\begin{cases} \Delta_x u(x, y) + \beta^2 c_\beta^{2/\beta} y^{2-2/\beta} \partial_y^2 u(x, y) = 0 & \text{for } y > 0, \\ u(x, 0) = f(x), \\ \partial_y u(x, 0) = \mathcal{L}f(x), \end{cases}$$

where  $c_\beta = 2^{-\beta} |\Gamma(-\frac{\beta}{2})| / \Gamma(\frac{\beta}{2})$  and where  $u(\cdot, y)$  is a function of class  $\mathcal{X}$  which depends continuously on  $y \in [0, \infty)$  and  $\|u(\cdot, y)\|_{\mathcal{X}}$  is bounded in  $y \in [0, \infty)$ .

# Fractional Laplacian (10 equivalent definitions)

☰ We denote by  $\mathbb{L}^p$  ( $p \in [1, \infty)$ ) the Lebesgue spaces,  $\mathcal{C}_0$  the space of continuous functions vanishing at infinity, and with  $\mathcal{C}_{bu}$  the space of bounded uniformly continuous functions.

## Theorem (Kwaśnicki 2017, Theorem 1.1)

Let  $\mathcal{X}$  be any of the spaces  $\mathbb{L}^p$ ,  $p \in [1, \infty)$ ,  $\mathcal{C}_0$  or  $\mathcal{C}_{bu}$ , and let  $f \in \mathcal{X}$ ,  $\beta = 2\alpha$ . The following definitions of  $\mathcal{L}f \in \mathcal{X}$  are equivalent.

In addition, in (c), (e), (f), (h) and (j), convergence in the uniform norm can be relaxed to pointwise convergence to a function in  $\mathcal{X}$  when  $\mathcal{X} = \mathcal{C}_0$  or  $\mathcal{X} = \mathcal{C}_{bu}$ . Finally, for  $\mathcal{X} = \mathbb{L}^p$  with  $p \in [1, \infty)$ , norm convergence in (e), (f), (h) or (j) implies pointwise convergence for almost all  $x$ .

⚙ Convergence properties described here are for the *full-space definitions* of the fractional Laplace operator  $\mathcal{L}$ .

💡 We can invent **numerical methods** starting from **each of these definitions**.



# Fractional Laplacian: equations on bounded domains

---

If  $\Omega$  is **bounded** we can modify our first definition as follows.

🔧 Take  $u : \Omega \rightarrow \mathbb{R}$  and extend it to zero outside of  $\Omega$ :

$$(-\Delta)^\alpha \tilde{u} = f \text{ in } \Omega, \quad \tilde{u} = 0 \text{ in } \Omega^c = \mathbb{R}^n \setminus \Omega.$$

where

$$(-\Delta)^\alpha \tilde{u} = c_{n,\alpha} \int_{\mathbb{R}^n} \frac{\tilde{u}(\mathbf{x}) - \tilde{u}(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|^{n+2s}} d\mathbf{y}$$

and thus  $\tilde{u}$  is the extension by zero to  $\mathbb{R}^n$  of a function  $u : \Omega \rightarrow \mathbb{R}$  in  $\mathbb{L}^2(\Omega)$ .

# Fractional Laplacian: equations on bounded domains

If  $\Omega$  is **bounded** we can modify our first definition as follows.

🔧 Take  $u : \Omega \rightarrow \mathbb{R}$  and extend it to zero outside of  $\Omega$ :

$$(-\Delta)^\alpha \tilde{u} = f \text{ in } \Omega, \quad \tilde{u} = 0 \text{ in } \Omega^c = \mathbb{R}^n \setminus \Omega.$$

where

$$(-\Delta)^\alpha \tilde{u} = c_{n,\alpha} \int_{\mathbb{R}^n} \frac{\tilde{u}(\mathbf{x}) - \tilde{u}(\mathbf{y})}{|\mathbf{x} - \mathbf{y}|^{n+2s}} d\mathbf{y}$$

and thus  $\tilde{u}$  is the extension by zero to  $\mathbb{R}^n$  of a function  $u : \Omega \rightarrow \mathbb{R}$  in  $\mathbb{L}^2(\Omega)$ .

## 🚶 Stochastic interpretation.

As we have seen when discussing the other derivatives, we can interpret also the Fractional Laplacian in a stochastic way. Indeed, one can prove that it is the infinitesimal generator of a **2 $\alpha$ -stable Lévy process**. The **boundary conditions** means that the particles are killed upon reaching  $\Omega^c$ .

# Fractional Laplacian: equations on bounded domains

---

The second definition relies instead on **spectral theory**.

- ⚙ Recall that  $-\Delta : \mathcal{D}(-\Delta) \subset \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)$  is an unbounded, positive and closed operator with dense domain  $\mathcal{D}(-\Delta) = \mathbb{H}_0^1(\Omega) \cap \mathbb{H}^2(\Omega)$  with a compact inverse.

# Fractional Laplacian: equations on bounded domains

---

The second definition relies instead on **spectral theory**.

- ⚙ Recall that  $-\Delta : \mathcal{D}(-\Delta) \subset \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)$  is an unbounded, positive and closed operator with dense domain  $\mathcal{D}(-\Delta) = \mathbb{H}_0^1(\Omega) \cap \mathbb{H}^2(\Omega)$  with a compact inverse.
- 📖 There is a *countable* collection of eigenpairs  $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^+ \times \mathbb{H}_0^1(\Omega)$  such that  $\{\varphi_k\}_{k \in \mathbb{N}}$  is an **orthonormal basis** of  $\mathbb{L}^2(\Omega)$  (and of  $\mathbb{H}_0^1(\Omega)$ ).

# Fractional Laplacian: equations on bounded domains

The second definition relies instead on **spectral theory**.

- ⚙ Recall that  $-\Delta : \mathcal{D}(-\Delta) \subset \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)$  is an unbounded, positive and closed operator with dense domain  $\mathcal{D}(-\Delta) = \mathbb{H}_0^1(\Omega) \cap \mathbb{H}^2(\Omega)$  with a compact inverse.
- 📖 There is a *countable* collection of eigenpairs  $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^+ \times \mathbb{H}_0^1(\Omega)$  such that  $\{\varphi_k\}_{k \in \mathbb{N}}$  is an **orthonormal basis** of  $\mathbb{L}^2(\Omega)$  (and of  $\mathbb{H}_0^1(\Omega)$ ).
- 🔧 The **fractional power of the Dirichlet Laplacian** can thus be defined  $\forall u \in \mathcal{C}_0^\infty$  as

$$(-\Delta)^\alpha u = \sum_{k=1}^{+\infty} \lambda_k^\alpha u_k \varphi_k, \quad u_k = \langle w, \varphi_k \rangle_{\mathbb{L}^2(\Omega)} = \int_{\Omega} w \varphi_k \, dx, \quad k \in \mathbb{N}$$

# Fractional Laplacian: equations on bounded domains

The second definition relies instead on **spectral theory**.

- ⚙ Recall that  $-\Delta : \mathcal{D}(-\Delta) \subset \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)$  is an unbounded, positive and closed operator with dense domain  $\mathcal{D}(-\Delta) = \mathbb{H}_0^1(\Omega) \cap \mathbb{H}^2(\Omega)$  with a compact inverse.
- 📖 There is a *countable* collection of eigenpairs  $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^+ \times \mathbb{H}_0^1(\Omega)$  such that  $\{\varphi_k\}_{k \in \mathbb{N}}$  is an **orthonormal basis** of  $\mathbb{L}^2(\Omega)$  (and of  $\mathbb{H}_0^1(\Omega)$ ).
- 🔧 The **fractional power of the Dirichlet Laplacian** can thus be defined  $\forall u \in \mathcal{C}_0^\infty$  as

$$(-\Delta)^\alpha u = \sum_{k=1}^{+\infty} \lambda_k^\alpha u_k \varphi_k, \quad u_k = \langle w, \varphi_k \rangle_{\mathbb{L}^2(\Omega)} = \int_{\Omega} w \varphi_k \, dx, \quad k \in \mathbb{N}$$

## Extension

This definition of  $(-\Delta)^\alpha$  can be extended by density to

$$\mathbb{H}^\alpha(\Omega) = \left\{ w = \sum_{k=1}^{+\infty} w_k \varphi_k : \sum_{k=1}^{+\infty} \lambda_k^\alpha w_k^2 < +\infty \right\}.$$

# 🤖 definitions on bounded domains aren't equivalent!

The **integral definition** of the Fractional Laplacian in

$$(-\Delta)^\alpha \tilde{u} = f \text{ in } \Omega, \quad \tilde{u} = 0 \text{ in } \Omega^c = \mathbb{R}^n \setminus \Omega,$$

and the **spectral definition**

$$(-\Delta)^\alpha u = \sum_{k=1}^{+\infty} \lambda_k^\alpha u_k \varphi_k, \quad u_k = \langle w, \varphi_k \rangle_{\mathbb{L}^2(\Omega)} = \int_{\Omega} w \varphi_k \, dx, \quad k \in \mathbb{N},$$

are **NOT EQUIVALENT!**

## Differences

Their difference is **positive** and **positivity preserving** (Musina and Nazarov 2014, Theorems 1 and 2). Furthermore, if we call  $d(x, \partial\Omega)$  the distance for  $x \in \Omega$  to the boundary  $\partial\Omega$  we find

$$(\text{integral}) \, u(x) \approx d(x, \partial\Omega)^\alpha + v(x), \quad (\text{spectral}) \, u(x) \approx \begin{cases} d(x, \partial\Omega)^{2\alpha} + v(x), & \alpha \in (0, 1/2), \\ d(x, \partial\Omega) + v(x), & \alpha \in (1/2, 1), \end{cases}$$

for a smooth  $v(x)$ .

## Equations of interest

---

Selecting the **right definition** for the problem the setting one has in mind (finite domain, infinite domain, ...) we can formulate several PDE with this new operator.

Diffusion-reaction  $\partial_t u + (-\Delta)^\alpha u + c(t, x)u = 0$ , Domain  $(0, +\infty) \times \mathbb{R}^n$ ,

Quasi-geostrophic  $\partial_t \theta + u \cdot \nabla \theta + \kappa(-\Delta)^\alpha \theta = f$ , Domain  $[0, T] \times \mathbb{R}^2$ ,

Cahn-Hilliard  $\partial_t u + (-\Delta)^\alpha (-\varepsilon^2 \Delta u + f(u)) = 0$ , Domain  $(0, T] \times (0, 2\pi)^2$ ,

Porous medium  $\partial_t u + (-\Delta)^\alpha (|u|^{m-1} \text{sign}(u)) = 0$ , Domain  $(0, +\infty) \times \mathbb{R}^n$ ,

Schrödinger  $i\hbar \partial_t \psi = D_\alpha (-\hbar^2 \Delta)^\alpha \psi + V(r, t)\psi$ , Domain  $(r, t) \in \mathbb{R}^3 \times (0, +\infty)$ ,

Ultrasound  $c_0^{-2} \partial_t^2 p = \nabla^2 p - \{\tau \partial_t (-\Delta)^\alpha + \eta (-\Delta)^{\alpha+1/2}\} p$ , Domain  $(-\infty, +\infty) \times \mathbb{R}^n$ .

👁 See **the review** (Lischke et al. [2020](#)) for an updated *list of references*.



# The Spectral Fractional Laplacian

---

Let us focus on problem using the **spectral Fractional Laplacian**

$$(-\Delta)^\alpha u = \sum_{k=1}^{+\infty} \lambda_k^\alpha u_k \varphi_k, \quad u_k = \langle w, \varphi_k \rangle_{\mathbb{L}^2(\Omega)} = \int_{\Omega} w \varphi_k \, dx, \quad k \in \mathbb{N}.$$

❓ How can we obtain **reliable numerical methods**?

# The Spectral Fractional Laplacian

Let us focus on problem using the **spectral Fractional Laplacian**

$$(-\Delta)^\alpha u = \sum_{k=1}^{+\infty} \lambda_k^\alpha u_k \varphi_k, \quad u_k = \langle w, \varphi_k \rangle_{\mathbb{L}^2(\Omega)} = \int_{\Omega} w \varphi_k \, dx, \quad k \in \mathbb{N}.$$

❓ How can we obtain **reliable numerical methods**?

## 💡 The Matrix-Transfer Technique

The idea from (Ilic et al. 2005, 2006) goes as follows, suppose that we have a *discretization scheme* for  $-\Delta$  on  $\Omega$ . That is, we can build  $A_n = -\Delta_h \approx -\Delta$  on a discrete  $\Omega_h$  ( $h \rightarrow 0$  for  $n \rightarrow +\infty$ ), then:

$$(-\Delta)^\alpha \approx (-\Delta_h)^\alpha = A_n^\alpha,$$

i.e., we have to compute a **matrix function** of (sparse) matrix discretizing the ordinary Laplacian on the domain of interest.

# The Finite Difference Example

---

The simplest example we can think of is using **finite differences** on  $\Omega = [0, 1]$  to solve for

$$\begin{cases} (-\Delta)^\alpha u = f(x), & x \in (0, 1), \\ u(0) = u(1) = 0. \end{cases}$$

This can be rewritten as

$$A_n = \frac{1}{h^2} T_{n-2}(2 - 2\cos(\theta)), \quad h = \frac{1}{n-1},$$

on the grid  $\{x_j = jh\}_{j=0}^n$ , and solved on the inner nodes

$$\mathbf{u}_n(2:n-1) = A_n^{-\alpha} \mathbf{f}(2:n-1),$$

via *diagonalization*.

# The Finite Difference Example

The simplest example we can think of is using **finite differences** on  $\Omega = [0, 1]$  to solve for

$$\begin{cases} (-\Delta)^\alpha u = f(x), & x \in (0, 1), \\ u(0) = u(1) = 0. \end{cases}$$

This can be rewritten as

$$A_n = \frac{1}{h^2} T_{n-2}(2 - 2\cos(\theta)), \quad h = \frac{1}{n-1},$$

on the grid  $\{x_j = jh\}_{j=0}^n$ , and solved on the inner nodes

$$\mathbf{u}_n(2:n-1) = A_n^{-\alpha} \mathbf{f}(2:n-1),$$

via *diagonalization*.

```
n = 100; h = 1/(n-1);  
x = linspace(0,1,n)';  
e = ones(n-2,1);  
An = spdiags([-e,2*e,-e]/h^2,-1:1,  
    ↪ n-2,n-2);  
f = sin(pi*x);  
u = [0;An\f(2:n-1);0];  
[U,L,V] = eig(full(An));  
ualpha = @(alpha)  
    ↪ [0;V'\'(L.^alpha\'(U\f(2:n-1))))];
```

# The Finite Difference Example

The simplest example we can think of is using **finite differences** on  $\Omega = [0, 1]$  to solve for

$$\begin{cases} (-\Delta)^\alpha u = f(x), & x \in (0, 1), \\ u(0) = u(1) = 0. \end{cases}$$


This can be rewritten as

$$A_n = \frac{1}{h^2} T_{n-2}(2 - 2\cos(\theta)), \quad h = \frac{1}{n-1},$$

on the grid  $\{x_j = jh\}_{j=0}^n$ , and solved on the inner nodes

$$\mathbf{u}_n(2:n-1) = A_n^{-\alpha} \mathbf{f}(2:n-1),$$

via *diagonalization*.



# The general case

---

😡 Somebody usually gets angry if we start diagonalizing stuff...

# The general case

---

😡 Somebody usually gets angry if we start diagonalizing stuff... so the right way to do is going for a *matrix function times vector* computation.

- ⚙️ We need to compute  $g(z) = z^{-\alpha}$  for  $\alpha \in (0, 1)$ ,
- ⚙️ on a matrix  $A_n$  that is either **symmetric and positive definite**, or of a matrix that is *similar* to an SPD matrix,
- ⚙️  $A_n$  has also a condition number that grows (at least quadratically) with its size, i.e., is **ill-conditioned**.

# The general case

---

😡 Somebody usually gets angry if we start diagonalizing stuff... so the right way to do is going for a *matrix function times vector* computation.

⚙️ We need to compute  $g(z) = z^{-\alpha}$  for  $\alpha \in (0, 1)$ ,

⚙️ on a matrix  $A_n$  that is either **symmetric and positive definite**, or of a matrix that is *similar* to an SPD matrix,

⚙️  $A_n$  has also a condition number that grows (at least quadratically) with its size, i.e., is **ill-conditioned**.

❓ What method do we select?



# The general case

---

😡 Somebody usually gets angry if we start diagonalizing stuff... so the right way to do is going for a *matrix function times vector* computation.

⚙️ We need to compute  $g(z) = z^{-\alpha}$  for  $\alpha \in (0, 1)$ ,

⚙️ on a matrix  $A_n$  that is either **symmetric and positive definite**, or of a matrix that is *similar* to an SPD matrix,

⚙️  $A_n$  has also a condition number that grows (at least quadratically) with its size, i.e., is **ill-conditioned**.

❓ What method do we select?

🔧  $A_n$  is sparse and, if we deal with a regular uniform grid maybe also Toeplitz, a Lanczos **polynomial Krylov** with fast convergence would be perfect if it reaches convergence with a number of iteration independent of the size  $n$ .

# The general case

---

😡 Somebody usually gets angry if we start diagonalizing stuff... so the right way to do is going for a *matrix function times vector* computation.

⚙️ We need to compute  $g(z) = z^{-\alpha}$  for  $\alpha \in (0, 1)$ ,

⚙️ on a matrix  $A_n$  that is either **symmetric and positive definite**, or of a matrix that is *similar* to an SPD matrix,

⚙️  $A_n$  has also a condition number that grows (at least quadratically) with its size, i.e., is **ill-conditioned**.

❓ What method do we select?

🔧  $A_n$  is sparse and, if we deal with a regular uniform grid maybe also Toeplitz, a Lanczos **polynomial Krylov** with fast convergence would be perfect if it reaches convergence with a number of iteration independent of the size  $n$ .

❓ Is this the case?

# The Polynomial Krylov Method

If we use a polynomial Krylov subspace


$$\mathcal{K}_\ell(A_n, \mathbf{v}) = \text{Span}\{\mathbf{v}, A_n \mathbf{v}, \dots, A_n^{\ell-1} \mathbf{v}\}$$

to solve the problem, then the behavior is controlled by the approximation property

$$\|\mathbf{x} - \mathbf{x}_\ell\| \leq C \cdot \min_{p(z) \in \mathbb{P}_{\ell-1}} \max_{z \in \Lambda(A_n)} |p(z) - z^{-\alpha}|$$

for  $\mathbb{P}_{\ell-1}$  the set of polynomial of degree  $\leq \ell$ , and  $C$  a constant *independent* of  $A$  and  $\ell$ .

```
ytrue = mpower(full(An), -alpha)*b;  
[Q,H] = arnoldi(An,b,l);  
for j=1:l  
    y = Q(:,1:j)*(mpower(H(1:j,1:j),  
        ↪ -alpha)*(Q(:,1:j)'*b));  
    err(j) = norm(y-ytrue)./norm(ytrue);  
end
```



# Rational Krylov Method

---

We need **better functions** for our approximation problem, i.e., *rational functions*!

# Rational Krylov Method

We need **better functions** for our approximation problem, i.e., *rational functions*!

## A general framework

Given a set of scalars  $\{\sigma_1, \dots, \sigma_{k-1}\} \subset \overline{\mathbb{C}}$  (the extended complex plane), that are not eigenvalues of  $A$ , let

$$q_{k-1}(z) = \prod_{j=1}^{k-1} (\sigma_j - z).$$

The **rational Krylov** subspace of order  $k$  associated with  $A$ ,  $\mathbf{v}$  and  $q_{k-1}$  is defined by

$$\mathcal{Q}_k(A, \mathbf{v}) = [q_{k-1}(A)]^{-1} \mathcal{K}_k(A, \mathbf{v}), \quad \mathcal{K}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, A\mathbf{v}, \dots, A^{k-1}\mathbf{v}\}.$$

# Rational Krylov Method

We need **better functions** for our approximation problem, i.e., *rational functions*!

## A general framework

Given a set of scalars  $\{\sigma_1, \dots, \sigma_{k-1}\} \subset \overline{\mathbb{C}}$  (the extended complex plane), that are not eigenvalues of  $A$ , let

$$q_{k-1}(z) = \prod_{j=1}^{k-1} (\sigma_j - z).$$

The **rational Krylov** subspace of order  $k$  associated with  $A$ ,  $\mathbf{v}$  and  $q_{k-1}$  is defined by

$$\mathcal{Q}_k(A, \mathbf{v}) = [q_{k-1}(A)]^{-1} \mathcal{K}_k(A, \mathbf{v}), \quad \mathcal{K}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, A\mathbf{v}, \dots, A^{k-1}\mathbf{v}\}.$$

## A matrix expression

Given  $\{\mu_1, \dots, \mu_{k-1}\} \subset \overline{\mathbb{C}}$  such that  $\sigma_j \neq \mu_j^{-2}$ , we define the matrices

$$C_j = (\mu_j \sigma_j A - I) (\sigma_j I - A)^{-1}, \text{ and } \mathcal{Q}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, C_1 \mathbf{v}, \dots, C_{k-1} \cdots C_2 C_1 \mathbf{v}\}.$$

# Rational Krylov Method

## A matrix expression

Given  $\{\mu_1, \dots, \mu_{k-1}\} \subset \overline{\mathbb{C}}$  such that  $\sigma_j \neq \mu_j^{-2}$ , we define the matrices

$$C_j = (\mu_j \sigma_j A - I) (\sigma_j I - A)^{-1}, \text{ and } \mathcal{Q}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, C_1 \mathbf{v}, \dots, C_{k-1} \cdots C_2 C_1 \mathbf{v}\}.$$

**Polynomial Krylov**  $\mathcal{W}_k(A, \mathbf{v}) = \mathcal{K}_k(A, \mathbf{v})$  set  $\mu_j = 1$  and  $\sigma_j = \infty$  for each  $j$ ,

# Rational Krylov Method

## A matrix expression

Given  $\{\mu_1, \dots, \mu_{k-1}\} \subset \overline{\mathbb{C}}$  such that  $\sigma_j \neq \mu_j^{-2}$ , we define the matrices

$$C_j = (\mu_j \sigma_j A - I) (\sigma_j I - A)^{-1}, \text{ and } \mathcal{Q}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, C_1 \mathbf{v}, \dots, C_{k-1} \cdots C_2 C_1 \mathbf{v}\}.$$

**Polynomial Krylov**  $\mathcal{W}_k(A, \mathbf{v}) = \mathcal{K}_k(A, \mathbf{v})$  set  $\mu_j = 1$  and  $\sigma_j = \infty$  for each  $j$ ,

**Extended Krylov**  $\mathcal{W}_{2k-1}(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, A^{-1}\mathbf{v}, A\mathbf{v}, \dots, A^{-(k-1)}\mathbf{v}, A^{k-1}\mathbf{v}\}$ , set

$$(\mu_j, \sigma_j) = \begin{cases} (1, \infty), & \text{for } j \text{ even,} \\ (0, 0), & \text{for } j \text{ odd.} \end{cases}$$



# Rational Krylov Method

## A matrix expression

Given  $\{\mu_1, \dots, \mu_{k-1}\} \subset \overline{\mathbb{C}}$  such that  $\sigma_j \neq \mu_j^{-2}$ , we define the matrices

$$C_j = (\mu_j \sigma_j A - I) (\sigma_j I - A)^{-1}, \text{ and } \mathcal{Q}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, C_1 \mathbf{v}, \dots, C_{k-1} \cdots C_2 C_1 \mathbf{v}\}.$$

**Polynomial Krylov**  $\mathcal{W}_k(A, \mathbf{v}) = \mathcal{K}_k(A, \mathbf{v})$  set  $\mu_j = 1$  and  $\sigma_j = \infty$  for each  $j$ ,

**Extended Krylov**  $\mathcal{W}_{2k-1}(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, A^{-1}\mathbf{v}, A\mathbf{v}, \dots, A^{-(k-1)}\mathbf{v}, A^{k-1}\mathbf{v}\}$ , set

$$(\mu_j, \sigma_j) = \begin{cases} (1, \infty), & \text{for } j \text{ even,} \\ (0, 0), & \text{for } j \text{ odd.} \end{cases}$$

**Shift-And-Invert**  $\mathcal{W}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, (\sigma I - A)^{-1}\mathbf{v}, \dots, (\sigma I - A)^{-(k-1)}\mathbf{v}\}$ , take  $\mu_j = 0$  and  $\sigma_j = \sigma$  for each  $j$ ,

# Rational Krylov Method

## A matrix expression

Given  $\{\mu_1, \dots, \mu_{k-1}\} \subset \overline{\mathbb{C}}$  such that  $\sigma_j \neq \mu_j^{-2}$ , we define the matrices

$$C_j = (\mu_j \sigma_j A - I) (\sigma_j I - A)^{-1}, \text{ and } \mathcal{Q}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, C_1 \mathbf{v}, \dots, C_{k-1} \cdots C_2 C_1 \mathbf{v}\}.$$

**Polynomial Krylov**  $\mathcal{W}_k(A, \mathbf{v}) = \mathcal{K}_k(A, \mathbf{v})$  set  $\mu_j = 1$  and  $\sigma_j = \infty$  for each  $j$ ,

**Extended Krylov**  $\mathcal{W}_{2k-1}(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, A^{-1}\mathbf{v}, A\mathbf{v}, \dots, A^{-(k-1)}\mathbf{v}, A^{k-1}\mathbf{v}\}$ , set


$$(\mu_j, \sigma_j) = \begin{cases} (1, \infty), & \text{for } j \text{ even,} \\ (0, 0), & \text{for } j \text{ odd.} \end{cases}$$

**Shift-And-Invert**  $\mathcal{W}_k(A, \mathbf{v}) = \text{Span}\{\mathbf{v}, (\sigma I - A)^{-1}\mathbf{v}, \dots, (\sigma I - A)^{-(k-1)}\mathbf{v}\}$ , take  $\mu_j = 0$  and  $\sigma_j = \sigma$  for each  $j$ ,

 We are left our usual problem: **how do we select the poles?**


# Pole Selection Strategies

---

 Given a function  $g(z)$  we find an **explicit (minimal) rational approximation**:

$$g(z) = \frac{P_\ell(z)}{Q_q(z)}, \quad P_\ell \in \mathbb{P}_\ell[x], \quad Q_q \in \mathbb{P}_q[x],$$

and use its poles for the RK-Method.

- ✓ Reasonably easy to get worst case scenario bounds;
- ✗ If we want an approximation of the same class with more poles we *usually* need to redo everything from scratch;
-  There exist **brute force** algorithm to get such approximations.

# Pole Selection Strategies

1] Given a function  $g(z)$  we find an **explicit (minimal) rational approximation**:

$$g(z) = \frac{P_\ell(z)}{Q_q(z)}, \quad P_\ell \in \mathbb{P}_\ell[x], \quad Q_q \in \mathbb{P}_q[x],$$


and use its poles for the RK-Method.

- ✓ Reasonably easy to get worst case scenario bounds;
- ✗ If we want an approximation of the same class with more poles we *usually* need to redo everything from scratch;
- 🔧 There exist **brute force** algorithm to get such approximations.

1] Obtain *optimal* rational approximation by solving **best approximation formulations**.

- ✗ It is only possible for certain combinations of complex plane regions and function classes;
- ✓ If one gets an explicit solution of the best approximation, there is usually a way to get nested poles;
- 🔧 In some cases is possible **brute force** our way through it.


# Pole Selection Strategies

 Given a function  $g(z)$  we find an **explicit (minimal) rational approximation**:

$$g(z) = \frac{P_\ell(z)}{Q_q(z)}, \quad P_\ell \in \mathbb{P}_\ell[x], \quad Q_q \in \mathbb{P}_q[x],$$

and use its poles for the RK-Method.

- ✓ Reasonably easy to get worst case scenario bounds;
- ✗ If we want an approximation of the same class with more poles we *usually* need to redo everything from scratch;
- 🔧 There exist **brute force** algorithm to get such approximations.

 Obtain *optimal* rational approximation by solving **best approximation formulations**.

- ✗ It is only possible for certain combinations of complex plane regions and function classes;
- ✓ If one gets an explicit solution of the best approximation, there is usually a way to get nested poles;
- 🔧 In some cases is possible **brute force** our way through it.



## Direct rational approximations

Sometimes it may be worth our while to use directly  $g(A_n)\mathbf{v} = Q_q(A_n)^{-1}P_\ell(A_n)\mathbf{v}$ .

# Best Uniform Rational Approximation (BURA)

---

We try to find the poles by solving the min-max problem

$$\max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = \min_{r_k(t) \in \mathbb{R}_{k,k}} \max_{t \in [0,1]} |t^\alpha - r_k(t)|, \quad \alpha \in (0,1),$$

for  $r_k(t)$  a  $(k, k)$ -rational function.

# Best Uniform Rational Approximation (BURA)

We try to find the poles by solving the min-max problem

$$\max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = \min_{r_k(t) \in \mathbb{R}_{k,k}} \max_{t \in [0,1]} |t^\alpha - r_k(t)|, \quad \alpha \in (0, 1),$$

for  $r_k(t)$  a  $(k, k)$ -rational function.

Theorem (Stahl 2003, Theorem 1)

$$E_{\alpha,k} = \max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = 4^{\alpha+1} |\sin(\alpha\pi)| e^{-2\pi\sqrt{\alpha k}}.$$

# Best Uniform Rational Approximation (BURA)

We try to find the poles by solving the min-max problem

$$\max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = \min_{r_k(t) \in \mathbb{R}_{k,k}} \max_{t \in [0,1]} |t^\alpha - r_k(t)|, \quad \alpha \in (0, 1),$$

for  $r_k(t)$  a  $(k, k)$ -rational function.

Theorem (Stahl 2003, Theorem 1)

$$E_{\alpha,k} = \max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = 4^{\alpha+1} |\sin(\alpha\pi)| e^{-2\pi\sqrt{\alpha k}}.$$



The matrix is approximated as  $A^{-\alpha} \approx \lambda_{1,h}^{-\alpha} r_{\alpha,k}(\lambda_{1,h} A^{-1})$ .



# Best Uniform Rational Approximation (BURA)

We try to find the poles by solving the min-max problem

$$\max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = \min_{r_k(t) \in \mathbb{R}_{k,k}} \max_{t \in [0,1]} |t^\alpha - r_k(t)|, \quad \alpha \in (0, 1),$$

for  $r_k(t)$  a  $(k, k)$ -rational function.

Theorem (Stahl 2003, Theorem 1)

$$E_{\alpha,k} = \max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = 4^{\alpha+1} |\sin(\alpha\pi)| e^{-2\pi\sqrt{\alpha k}}.$$

🔧 The matrix is approximated as  $A^{-\alpha} \approx \lambda_{1,h}^{-\alpha} r_{\alpha,k}(\lambda_{1,h} A^{-1})$ .

❓ But how do we compute  $r_{\alpha,k}(t)$  in practice?

# Best Uniform Rational Approximation (BURA)

We try to find the poles by solving the min-max problem

$$\max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = \min_{r_k(t) \in \mathbb{R}_{k,k}} \max_{t \in [0,1]} |t^\alpha - r_k(t)|, \quad \alpha \in (0, 1),$$

for  $r_k(t)$  a  $(k, k)$ -rational function.

Theorem (Stahl 2003, Theorem 1)

$$E_{\alpha,k} = \max_{t \in [0,1]} |t^\alpha - r_{\alpha,k}(t)| = 4^{\alpha+1} |\sin(\alpha\pi)| e^{-2\pi\sqrt{\alpha k}}.$$

- 🔧 The matrix is approximated as  $A^{-\alpha} \approx \lambda_{1,h}^{-\alpha} r_{\alpha,k}(\lambda_{1,h} A^{-1})$ .
- ❓ But how do we compute  $r_{\alpha,k}(t)$  in practice?
- 🔧 There is no *explicit solution*, thus we need to use a **numerical method**.

# Best Uniform Rational Approximation (BURA)

The *workhorse* for computing BURA is the **Remez algorithm** (Braess 1986, § 6.B)

- 💡 Determine the points at which the error of the BURA equioscillates.
- 🔧 Starting with a *suitable initial guess*, it iteratively determines a rational approximation passing through these points while shifting one or more toward a nearby local maximum.
- 🔧 Implementation is **delicate matter**, observe we want both stability and possibly quadratic convergence.

Chose  $P^{(0)}/Q^{(0)} \in \mathbb{R}_{m,n}$  and  $l$  points  $\{x_i^1\}_{i=1}^l$ ;  
 $k \leftarrow 1$ ;

**while** *not satisfied* **do**

Determine  $P^{(k)}/Q^{(k)} \in \mathbb{R}_{m,n}$  and  $\eta_k \in \mathbb{R}$  such that for  $i = 1, 2, \dots, l$

$$f(x_i^k) - P^{(k)}(x_i^k)/Q^{(k)}(x_i^k) = (-1)^i \eta_k$$

Determine  $x_1^{k+1} < x_2^{k+1} < \dots < x_l^{k+1}$  such that for  $i = 1, 2, \dots, l$


$$s(-1)^i (f - P^{(k)}/Q^{(k)})(x_i^{k+1}) \geq |\eta_k|,$$

and that for one  $i \in \{1, 2, \dots, l\}$  the left-hand side equals  $\|f - P^{(k)}/Q^{(k)}\|$ ,  $s = \pm 1$ ;

$k \leftarrow k + 1$ ;

**end**

# Best Uniform Rational Approximation (BURA)


A **recent** and **available** implementation is given in the  Python `baryrat` package, see (Hofreither 2021).

```
import numpy as np
import baryrat

alpha = 0.5
def f(x): return x**alpha
r = baryrat.brasil(f, [0,1], 5)
```


That gives us the `r.poles()`:

$$\sigma = \{-3.21294874e + 00, -1.62633499e - 01, \\ -1.27958136e - 02, -6.62129541e - 04, \\ -1.22326563e - 05\}.$$



$$k = 5, \alpha = 1/2$$

# Best Uniform Rational Approximation (BURA)


A **recent** and **available** implementation is given in the  Python `baryrat` package, see (Hofreither 2021).

```
import numpy as np
import baryrat

alpha = 0.5
def f(x): return x**alpha
r = baryrat.brasil(f, [0,1], 5)
```


That gives us the `r.poles()`:

$$\sigma = \{-3.21294874e + 00, -1.62633499e - 01, \\ -1.27958136e - 02, -6.62129541e - 04, \\ -1.22326563e - 05\}.$$



$$k = 10, \alpha = 1/2$$

# Best Uniform Rational Approximation (BURA)


A **recent** and **available** implementation is given in the  Python `baryrat` package, see (Hofreither 2021).

```
import numpy as np
import baryrat

alpha = 0.5
def f(x): return x**alpha
r = baryrat.brasil(f, [0,1], 5)
```


That gives us the `r.poles(0)`:

$$\sigma = \{-3.21294874e + 00, -1.62633499e - 01, \\ -1.27958136e - 02, -6.62129541e - 04, \\ -1.22326563e - 05\}.$$



$$k = 20, \alpha = 1/2$$

# Best Uniform Rational Approximation (BURA)


A **recent** and **available** implementation is given in the  Python `baryrat` package, see (Hofreither 2021).

```
import numpy as np
import baryrat

alpha = 0.5
def f(x): return x**alpha
r = baryrat.brazil(f, [0,1], 5)
```

That gives us the `r.poles(0)`:

$$\sigma = \{-3.21294874e + 00, -1.62633499e - 01, \\ -1.27958136e - 02, -6.62129541e - 04, \\ -1.22326563e - 05\}.$$



$$k = 40, \alpha = 1/2$$

# Best Uniform Rational Approximation (BURA)

---

One can couple the error analysis with the one coming from the discretization of the Laplacian to get overall results (Harizanov et al. 2020).



# Best Uniform Rational Approximation (BURA)

One can couple the error analysis with the one coming from the discretization of the Laplacian to get overall results (Harizanov et al. 2020).

Theorem (Harizanov et al. 2020, Theorem 4.2).

Let  $\Omega \subset \mathbb{R}^2$  and suppose that the solution is in  $\mathbb{H}^2(\Omega) \cup \mathbb{H}_0^1(\Omega)$  and satisfies  $\|(-\Delta)^{-\alpha} f\|_{\mathbb{H}^2(\Omega)} \leq c\|f\|$ . Then for  $f \in \mathbb{H}^{1+\gamma}(\Omega)$ ,  $\gamma > 0$ , the solution  $\mathbf{u}_h$  given by

$$\mathbf{u}_h = \lambda_{1,h}^{-\alpha} (\lambda_{1,h} A^{-1})^{\alpha} I_h f, \quad A = M_n^{-1} A_n, \quad I_h \text{ Interpolation,}$$

satisfies

$$\|(-\Delta)^{-\alpha} f - \mathbf{u}_h\| \leq C(h^{2\alpha} + h^{1+\gamma})\|f\|_{\mathbb{H}^{1+\gamma}(\Omega)}.$$

# Best Uniform Rational Approximation (BURA)

One can couple the error analysis with the one coming from the discretization of the Laplacian to get overall results (Harizanov et al. 2020).

Theorem (Harizanov et al. 2020, Theorem 4.2).

Let  $\Omega \subset \mathbb{R}^2$  and suppose that the solution is in  $\mathbb{H}^2(\Omega) \cup \mathbb{H}_0^1(\Omega)$  and satisfies  $\|(-\Delta)^{-\alpha} f\|_{\mathbb{H}^2(\Omega)} \leq c\|f\|$ . Then for  $f \in \mathbb{H}^{1+\gamma}(\Omega)$ ,  $\gamma > 0$ , the solution  $\mathbf{u}_h$  given by

$$\mathbf{u}_h = \lambda_{1,h}^{-\alpha} (\lambda_{1,h} A^{-1})^{\alpha} I_h f, \quad A = M_n^{-1} A_n, \quad I_h \text{ Interpolation,}$$

satisfies

$$\|(-\Delta)^{-\alpha} f - \mathbf{u}_h\| \leq C(h^{2\alpha} + h^{1+\gamma})\|f\|_{\mathbb{H}^{1+\gamma}(\Omega)}.$$

➤ Using **lumped FEM**, it is possible to have the error of the **fully discrete scheme** (Harizanov et al. 2020, Corollary 4.3), and then balance the discretization and the BURA error.

# Best Uniform Rational Approximation (BURA)

One can couple the error analysis with the one coming from the discretization of the Laplacian to get overall results (Harizanov et al. 2020).

Theorem (Harizanov et al. 2020, Theorem 4.2).

Let  $\Omega \subset \mathbb{R}^2$  and suppose that the solution is in  $\mathbb{H}^2(\Omega) \cup \mathbb{H}_0^1(\Omega)$  and satisfies  $\|(-\Delta)^{-\alpha} f\|_{\mathbb{H}^2(\Omega)} \leq c\|f\|$ . Then for  $f \in \mathbb{H}^{1+\gamma}(\Omega)$ ,  $\gamma > 0$ , the solution  $\mathbf{u}_h$  given by

$$\mathbf{u}_h = \lambda_{1,h}^{-\alpha} (\lambda_{1,h} A^{-1})^\alpha I_h f, \quad A = M_n^{-1} A_n, \quad I_h \text{ Interpolation,}$$

satisfies

$$\|(-\Delta)^{-\alpha} f - \mathbf{u}_h\| \leq C(h^{2\alpha} + h^{1+\gamma})\|f\|_{\mathbb{H}^{1+\gamma}(\Omega)}.$$

🔧 Using **lumped FEM**, it is possible to have the error of the **fully discrete scheme** (Harizanov et al. 2020, Corollary 4.3), and then balance the discretization and the BURA error.

🔧 The intend usage of these scheme is *outside* of a Krylov method.

# Quadrature-based approaches

Another viable approach is to use a *rational approximation* based on a **quadrature formula**.

- ☰ There is more than a *connection* between **quadrature formulas** and **rational approximations**.
- 📄 Padé approximants can be viewed as formal Gaussian quadrature methods (Brezinski 1980, Page 34).
- 🕒 This connection was already known to Gauß  
*C. F. Gauss, Methodus nova integralium valores per approximationem inveniendi, Comment. Soc. Reg. Scient. Gotting. Recent., 1814*



C.F. Gauß  
(1777-1855)

# Quadrature-based approaches

Another viable approach is to use a *rational approximation* based on a **quadrature formula**.

☰ There is more than a *connection* between **quadrature formulas** and **rational approximations**.

📄 Padé approximants can be viewed as formal Gaussian quadrature methods (Brezinski 1980, Page 34).

🕒 This connection was already known to Gauß  
*C. F. Gauss, Methodus nova integralium valores per approximationem inveniendi, Comment. Soc. Reg. Scient. Gotting. Recent., 1814*



C.F. Gauß  
(1777-1855)

💡 The idea is always the same **1.** Find an integral representation of the function of interest. **2.** Find a change of variables that makes a Gauss-type weight appear. **3.** Rational approximation is obtained by the Gauss quadrature formula. **4.** The error analysis relies on the analysis for the formula.

# The Gauss-Jacobi approach

---

This is an idea from (Aceto, Bertaccini, et al. 2019; Aceto and Novati 2018).

# The Gauss-Jacobi approach

---

This is an idea from (Aceto, Bertaccini, et al. 2019; Aceto and Novati 2018). We do **step 1** by looking through a book:

Proposition (Bhatia 1997, example V.1.10, 21, section 5.5.5)

Let  $A \in \mathbb{R}^{n \times n}$  be such that  $\Lambda(A) \subset \mathbb{C} \setminus (-\infty, 0]$ . For  $\alpha \in (0, 1)$  the following representation holds

$$A^\alpha = \frac{\sin(\alpha\pi)}{\alpha\pi} A \int_0^\infty \left( \rho^{1/\alpha} I + A \right)^{-1} d\rho.$$

# The Gauss-Jacobi approach

This is an idea from (Aceto, Bertaccini, et al. 2019; Aceto and Novati 2018). We do **step 1** by looking through a book:

Proposition (Bhatia 1997, example V.1.10, 21, section 5.5.5)

Let  $A \in \mathbb{R}^{n \times n}$  be such that  $\Lambda(A) \subset \mathbb{C} \setminus (-\infty, 0]$ . For  $\alpha \in (0, 1)$  the following representation holds

$$A^\alpha = \frac{\sin(\alpha\pi)}{\alpha\pi} A \int_0^\infty \left( \rho^{1/\alpha} I + A \right)^{-1} d\rho.$$

Now do **step 2**, i.e., a *change of variables*:

$$\rho^{1/\alpha} = \tau \frac{1-t}{1+t}, \quad \tau > 0.$$



# The Gauss-Jacobi approach

---

By plugging the change of variables in the integral, we find

$$A^\alpha = \frac{2 \sin(\alpha\pi) \tau^\alpha}{\pi} A \int_{-1}^1 (1-t)^{\alpha-1} (1+t)^{-\alpha} (\tau(1-t)I + (1+t)A)^{-1} dt.$$

# The Gauss-Jacobi approach

---

By plugging the change of variables in the integral, we find

$$A^\alpha = \frac{2 \sin(\alpha\pi) \tau^\alpha}{\pi} A \int_{-1}^1 (1-t)^{\alpha-1} (1+t)^{-\alpha} (\tau(1-t)I + (1+t)A)^{-1} dt.$$

We made the **weights of the Gauss-Jacobi quadrature** appear, thus

$$\left(\frac{1}{\tau}A\right)^{\aleph} \approx \frac{1}{\tau}A \sum_{j=1}^k \frac{2 \sin(\alpha\pi)}{\pi} \frac{\omega_j}{1+\theta_j} \left(\frac{1-\theta_j}{1+\theta_j} + \frac{1}{\tau}A\right)^{-1},$$

- ⚙  $\omega_j$  and  $\theta_j$  are, respectively, the weights and nodes of the Gauss-Jacobi quadrature formula with weight function  $(1-t)^{\alpha-1}(1+t)^{-\alpha}$ ,
- 🔧 we should use *error analysis* to fix the  $\tau$  parameter.
- 📄 From (Frommer, Güttel, and Schweitzer 2014, Lemma 4.4) we know that the  $k$ -point Gauss-Jacobi quadrature corresponds to the  $(k-1, k)$ -Padé approximant of  $(z/\tau)^{\alpha-1}$  centered at 1.

# The Gauss-Jacobi approach

As we have seen from the BURA example, we may be interested in  $g(z) = z^{-\alpha}$ ,  $\alpha \in (0, 1)$ , but it is easy to rewrite the approximation as

$$z^{-\alpha/2} \approx \sum_{j=1}^k \frac{2 \sin(\alpha\pi) \tau^{1-\alpha/2}}{\pi} \frac{\omega_j}{1 + \theta_j} \left( \frac{\tau(1 - \theta_j)}{1 + \theta_j} + z \right)^{-1} \triangleq R_{k-1,k}(z), \quad \tau > 0$$

⚙  $\omega_j$  and  $\theta_j$  are the weights and nodes of the Gauss-Jacobi quadrature formula with weight  $(1-x)^{-\alpha}(1+x)^{\alpha-1}$ .

🔧 If we rearrange the expression we then find

$$R_{k-1,k}(z) = \frac{p_{k-1}(z)}{q_k(z)} = \frac{\chi \prod_{r=1}^{k-1} (z + \epsilon_r)}{\prod_{j=1}^k (z + \eta_j)}, \quad \chi = \frac{\eta_k}{\tau^\alpha} \frac{\binom{k+\alpha/2-1}{k-1}}{\binom{k-\alpha}{k}} \prod_{j=1}^{k-1} \frac{\eta_j}{\epsilon_j}.$$

for

$$\epsilon_r = \tau \frac{1 - \zeta_r}{1 + \zeta_r}, \quad r = 1, 2, \dots, k-1, \quad \eta_j = \frac{\tau(1 - \theta_j)}{1 + \theta_j}, \quad j = 1, 2, \dots, k.$$

# The Gauss-Jacobi approach

---

To fix the  $\tau > 0$  parameter we need the error analysis from (Aceto and Novati 2019) to bound the *truncation error*:

$$E_{k-1,k}(\lambda/\tau) \triangleq (\lambda/\tau)^{-\alpha} - R_{k-1,k}(\lambda/\tau).$$

- When working with these expression, usually one can manipulate and express them in terms of *Gauss-Hypergeometric functions*, then use their asymptotic to produce the bound, e.g., in this case

$$z = 1 - \frac{\lambda}{t}, \quad (1 - z)^{-\alpha} = {}_2F_1 \left( \begin{matrix} 1, \alpha \\ 1 \end{matrix}; z \right), \quad |\arg(1 - z)| < \pi.$$

# The Gauss-Jacobi approach

To fix the  $\tau > 0$  parameter we need the error analysis from (Aceto and Novati 2019) to bound the *truncation error*:

$$E_{k-1,k}(\lambda/\tau) \triangleq (\lambda/\tau)^{-\alpha} - R_{k-1,k}(\lambda/\tau).$$

- When working with these expression, usually one can manipulate and express them in terms of *Gauss-Hypergeometric functions*, then use their asymptotic to produce the bound, e.g., in this case

$$z = 1 - \frac{\lambda}{t}, \quad (1 - z)^{-\alpha} = {}_2F_1 \left( \begin{matrix} 1, \alpha \\ 1 \end{matrix}; z \right), \quad |\arg(1 - z)| < \pi.$$

Proposition (Aceto and Novati 2019, Proposition 2)

For large values of  $k$ , the following representation for the truncation error holds

$$E_{k-1,k}(\lambda/\tau) = 2 \sin(\alpha\pi) (\lambda/\tau)^{-\alpha} \left[ \frac{\sqrt{\lambda} - \sqrt{\tau}}{\sqrt{\lambda} + \sqrt{\tau}} \right]^{2k} (1 + O(1/k)).$$

# The Gauss-Jacobi approach

Theorem (Aceto and Novati 2019, Theorem 2)

If  $\mathcal{L}$  is a self-adjoint positive operator on a separable Hilbert space  $\mathbb{H}$  with spectrum  $\Lambda(\mathcal{L}) \subset [c, +\infty)$ ,  $c > 0$  having a compact inverse, then

$$\left\| \mathcal{L}^{-\alpha} - \tau_k^{-\alpha} R_{k-1,k} \left( \frac{1}{\tau_k} \mathcal{L} \right) \right\|_{\mathbb{H} \rightarrow \mathbb{H}} \leq 2 \sin(\alpha\pi) c^{-\alpha} \left( \frac{2k\sqrt{e}}{\alpha} \right)^{-4\alpha} \left[ 2 \ln \left( \frac{2k}{\alpha} \right) + 1 \right]^{2\alpha} (1 + O(k^{-2})),$$

for

$$\tau_k = c \left( \frac{\alpha}{2ke} \right)^2 \exp \left( 2W \left( \frac{4k^2 e}{\alpha^2} \right) \right),$$

where  $W$  denotes the Lambert  $W$ -function.

👁 It becomes **increasingly difficult** if the spectrum is close to the branch point of  $z^{-\alpha}$ .

# The Gauss-Jacobi approach (bounded operators)

---

If  $\mathcal{L}_N$  is a **bounded operator**, i.e.,  $\Lambda(\mathcal{L}_N) \in [c, \lambda_N]$  then the min-max problem for  $|E_{k-1,k}(\lambda/\tau)|$  have two different solutions for *small* and *large* values of  $k$ .

We call  $\bar{\lambda} = \frac{\tau}{\alpha^2}(k + \sqrt{k^2 + 1})^2$

$\bar{\lambda} < \lambda_N$  ( $k$  small) The previous estimate is still good, i.e.,

$$\tau_k = c \left( \frac{\alpha}{2ke} \right)^2 \exp \left( 2W \left( \frac{4k^2 e}{\alpha^2} \right) \right),$$

$\bar{\lambda} > \lambda_N$  ( $k$  large) then

$$\hat{\tau}_k = \left( -\frac{\alpha\sqrt{(\lambda_N)}}{8k} \ln \left( \frac{\lambda_N}{c} \right) + \sqrt{\left( \frac{\alpha\sqrt{\lambda_N}}{8k} \ln \left( \frac{\lambda_N}{c} \right) \right)^2 + \sqrt{c\lambda_N}} \right)^2.$$

# The Gauss-Jacobi approach (bounded operators)

Theorem (Aceto and Novati 2019, Theorem 3)

Let  $\bar{k}$  be such that for each  $k \geq \bar{k}$  we have  $\bar{\lambda} = \bar{\lambda}(k) > \lambda_N$ . Then for each  $k \geq \bar{k}$ , taking  $\tau = \hat{\tau}_k$ , the following bound holds

$$\left\| \mathcal{L}_N^{-\alpha} - \hat{\tau}_k^{-\alpha} R_{k-1,k} \left( \frac{1}{\hat{\tau}_k} \mathcal{L}_N \right) \right\|_2 \leq 2 \sin(\alpha\pi) (c\lambda_N)^{-\alpha/2} \exp \left( -4k \left( \frac{c}{\lambda_N} \right)^{1/4} \right) (1 + O(k^{-1})).$$

- 👁 The bound gets worse when we refine the discretization of the differential operator!
- 🔧 The choice of  $\tau$  is better than the asymptotically selected value  $\tau_\infty = \sqrt{c\Lambda_N}$ .



# The Gauss-Jacobi approach (bounded operators)

Theorem (Aceto and Novati 2019, Theorem 3)

Let  $\bar{k}$  be such that for each  $k \geq \bar{k}$  we have  $\bar{\lambda} = \bar{\lambda}(k) > \lambda_N$ . Then for each  $k \geq \bar{k}$ , taking  $\tau = \hat{\tau}_k$ , the following bound holds

$$\left\| \mathcal{L}_N^{-\alpha} - \hat{\tau}_k^{-\alpha} R_{k-1,k} \left( \frac{1}{\hat{\tau}_k} \mathcal{L}_N \right) \right\|_2 \leq 2 \sin(\alpha\pi) (c\lambda_N)^{-\alpha/2} \exp \left( -4k \left( \frac{c}{\lambda_N} \right)^{1/4} \right) (1 + O(k^{-1})).$$

👁 The bound gets worse when we refine the discretization of the differential operator!

🔧 The choice of  $\tau$  is better than the asymptotically selected value  $\tau_\infty = \sqrt{c\lambda_N}$ .

The choice is made as

$$\tau_{k,N} = \begin{cases} \tau_k, & k < \bar{k}, \\ \hat{\tau}_k, & k \geq \bar{k}, \end{cases} \quad \text{for } \bar{k} = \left\lceil \frac{\alpha}{2\sqrt{2}} \sqrt{\ln \left( \frac{\lambda_N}{c} e^2 \right)} \left( \frac{\lambda_N}{c} \right)^{\frac{1}{4}} \right\rceil.$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

---

We start again from an **integral representation** (Bonito and Pasciak 2015)

$$\mathcal{L}^{-\alpha} = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^{+\infty} t^{2\alpha-1} (\mathcal{I} + t^2 \mathcal{L})^{-1} dt, \quad \alpha \in (0, 1).$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

We start again from an **integral representation** (Bonito and Pasciak 2015)

$$\mathcal{L}^{-\alpha} = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^{+\infty} t^{2\alpha-1} (\mathcal{I} + t^2 \mathcal{L})^{-1} dt, \quad \alpha \in (0, 1).$$

Then, we go for the **change of variables**  $y = \ln t$  we obtain

$$\mathcal{L}^{-\alpha} = \frac{2 \sin(\alpha\pi)}{\pi} \int_{-\infty}^{+\infty} e^{2\alpha y} (\mathcal{I} + e^{2y} \mathcal{L})^{-1} dy, \quad \alpha \in (0, 1).$$

$$= \int_{-\infty}^0 e^{2\alpha y} (\mathcal{I} + e^{2y} \mathcal{L})^{-1} dy + \int_0^{+\infty} e^{2\alpha y} (\mathcal{I} + e^{2y} \mathcal{L})^{-1} dy$$

$$\begin{array}{l} 2\alpha y = -x \\ 2(1-\alpha)y = x \end{array} \rightarrow = \frac{1}{2\alpha} \int_0^{+\infty} e^{-x} (\mathcal{I} + e^{-x/\alpha} \mathcal{L})^{-1} dx + \frac{1}{2(1-\alpha)} \int_0^{+\infty} e^{-x} (e^{-x/(1-\alpha)} \mathcal{I} + \mathcal{L})^{-1} dx.$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

We start again from an **integral representation** (Bonito and Pasciak 2015)

$$\mathcal{L}^{-\alpha} = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^{+\infty} t^{2\alpha-1} (\mathcal{I} + t^2 \mathcal{L})^{-1} dt, \quad \alpha \in (0, 1).$$

Then, we go for the **change of variables**  $y = \ln t$  we obtain

$$\mathcal{L}^{-\alpha} = \frac{\sin(\alpha\pi)}{\alpha\pi} I^{(1)}(\mathcal{L}) + \frac{\sin(\alpha\pi)}{(1-\alpha)\pi} I^{(2)}(\mathcal{L}),$$

for

$$I^{(1)}(\lambda) = \int_0^{+\infty} e^{-x} (1 + e^{-x/\alpha} \lambda)^{-1} dx, \quad I^{(2)}(\lambda) = \int_0^{+\infty} e^{-x} (e^{-x/(1-\alpha)} + \lambda)^{-1} dx.$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

We start again from an **integral representation** (Bonito and Pasciak 2015)

$$\mathcal{L}^{-\alpha} = \frac{2 \sin(\alpha\pi)}{\pi} \int_0^{+\infty} t^{2\alpha-1} (\mathcal{I} + t^2 \mathcal{L})^{-1} dt, \quad \alpha \in (0, 1).$$

Then, we go for the **change of variables**  $y = \ln t$  we obtain

$$\mathcal{L}^{-\alpha} =$$

for

$$I^{(1)}(\lambda) = \int_0^{+\infty} e^{-x} (1 + e^{-x/\alpha} \lambda)^{-1} dx, \quad I^{(2)}(\lambda) = \int_0^{+\infty} e^{-x} (e^{-x/(1-\alpha)} + \lambda)^{-1} dx.$$

The weight  $\omega(x) = e^{-x}$ , is the weight of **Gauss-Laguerre** formulas.

# A Gauss-Laguerre approach (Aceto and Novati 2022)

If we call the weights  $w_j^{(n)}$  and nodes  $\vartheta_j^{(n)}$  (in ascending order) of the Gauss-Laguerre formula, then we obtain the following  $(2n-1, 2n)$  rational approximation:

$$\mathcal{L}^{-\alpha} \approx \frac{\sin(\alpha\pi)}{\alpha\pi} R_{n-1,n}^{(1)}(\mathcal{L}) + \frac{\sin(\alpha\pi)}{(1-\alpha)\pi} R_{n-1,n}^{(2)}(\mathcal{L}) \triangleq R_{2n-1,2n}(\mathcal{L}),$$

where

$$\begin{aligned} R_{n-1,n}^{(1)}(\lambda) &= \sum_{j=1}^n w_j^{(n)} \left( 1 + e^{-\vartheta_j^{(n)}/\alpha\lambda} \right)^{-1}, \\ R_{n-1,n}^{(2)}(\lambda) &= \sum_{j=1}^n w_j^{(n)} \left( e^{-\vartheta_j^{(n)/(1-\alpha)} + \lambda} \right)^{-1}. \end{aligned}$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

If we call the weights  $w_j^{(n)}$  and nodes  $\vartheta_j^{(n)}$  (in ascending order) of the Gauss-Laguerre formula, then we obtain the following  $(2n-1, 2n)$  rational approximation:

$$\mathcal{L}^{-\alpha} \approx \frac{\sin(\alpha\pi)}{\alpha\pi} R_{n-1,n}^{(1)}(\mathcal{L}) + \frac{\sin(\alpha\pi)}{(1-\alpha)\pi} R_{n-1,n}^{(2)}(\mathcal{L}) \triangleq R_{2n-1,2n}(\mathcal{L}),$$

where

$$\begin{aligned} R_{n-1,n}^{(1)}(\lambda) &= \sum_{j=1}^n w_j^{(n)} \left( 1 + e^{-\vartheta_j^{(n)}/\alpha\lambda} \right)^{-1}, \\ R_{n-1,n}^{(2)}(\lambda) &= \sum_{j=1}^n w_j^{(n)} \left( e^{-\vartheta_j^{(n)/(1-\alpha)}} + \lambda \right)^{-1}. \end{aligned}$$

🔧 Third step is using **error estimate for Gauss-Laguerre formulas** to get the bound.

# A Gauss-Laguerre approach (Aceto and Novati 2022)

---

The analysis treats separately the two integrals and requires expressing the error as a *contour integral*:

$$E_n(f) = \frac{1}{2\pi i} \int_{\Gamma} \frac{q_n(z)}{L_n(z)} f(z) dz,$$


here  $L_n(z)$  is the Laguerre polynomial,  $q_n(z)$  is the so-called associated function defined by

$$q_n(z) = \int_0^{+\infty} \frac{e^{-x} L_n(x)}{z - x} dx, \quad z \notin [0, +\infty),$$

and  $\Gamma$  is a contour containing  $[0, +\infty)$  with the additional property that **no singularity** of  $f(z)$  **lies on or within this contour**; see (Davis and Rabinowitz 1984, §4.6).



# A Gauss-Laguerre approach (Aceto and Novati 2022)




The error can be written as

$$E_n(f) = \frac{1}{2\pi i} \left\{ \int_{\Gamma_R} - \int_{C_1} - \int_{C_2} \right\} \frac{q_n(z)}{L_n(z)} f(z) dz.$$

Denote with  $C_1$  and  $C_2$  two arbitrary small circles surrounding the two poles and define

$$\Gamma = \Gamma_R \cup C_1 \cup C_2.$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)



Denote with  $C_1$  and  $C_2$  two arbitrary small circles surrounding the two poles and define

$$\Gamma = \Gamma_R \cup C_1 \cup C_2.$$


The error can be written as

$$E_n(f) = \frac{1}{2\pi i} \left\{ \int_{\Gamma_R} - \int_{C_1} - \int_{C_2} \right\} \frac{q_n(z)}{L_n(z)} f(z) dz.$$

Then using:

$$\begin{aligned} \frac{q_n(z)}{L_n(z)} &= 2\pi e^{-z} [\exp(\sqrt{-z})]^{-2\sqrt{n}} \times \\ &\times \left( 1 + O\left(\frac{1}{n}\right) \right), \quad z \notin [0, +\infty), \end{aligned}$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)



Denote with  $C_1$  and  $C_2$  two arbitrary small circles surrounding the two poles and define

$$\Gamma = \Gamma_R \cup C_1 \cup C_2.$$


The error can be written as

$$E_n(f) = \frac{1}{2\pi i} \left\{ \int_{\Gamma_R} - \int_{C_1} - \int_{C_2} \right\} \frac{q_n(z)}{L_n(z)} f(z) dz.$$

One arrives at

$$\begin{aligned} |E_n(f)| &\leq 4\pi \left| \text{Res}(f(z), z_0) e^{-z_0} \right| \times \\ &\quad \times \left[ \exp(\text{Re}(\sqrt{-z_0})) \right]^{-2\sqrt{n}} \times \\ &\quad \times \left( 1 + O\left(\frac{1}{n}\right) \right). \end{aligned}$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)



Denote with  $C_1$  and  $C_2$  two arbitrary small circles surrounding the two poles and define  $\Gamma = \Gamma_R \cup C_1 \cup C_2$ .

The error can be written as

$$E_n(f) = \frac{1}{2\pi i} \left\{ \int_{\Gamma_R} - \int_{C_1} - \int_{C_2} \right\} \frac{q_n(z)}{L_n(z)} f(z) dz.$$

One arrives at

$$\begin{aligned} |E_n(f)| &\leq 4\pi \left| \text{Res}(f(z), z_0) e^{-z_0} \right| \times \\ &\quad \times \left[ \exp(\text{Re}(\sqrt{-z_0})) \right]^{-2\sqrt{n}} \times \\ &\quad \times \left( 1 + O\left(\frac{1}{n}\right) \right). \end{aligned}$$

## ⚙️ Procedure

Apply the idea at  $f(z) = (1 + e^{-z/\alpha}\lambda)^{-1}$ , and  $f(z) = (e^{-z/(1-\alpha)} + \lambda)^{-1}$ . For the two integrals.

# A Gauss-Laguerre approach (Aceto and Novati 2022)

Theorem (Aceto and Novati 2022, Proposition 5.3)

Let  $R_{2n-1,2n}(\mathcal{L})$  be the Gauss-Laguerre rational approximation. Then, with respect to the operator norm in  $\mathbb{H}$  we have for  $n$  large enough

$$\|\mathcal{L}^{-\alpha} - R_{2n-1,2n}(\mathcal{L})\| \leq 4 \sin(\alpha\pi) \exp\left(-3(n\alpha^2\pi^2)^{1/3}\right) \left(1 + O\left(n^{-1/3}\right)\right).$$

# A Gauss-Laguerre approach (Aceto and Novati 2022)

Theorem (Aceto and Novati 2022, Proposition 5.3)

Let  $R_{2n-1,2n}(\mathcal{L})$  be the Gauss-Laguerre rational approximation. Then, with respect to the operator norm in  $\mathbb{H}$  we have for  $n$  large enough

$$\|\mathcal{L}^{-\alpha} - R_{2n-1,2n}(\mathcal{L})\| \leq 4 \sin(\alpha\pi) \exp\left(-3(n\alpha^2\pi^2)^{1/3}\right) \left(1 + O\left(n^{-1/3}\right)\right).$$

- 😊 The **convergence** is now **independent of the spectral information** of the matrix, we just need to scale  $A$  to have spectrum in  $[1, +\infty)$ .

# A Gauss-Laguerre approach (Aceto and Novati 2022)

## Theorem (Aceto and Novati 2022, Proposition 5.3)

Let  $R_{2n-1,2n}(\mathcal{L})$  be the Gauss-Laguerre rational approximation. Then, with respect to the operator norm in  $\mathbb{H}$  we have for  $n$  large enough

$$\|\mathcal{L}^{-\alpha} - R_{2n-1,2n}(\mathcal{L})\| \leq 4 \sin(\alpha\pi) \exp\left(-3(n\alpha^2\pi^2)^{1/3}\right) \left(1 + O\left(n^{-1/3}\right)\right).$$

- 😊 The **convergence** is now **independent of the spectral information** of the matrix, we just need to scale  $A$  to have spectrum in  $[1, +\infty)$ .
- 🔧 Truncation and balancing strategies can be applied to the quadratures observing that nodes and weights decay exponentially, i.e., apply

$$\mathcal{L}^{-\alpha} \approx \frac{\sin(\alpha\pi)}{\alpha\pi} R_{k_{n_1}-1, k_{n_1}}^{(1)}(\mathcal{L}) + \frac{\sin(\alpha\pi)}{(1-\alpha)\pi} R_{k_{n_2}-1, k_{n_2}}^{(2)}(\mathcal{L}).$$

# Laplace-Stieltjes and Cauchy-Stieltjes functions

---

Functions expressed as Stieltjes integrals admit a representation of the form:

$$f(z) = \int_0^{\infty} g(t, z) \mu(t) \, dt,$$

where

- $\mu(t)dt$  is a (non-negative) on  $[0, \infty]$ , measure,
- $g(t, z)$  is integrable with respect to that measure.



# Laplace-Stieltjes and Cauchy-Stieltjes functions

Functions expressed as Stieltjes integrals admit a representation of the form:

$$f(z) = \int_0^{\infty} g(t, z) \mu(t) \, dt,$$

where

- $\mu(t)dt$  is a (non-negative) on  $[0, \infty]$ , measure,
- $g(t, z)$  is integrable with respect to that measure.

## Cauchy-Stieltjes

Let  $f(z)$  be a function defined on  $\mathbb{C} \setminus \mathbb{R}_-$ . Then,  $f(z)$  is a *Cauchy-Stieltjes* function if there is a positive measure  $\mu(t)dt$  on  $\mathbb{R}_+$  such that

$$f(z) = \int_0^{\infty} \frac{\mu(t)}{t+z} \, dt.$$

# Laplace-Stieltjes and Cauchy-Stieltjes functions

Functions expressed as Stieltjes integrals admit a representation of the form:

$$f(z) = \int_0^{\infty} g(t, z) \mu(t) dt,$$

where

- $\mu(t)dt$  is a (non-negative) on  $[0, \infty]$ , measure,
- $g(t, z)$  is integrable with respect to that measure.

## Cauchy-Stieltjes

Let  $f(z)$  be a function defined on  $\mathbb{C} \setminus \mathbb{R}_-$ . Then,  $f(z)$  is a *Cauchy-Stieltjes* function if there is a positive measure  $\mu(t)dt$  on  $\mathbb{R}_+$  such that

$$f(z) = \int_0^{\infty} \frac{\mu(t)}{t + z} dt.$$

The function we are interested in is of this class for  $\alpha \in (0, 1)$ :

$$f(z) = z^{-\alpha} = \frac{\sin(\alpha\pi)}{\pi} \int_0^{\infty} \frac{t^{-\alpha}}{t + z} dt.$$

In (Massei and Robol [2021](#)) is given a general bound for the whole class of functions.

## ◀ Back to Zolotarev

---

To **obtain the poles** we consider the approach of minimizing the expression of the error within the Krylov space for the entire class of functions: we **return to Zolotarev**.

## ◀ Back to Zolotarev

---

To **obtain the poles** we consider the approach of minimizing the expression of the error within the Krylov space for the entire class of functions: we **return to Zolotarev**.

🔧 Let us write **compactly**:  $\mathcal{W} = \mathcal{K}(A, \mathbf{v}, \Psi)$  for the rational Krylov subspace with poles  $\Psi$ . Then we can write **the approximation error** as:

$$\|\mathbf{x}_{\mathcal{W}} - \mathbf{x}\|_2 \leq 2 \cdot \|\mathbf{v}\|_2 \cdot \min_{r(z) \in \frac{\mathbb{P}_\ell}{\Psi}} \max_{z \in [a, b]} |f(z) - r(z)|.$$

where  $\mathbf{x}_{\mathcal{W}} = Uf(U^H A U)U^H \mathbf{v}$  for  $U$  an orthonormal basis of  $\mathcal{W}$ , and  $\mathbf{x} = f(A)\mathbf{v}$ .

## ◀ Back to Zolotarev

To **obtain the poles** we consider the approach of minimizing the expression of the error within the Krylov space for the entire class of functions: we **return to Zolotarev**.

🔧 Let us write **compactly**:  $\mathcal{W} = \mathcal{K}(A, \mathbf{v}, \Psi)$  for the rational Krylov subspace with poles  $\Psi$ . Then we can write **the approximation error** as:

$$\|\mathbf{x}_{\mathcal{W}} - \mathbf{x}\|_2 \leq 2 \cdot \|\mathbf{v}\|_2 \cdot \min_{r(z) \in \frac{\mathbb{P}_{\ell}}{\Psi}} \max_{z \in [a, b]} |f(z) - r(z)|.$$

where  $\mathbf{x}_{\mathcal{W}} = Uf(U^H A U)U^H \mathbf{v}$  for  $U$  an orthonormal basis of  $\mathcal{W}$ , and  $\mathbf{x} = f(A)\mathbf{v}$ .

👁 Now comes the clever observation, the function we want to approximate is of the form

$$f(A)\mathbf{v} = \int_0^{\infty} g(t, A) \mu(t) \, dt, \quad g(t, A) \in \{e^{-tA}, (tI + A)^{-1}\}$$

⇒ Since the **projection is linear** we need **poles** to **approximate uniformly well** (in  $t$ ) the matrix exponentials and resolvents.

# Cauchy-Stieltjes functions

For Cauchy-Stieltjes function, we just need the result for the resolvent function.

Theorem (Massei and Robol 2021, Theorem 1)

Let  $A$  be Hermitian positive definite with spectrum contained in  $[a, b]$  and  $U$  be an orthonormal basis of  $\mathcal{U}_{\mathcal{R}} = \mathcal{K}_{\ell}(A, \mathbf{v}, \Psi)$ . Then,  $\forall t \in [0, \infty)$ , we have the following inequality:

$$\|(tI + A)^{-1}\mathbf{v} - U(tI + A_{\ell})^{-1}\mathbf{v}_{\ell}\|_2 \leq \frac{2}{t + a} \|\mathbf{v}\|_2 \min_{r(z) \in \frac{\mathcal{P}_{\ell}}{\Psi}} \frac{\max_{z \in [a, b]} |r(z)|}{\min_{z \in (-\infty, 0]} |r(z)|}$$

where  $A_{\ell} = U^H A U$  and  $\mathbf{v}_{\ell} = U^H \mathbf{v}$ .

# Cauchy-Stieltjes functions

For Cauchy-Stieltjes function, we just need the result for the resolvent function.

Theorem (Massei and Robol 2021, Theorem 1)

Let  $A$  be Hermitian positive definite with spectrum contained in  $[a, b]$  and  $U$  be an orthonormal basis of  $\mathcal{U}_{\mathcal{R}} = \mathcal{K}_{\ell}(A, \mathbf{v}, \Psi)$ . Then,  $\forall t \in [0, \infty)$ , we have the following inequality:

$$\|(tI + A)^{-1}\mathbf{v} - U(tI + A_{\ell})^{-1}\mathbf{v}_{\ell}\|_2 \leq \frac{2}{t+a} \|\mathbf{v}\|_2 \min_{r(z) \in \frac{\mathcal{P}_{\ell}}{\Psi}} \frac{\max_{z \in [a,b]} |r(z)|}{\min_{z \in (-\infty, 0]} |r(z)|}$$

where  $A_{\ell} = U^H A U$  and  $\mathbf{v}_{\ell} = U^H \mathbf{v}$ .

✎ We got back to our favorite 4<sup>th</sup> problem of Zolotarev! Than we do not know how to solve in close form in general...

# Cauchy-Stieltjes functions

For Cauchy-Stieltjes function, we just need the result for the resolvent function.

Theorem (Massei and Robol 2021, Theorem 1)

Let  $A$  be Hermitian positive definite with spectrum contained in  $[a, b]$  and  $U$  be an orthonormal basis of  $\mathcal{U}_{\mathcal{R}} = \mathcal{K}_{\ell}(A, \mathbf{v}, \Psi)$ . Then,  $\forall t \in [0, \infty)$ , we have the following inequality:

$$\|(tI + A)^{-1}\mathbf{v} - U(tI + A_{\ell})^{-1}\mathbf{v}_{\ell}\|_2 \leq \frac{2}{t+a} \|\mathbf{v}\|_2 \min_{r(z) \in \frac{\mathcal{P}_{\ell}}{\Psi}} \frac{\max_{z \in [a,b]} |r(z)|}{\min_{z \in (-\infty, 0]} |r(z)|}$$

where  $A_{\ell} = U^H A U$  and  $\mathbf{v}_{\ell} = U^H \mathbf{v}$ .

✎ We got back to our favorite 4<sup>th</sup> problem of Zolotarev! Than we do not know how to solve in close form in general...

! this is not the general case, this is the case of two intervals  $[a, b]$  and  $(-\infty, 0]$  😊



## Solving this particular Zolotarev instance

---

### The Zolotarev constant

Let  $\Psi = \{\psi_1, \dots, \psi_\ell\} \subset \overline{\mathbb{C}}$  be a finite set, and  $I_1, I_2$  closed subsets of  $\overline{\mathbb{C}}$ . Then, we define

$$\theta_\ell(I_1, I_2, \Psi) = \min_{r(z) \in \frac{\mathcal{P}_\ell}{\Psi}} \frac{\max_{I_1} |r(z)|}{\min_{I_2} |r(z)|}.$$

# Solving this particular Zolotarev instance

## Theorem (Zolotarev)

Let  $I = [a, b]$ , with  $0 < a < b$ . Then

$$\min_{\Psi \subset \overline{\mathbb{C}}, |\Psi|=\ell} \theta_\ell(I, -I, \Psi) \leq 4\rho_{[a,b]}^\ell, \quad \rho_{[a,b]} = \exp\left(-\frac{\pi^2}{\log(4\kappa)}\right), \quad \kappa = \frac{b}{a}.$$

In addition, the optimal rational function  $r_\ell^{[a,b]}(z)$  that realizes the minimum has the form

$$r_\ell^{[a,b]}(z) = \frac{p_\ell^{[a,b]}(z)}{p_\ell^{[a,b]}(-z)}, \quad p_\ell^{[a,b]}(z) = \prod_{j=1}^{\ell} (z + \psi_{j,\ell}^{[a,b]}), \quad \psi_{j,\ell}^{[a,b]} \in -I.$$

We denote by  $\Psi_\ell^{[a,b]} = \{\psi_{1,\ell}^{[a,b]}, \dots, \psi_{\ell,\ell}^{[a,b]}\}$  the set of poles of  $r_\ell^{[a,b]}(z)$ .

## Solving this particular Zolotarev instance

### Theorem (Zolotarev)


Let  $I = [a, b]$ , with  $0 < a < b$ . Then

$$\min_{\Psi \subset \overline{\mathbb{C}}, |\Psi|=\ell} \theta_\ell(I, -I, \Psi) \leq 4\rho_{[a,b]}^\ell, \quad \rho_{[a,b]} = \exp\left(-\frac{\pi^2}{\log(4\kappa)}\right), \quad \kappa = \frac{b}{a}.$$

In addition, the optimal rational function  $r_\ell^{[a,b]}(z)$  that realizes the minimum has the form

$$r_\ell^{[a,b]}(z) = \frac{p_\ell^{[a,b]}(z)}{p_\ell^{[a,b]}(-z)}, \quad p_\ell^{[a,b]}(z) = \prod_{j=1}^{\ell} (z + \psi_{j,\ell}^{[a,b]}), \quad \psi_{j,\ell}^{[a,b]} \in -I.$$

We denote by  $\Psi_\ell^{[a,b]} = \{\psi_{1,\ell}^{[a,b]}, \dots, \psi_{\ell,\ell}^{[a,b]}\}$  the set of poles of  $r_\ell^{[a,b]}(z)$ .

 This solution is for  $I_1 = [a, b]$  and  $I_2 = [-b, -a]$ : we had  $[a, b]$  and  $(-\infty, 0]$ !

# Solving this particular Zolotarev instance

## The Zolotarev constant

Let  $\Psi = \{\psi_1, \dots, \psi_\ell\} \subset \overline{\mathbb{C}}$  be a finite set, and  $I_1, I_2$  closed subsets of  $\overline{\mathbb{C}}$ . Then, we define

$$\theta_\ell(I_1, I_2, \Psi) = \min_{r(z) \in \frac{\mathcal{P}_\ell}{\Psi}} \frac{\max_{I_1} |r(z)|}{\min_{I_2} |r(z)|}.$$

For any  $I_1, I_2$  be subsets of the complex plane, and  $\Psi \subset \overline{\mathbb{C}}$  we have


**shift invariance** For any  $t \in \mathbb{C}$ , it holds  $\theta_\ell(I_1 + t, I_2 + t, \Psi + t) = \theta_\ell(I_1, I_2, \Psi)$ .

**monotonicity**  $\theta_\ell(I_1, I_2, \Psi)$  is monotonic with respect to the inclusion on the parameters  $I_1$  and  $I_2$ :  $I_1 \subseteq I'_1, I_2 \subseteq I'_2 \implies \theta_\ell(I_1, I_2, \Psi) \leq \theta_\ell(I'_1, I'_2, \Psi)$ .

**Möbius invariance** If  $M(z)$  is a Möbius transform, that is a rational function

$M(z) = (\alpha z + \beta)/(\gamma z + \delta)$  with  $\alpha\delta \neq \beta\gamma$ , then

$$\theta_\ell(I_1, I_2, \Psi) = \theta_\ell(M(I_1), M(I_2), M(\Psi)).$$

 This solution is for  $I_1 = [a, b]$  and  $I_2 = [-b, -a]$ : we had  $[a, b]$  and  $(-\infty, 0]!$

## Solving this particular Zolotarev instance

We just need to build the right Möbius transform to map

$$(-\infty, 0] \cup [a, b] \mapsto -I \cup I, \quad I = [a', b'], \quad 0 < a' < b'.$$

Lemma (Massei and Robol [2021](#), Lemma 4)

The Möbius transformation

$$T_C(z) = \frac{\Delta + z - b}{\Delta - z + b}, \quad \Delta = \sqrt{b^2 - ab},$$

maps  $[-\infty, 0] \cup [a, b]$  into  $[-1, -\hat{a}] \cup [\hat{a}, 1]$ , with  $\hat{a} = \frac{\Delta + a - b}{\Delta - a + b} = \frac{b - \Delta}{\Delta + b}$ . The inverse map  $T_C(z)^{-1}$  is given by:

$$T_C^{-1}(z) = \frac{(b + \Delta)z + b - \Delta}{1 + z}.$$

Moreover, for any  $0 < a < b$  it holds  $\hat{a}^{-1} \leq \frac{4b}{a}$ , and therefore  $\rho_{[\hat{a}, 1]} \leq \rho_{[a, 4b]}$ .

# Cauchy-Stieltjes functions

- ⚙️ We map the interval  $[a, b]$  to  $[\hat{a}, 1]$ ,
- 🔧 solve *explicitly* the Zolotarev problem there,
- 📖 read the poles for our problem.

## Proposition (Massei and Robol 2021, Corollary 4)

Let  $f(z)$  be a Cauchy-Stieltjes function,  $A$  be Hermitian positive definite with spectrum contained in  $[a, b]$ ,  $U$  be an orthonormal basis of  $\mathcal{K}_\ell(A, \mathbf{v}, \Psi_{C,\ell}^{[a,b]})$  with  $\Psi_{C,\ell}^{[a,b]}$  given by

$$\Psi_{C,\ell}^{[a,b]} = T_C^{-1}(\Psi_\ell^{[\hat{a},1]})$$

and  $\mathbf{x}_\ell = Uf(A_\ell)\mathbf{v}_\ell$  with  $A_\ell = U^H A U$  and  $\mathbf{v}_\ell = U^H \mathbf{v}$ . Then

$$\|f(A)\mathbf{v} - \mathbf{x}_\ell\|_2 \leq 8f(a)\|\mathbf{v}\|_2 \rho_{[a,4b]}^\ell = 8f(a) \exp\left(-\ell \frac{\pi^2}{\log(16b/a)}\right).$$

# Nesting the poles

The poles built this way are still **not nested**. In (Massei and Robol 2021) a technique called method of equidistributed sequences (EDS) is proposed to generate them:

1. Select  $\zeta \in \mathbb{R}^+ \setminus \mathbb{Q}$  and generate the sequence  $\{s_j\}_{j \in \mathbb{N}} = \{0, \zeta - \lfloor \zeta \rfloor, 2\zeta - \lfloor 2\zeta \rfloor, 3\zeta - \lfloor 3\zeta \rfloor, \dots\}$ , where  $\lfloor \cdot \rfloor$  indicates the greatest integer less than or equal to the argument; this sequence has as asymptotic distribution (in the sense of EDS) the Lebesgue measure on  $[0, 1]$ .
2. Compute the sequence  $\{t_j\}_{j \in \mathbb{N}}$  such that  $g(t_j) = s_j$  where

$$g(t) = \frac{1}{2M} \int_{a^2}^t \frac{dy}{\sqrt{(y - a^2)y(1 - y)}}, \quad M = \int_0^1 \frac{dy}{\sqrt{(1 - y^2)(1 - (1 - a^2)y^2)}},$$

3. Define  $\tilde{\sigma}_j = \sqrt{t_j}$ .

➡ The EDS associated with  $\Psi_\ell^{[a,b]}$ ,  $\Psi_{C,\ell}^{[a,b]}$  are obtained by applying either a scaling or the Möbius transformation to the EDS for  $\Psi_\ell^{[a,1]}$ .

# Brute force approaches

---

It is also possible to try and solve numerically rational approximation problems.

**RKFIT** (Berljafa and Güttel 2017) Is an iterative method for solving rational Least-Square problems,  $\{A, F\} \in \mathbb{C}^{n \times n}$  and  $\mathbf{b} \in \mathbb{C}^n$  find a ration function  $r$  such that

$$\|F\mathbf{b} - r(A)\mathbf{b}\|_2^2 \rightarrow \min.$$

**AAA** (Nakatsukasa, Sète, and Trefethen 2018) Find a representation of the rational approximant in barycentric form with interpolation at certain support points while performing a greedy selection of them to avoid exponential instabilities.

If we have an idea of *where* the approximation should work, these approaches deliver reasonably good results.



# An Application to Complex Networks

---

The **spectral definition** makes the procedure ideal also in more *exotic* cases.

# An Application to Complex Networks

---

The **spectral definition** makes the procedure ideal also in more *exotic* cases.

⚙️ A *weighted directed graph* (digraph) is a pair  $G = (V, E, W)$ , where  $V = \{v_1, \dots, v_n\}$  is a **set of nodes** (or vertices), and  $E \subseteq V \times V$  is a **set of ordered pairs** of nodes called **edges**, and  $W \in \mathbb{R}^{n \times n}$  such that  $(W)_{ij} \neq 0$  iff  $(v_i, v_j) \in E$ .

# An Application to Complex Networks

The **spectral definition** makes the procedure ideal also in more *exotic* cases.

In matrix language

⚙️ A *weighted directed graph* (digraph) is a pair  $G = (V, E, W)$ , where  $V = \{v_1, \dots, v_n\}$  is a **set of nodes** (or vertices), and  $E \subseteq V \times V$  is a **set of ordered pairs** of nodes called **edges**, and  $W \in \mathbb{R}^{n \times n}$  such that  $(W)_{ij} \neq 0$  iff  $(v_i, v_j) \in E$ .

🔧 If all the weights are equal to one, the **adjacency matrix**  $A \in \mathbb{R}^{n \times n}$  is

$$(A)_{ij} = a_{ij} = \begin{cases} 1, & \text{if } (v_i, v_j) \in E, \\ 0, & \text{otherwise.} \end{cases}$$

otherwise,  $A \equiv W$ .

⚙️ We call *in-degrees* and *out-degrees*

$$d_i^{(\text{in})} = \deg_{\text{in}}(v_i) = \sum_{j: (v_j, v_i) \in E} w_{j,i},$$

$$d_i^{(\text{out})} = \deg_{\text{out}}(v_i) = \sum_{j: (v_i, v_j) \in E} w_{i,j},$$

# An Application to Complex Networks

The **spectral definition** makes the procedure ideal also in more *exotic* cases.

⚙️ A *weighted directed graph* (digraph) is a pair  $G = (V, E, W)$ , where  $V = \{v_1, \dots, v_n\}$  is a **set of nodes** (or vertices), and  $E \subseteq V \times V$  is a **set of ordered pairs** of nodes called **edges**, and  $W \in \mathbb{R}^{n \times n}$  such that  $(W)_{ij} \neq 0$  iff  $(v_i, v_j) \in E$ .

⚙️ We call *in-degrees* and *out-degrees*

$$d_i^{(\text{in})} = \deg_{\text{in}}(v_i) = \sum_{j: (v_j, v_i) \in E} w_{j,i},$$

$$d_i^{(\text{out})} = \deg_{\text{out}}(v_i) = \sum_{j: (v_i, v_j) \in E} w_{i,j},$$

In matrix language

🔧 If all the weights are equal to one, the **adjacency matrix**  $A \in \mathbb{R}^{n \times n}$  is

$$(A)_{ij} = a_{ij} = \begin{cases} 1, & \text{if } (v_i, v_j) \in E, \\ 0, & \text{otherwise.} \end{cases}$$

otherwise,  $A \equiv W$ .

🔧 **Degree diagonal matrices**

$$\begin{aligned} D_{\text{in}} &= \text{diag}(\deg_{\text{in}}(v_1), \dots, \deg_{\text{in}}(v_n)) \\ &= \text{diag}(d_1^{(\text{in})}, \dots, d_n^{(\text{in})}), \end{aligned}$$

$$\begin{aligned} D_{\text{out}} &= \text{diag}(\deg_{\text{out}}(v_1), \dots, \deg_{\text{out}}(v_n)) \\ &= \text{diag}(d_1^{(\text{out})}, \dots, d_n^{(\text{out})}). \end{aligned}$$

# Laplacians on Graphs

## Undirected case

Let  $G = (V, E)$  be a weighted undirected graph with weight matrix  $W$ , weighted degree matrix  $D$  and weighted incidence matrix  $B$ . Then the **graph Laplacian**  $L$  of  $G$  is

$$L = D - W.$$

The *normalized random walk* version of the graph Laplacian is

$$D^{-1}L = I - D^{-1}W,$$

where  $I$  is the identity matrix. Observe that  $D^{-1}W$  is a row-stochastic matrix, i.e. it is nonnegative with row sums equal to 1. The *normalized symmetric* version is

$$D^{-\frac{1}{2}}LD^{-\frac{1}{2}} = I - D^{-\frac{1}{2}}WD^{-\frac{1}{2}}.$$


If  $G$  is unweighted then  $W = A$  in the above definitions. Here we assume that every vertex has nonzero degree.

# Laplacians on Graphs

## Directed case

Let  $G = (V, E, W)$  be a weighted directed graph, with degree matrices  $D_{\text{out}}$  and  $D_{\text{in}}$ . The **nonnormalized directed graph Laplacian**  $L_{\text{out}}$  and  $L_{\text{in}}$  of  $G$  are

$$L_{\text{out}} = D_{\text{out}} - W, \quad L_{\text{in}} = D_{\text{in}} - W.$$


 To define the **normalized versions**, we need to invert either the  $D_{\text{in}}$  or the  $D_{\text{out}}$  matrices, but the absence of isolated vertices is no longer sufficient to ensure this!

# Laplacians on Graphs

## Directed case

Let  $G = (V, E, W)$  be a weighted directed graph, with degree matrices  $D_{\text{out}}$  and  $D_{\text{in}}$ . The **nonnormalized directed graph Laplacian**  $L_{\text{out}}$  and  $L_{\text{in}}$  of  $G$  are

$$L_{\text{out}} = D_{\text{out}} - W, \quad L_{\text{in}} = D_{\text{in}} - W.$$

 To define the **normalized versions**, we need to invert either the  $D_{\text{in}}$  or the  $D_{\text{out}}$  matrices, but the absence of isolated vertices is no longer sufficient to ensure this!

It is **interesting to look at diffusion on graphs**:

$$\begin{aligned} &\text{find } u : [0, T] \longrightarrow \mathbb{R}^n \\ &\text{s.t. } \begin{cases} \frac{d}{dt} u(t) = -\kappa L_{\cdot/\text{in}/\text{out}} u(t), & t \in (0, T], \\ u(0) = u_0, & \text{prescribed,} \end{cases} \end{aligned}$$

# Laplacians on Graphs

## Directed case

Let  $G = (V, E, W)$  be a weighted directed graph, with degree matrices  $D_{\text{out}}$  and  $D_{\text{in}}$ . The **nonnormalized directed graph Laplacian**  $L_{\text{out}}$  and  $L_{\text{in}}$  of  $G$  are

$$L_{\text{out}} = D_{\text{out}} - W, \quad L_{\text{in}} = D_{\text{in}} - W.$$

⚠ To define the **normalized versions**, we need to invert either the  $D_{\text{in}}$  or the  $D_{\text{out}}$  matrices, but the absence of isolated vertices is no longer sufficient to ensure this!

It is **interesting to look at diffusion on graphs**:

$$\begin{aligned} &\text{find } u : [0, T] \longrightarrow \mathbb{R}^n \\ &\text{s.t. } \begin{cases} \frac{d}{dt}u(t) = -\kappa L_{\cdot/\text{in}/\text{out}}^{\alpha} u(t), & t \in (0, T], \\ u(0) = u_0, & \text{prescribed,} \end{cases} \quad \alpha \in (0, 1], \end{aligned}$$

$\Rightarrow$  it *could be* interesting to look at **fractional diffusion** on graphs.



# Laplacians on Graphs

---

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

# Laplacians on Graphs

---

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

# Laplacians on Graphs


---

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

  $L_{\text{out}}$  is a **singular  $M$ -matrix**,

  $L_{\text{out}}\mathbf{1} = \mathbf{0}$ ,

  $0$  is a *semisimple* eigenvalue of  $L_{\text{out}}$ .

# Laplacians on Graphs

---

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

🏛️  $L_{\text{out}}$  is a **singular  $M$ -matrix**,

🏛️  $L_{\text{out}} \mathbf{1} = \mathbf{0}$ ,

🏛️ 0 is a *semisimple* eigenvalue of  $L_{\text{out}}$ .

⚖️ We need to prove that  $L_{\text{out}}^\alpha$  can be **defined** and **respect all the properties**.

# Laplacians on Graphs

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

🏛️  $L_{\text{out}}$  is a **singular**  $M$ -matrix,

🏛️  $L_{\text{out}} \mathbf{1} = \mathbf{0}$ ,

🏛️ 0 is a *semisimple* eigenvalue of  $L_{\text{out}}$ .

⚖️ We need to prove that  $L_{\text{out}}^\alpha$  can be **defined** and **respect all the properties**.

Proposition (Benzi, Bertaccini, et al. 2020)


Given a weighted graph  $G = (V, E, W)$  and its Laplacian with respect to the out degree  $L_{\text{out}}$ , the function  $f(x) = x^\alpha$  is defined on the spectrum of  $L_{\text{out}}$  and induces a matrix function for all  $\alpha \in (0, 1]$ .


# Laplacians on Graphs

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

  $L_{\text{out}}$  is a **singular  $M$ -matrix**,

  $L_{\text{out}} \mathbf{1} = \mathbf{0}$ ,

  $0$  is a *semisimple* eigenvalue of  $L_{\text{out}}$ .

 We need to prove that  $L_{\text{out}}^\alpha$  can be **defined** and **respect all the properties**.

Proposition (Benzi, Bertaccini, et al. 2020)

If  $A$  is a singular  $M$ -matrix with  $0$  as a semisimple eigenvalue, then there exists a determination of  $A^\alpha$  for every  $\alpha \in (0, 1]$  that is a singular  $M$ -matrix.


# Laplacians on Graphs

If  $G$  is *undirected*, i.e.,  $L = L^T$ , everything follows by **diagonalization**, see, e.g., (Riascos and Mateos 2014).

If  $L$  is either  $L_{\text{out}}$  or  $L_{\text{in}}$  this **needs more care**.

  $L_{\text{out}}$  is a **singular  $M$ -matrix**,


  $L_{\text{out}} \mathbf{1} = \mathbf{0}$ ,

  $0$  is a *semisimple* eigenvalue of  $L_{\text{out}}$ .

 We need to prove that  $L_{\text{out}}^\alpha$  can be **defined** and **respect all the properties**.

Proposition (Benzi, Bertaccini, et al. 2020)

If  $A$  is a singular  $M$ -matrix with  $0$  as a semisimple eigenvalue, then there exists a determination of  $A^\alpha$  for every  $\alpha \in (0, 1]$  that is a singular  $M$ -matrix.

 We could also investigate the **the decay of the entries** of the fractional power, but leave the subject aside and refer to (Benzi, Bertaccini, et al. 2020).

# Laplacian on Graphs: computation

---

⚠ For the computation of the products  $L_{\text{out}}^\alpha \mathbf{v}$  it is necessary to **modify the strategies** we have seen: all the bounds and constructions required that **0 was not in the spectrum**.



# Laplacian on Graphs: computation

⚠ For the computation of the products  $L_{\text{out}}^\alpha \mathbf{v}$  it is necessary to **modify the strategies** we have seen: all the bounds and constructions required that **0 was not in the spectrum**.

🔧 In (Benzi and Simunec 2022) different strategies for accommodating this feature of  $L_{\text{out}}$  are investigated:

1 Use a **rank-one** shift, since the right and left eigenvectors  $\mathbf{1}$  and  $\vec{z}$  of  $L_{\text{out}}$  can be easily computed, we compute

$$f(L^T)\mathbf{b} = f(L^T + \theta \mathbf{z}\mathbf{1}^T)\mathbf{b} + [f(0) - f(\theta)]\mathbf{z}, \text{ for any } \theta > 0,$$

and in the rational Krylov subspace we solve the linear system at the same cost at which we solve the ones for  $L^T$  via Sherman-Morrison:

$$(L^T + \theta \mathbf{z}\mathbf{1}^T - \xi I)^{-1} = (L^T - \xi I)^{-1} + \frac{\theta}{\xi(\theta - \xi)} \mathbf{z}\mathbf{1}^T.$$

# Laplacian on Graphs: computation

⚠ For the computation of the products  $L_{\text{out}}^\alpha \mathbf{v}$  it is necessary to **modify the strategies** we have seen: all the bounds and constructions required that **0 was not in the spectrum**.

🔧 In (Benzi and Simunec 2022) different strategies for accommodating this feature of  $L_{\text{out}}$  are investigated:

1 Use a **rank-one** shift, since the right and left eigenvectors  $\mathbf{1}$  and  $\vec{z}$  of  $L_{\text{out}}$  can be easily computed, we compute

$$f(L^T)\mathbf{b} = f(L^T + \theta \mathbf{z} \mathbf{1}^T)\mathbf{b} + [f(0) - f(\theta)]\mathbf{z}, \text{ for any } \theta > 0,$$

and in the rational Krylov subspace we solve the linear system at the same cost at which we solve the ones for  $L^T$  by doing

$$(L^T + \theta \mathbf{z} \mathbf{1}^T - \xi I)^{-1} \mathbf{w} = \boldsymbol{\psi} + \frac{\mathbf{1}^T \mathbf{w}}{\theta - \xi} \mathbf{z} \quad \text{and} \quad (L^T - \xi I) \boldsymbol{\psi} = \mathbf{w} - (\mathbf{1}^T \mathbf{w}) \mathbf{z},$$

to **avoid cancellation** for  $\xi \approx 0$ .

# Laplacian on Graphs: computation

⚠ For the computation of the products  $L_{\text{out}}^\alpha \mathbf{v}$  it is necessary to **modify the strategies** we have seen: all the bounds and constructions required that **0 was not in the spectrum**.

🔧 In (Benzi and Simunec 2022) different strategies for accommodating this feature of  $L_{\text{out}}$  are investigated:

1️⃣ Use a **rank-one** shift, since the right and left eigenvectors  $\mathbf{1}$  and  $\vec{z}$  of  $L_{\text{out}}$  can be easily computed, we compute

$$f(L^T)\mathbf{b} = f(L^T + \theta \mathbf{z}\mathbf{1}^T)\mathbf{b} + [f(0) - f(\theta)]\mathbf{z}, \text{ for any } \theta > 0.$$

1️⃣ Project  $L$  on the  $n-1$  dimensional subspace  $\mathcal{S} = \text{Span}\{\mathbf{1}\}^\perp = \text{Range}(\tilde{Q})$  and compute


$$\begin{aligned} f(L^T)\mathbf{b} &= f(L^T)\mathbf{v} + \beta f(L^T)\mathbf{z} && \leftarrow 0 \neq \beta = \mathbf{1}^T \mathbf{b} \text{ and } \mathbf{b} = \mathbf{v} + \beta \mathbf{z} \text{ for } \mathbf{v} \perp \mathbf{1} \\ &= Qf(Q^T L^T Q)Q^T \mathbf{v} + \beta f(0)\mathbf{z} && \leftarrow QQ^T = I - \mathbf{1}\mathbf{1}^T/n, \quad Q = [\tilde{Q}, \mathbf{1}/\sqrt{n}]. \end{aligned}$$

❗  $Q$  can be built so that  $\{Q, Q^T\}\mathbf{v}$  costs  $O(n)$ .

# A gallery of open problems

---



*“When sorrows come, they come  
not single spies, but in battalions”*  
Hamlet, Act IV, Scene V.

-  Of the many problems we have discussed along the way, one that came back many times was the **selection of optimal poles** for the different matrix-equation/Rational Krylov based solvers (e.g., *all-at-once*, multi-dimensional approaches);

# A gallery of open problems

---

*“When sorrows come, they come  
not single spies, but in battalions”*  
Hamlet, Act IV, Scene V.



-  Of the many problems we have discussed along the way, one that came back many times was the **selection of optimal poles** for the different matrix-equation/Rational Krylov based solvers (e.g., *all-at-once*, multi-dimensional approaches);
-  Inventing **reduced memory** methods for the integration of fractional partial differential equations in time and space, i.e.,

$${}^{\text{CA}}D_t^\alpha \mathbf{u} = \mathcal{L}(\mathbf{u}; t), \quad \mathcal{L} \text{ non linear, and fractional;}$$


# A gallery of open problems

---

*“When sorrows come, they come  
not single spies, but in battalions”*  
Hamlet, Act IV, Scene V.

-  Of the many problems we have discussed along the way, one that came back many times was the **selection of optimal poles** for the different matrix-equation/Rational Krylov based solvers (e.g., *all-at-once*, multi-dimensional approaches);
-  Inventing **reduced memory** methods for the integration of fractional partial differential equations in time and space, i.e.,



$${}^{CA}D_t^\alpha \mathbf{u} = \mathcal{L}(\mathbf{u}; t), \quad \mathcal{L} \text{ non linear, and fractional;}$$

-  **Error analysis** entangling convergence of the Rational Krylov method and Finite Element (Isogeometric) Discretizations for FPDEs;



# A gallery of open problems

---

*“When sorrows come, they come  
not single spies, but in battalions”*  
Hamlet, Act IV, Scene V.

-  Of the many problems we have discussed along the way, one that came back many times was the **selection of optimal poles** for the different matrix-equation/Rational Krylov based solvers (e.g., *all-at-once*, multi-dimensional approaches);
-  Inventing **reduced memory** methods for the integration of fractional partial differential equations in time and space, i.e.,

$${}^{CA}D_t^\alpha \mathbf{u} = \mathcal{L}(\mathbf{u}; t), \quad \mathcal{L} \text{ non linear, and fractional;}$$

-  **Error analysis** entangling convergence of the Rational Krylov method and Finite Element (Isogeometric) Discretizations for FPDEs;
-  Solving FPDEs on **unlimited spatial domains**.

# Fractional Schrödinger equation

---

As we have discussed at the beginning of the lecture, there are several formulations of the Fractional Laplacian that should be naturally **considered on the whole space**.

An example is the **Schrödinger equation**

$$i\hbar^\beta {}^{CA}D^\beta \psi = -D_\alpha (-\hbar^2 \Delta)^{\alpha/2} \psi + V(\mathbf{x}, t) \psi,$$

that is naturally defined on the whole space.

To treat it numerically, the usual procedure is to couple it with **artificial boundary conditions of absorbing type**. It might be of interest to have **numerical methods** that can work with **infinite** or **semi-infinite matrices** that do not need this artificial correction.



# Conclusions

---





- ⚙ We focused on *few discretization*, there are many other viable approaches (*collocation, finite elements, IgA,...*). Most of the reasoning we did can be **adapted to these other cases**.
- 🔧 There are **other classical problems** that admits a fractional extension, e.g., optimal control, model order reduction, eigenvalue problems,...

*"The universe (which others call the Library) is composed of an indefinite and perhaps infinite number of hexagonal galleries, with vast air shafts between, surrounded by very low railings. From any of the hexagons one can see, interminably, the upper and lower floors. The distribution of the galleries is invariable."*

Jorge Luis Borges, The Library of Babel.





# Bibliography I

---

-  Aceto, L., D. Bertaccini, et al. (2019). “Rational Krylov methods for functions of matrices with applications to fractional partial differential equations”. In: *J. Comput. Phys.* 396, pp. 470–482. ISSN: 0021-9991. DOI: [10.1016/j.jcp.2019.07.009](https://doi.org/10.1016/j.jcp.2019.07.009). URL: <https://doi.org/10.1016/j.jcp.2019.07.009>.
-  Aceto, L. and P. Novati (2018). “Efficient implementation of rational approximations to fractional differential operators”. In: *J. Sci. Comput.* 76.1, pp. 651–671. ISSN: 0885-7474. DOI: [10.1007/s10915-017-0633-2](https://doi.org/10.1007/s10915-017-0633-2). URL: <https://doi.org/10.1007/s10915-017-0633-2>.
-  — (2019). “Rational approximations to fractional powers of self-adjoint positive operators”. In: *Numer. Math.* 143.1, pp. 1–16. ISSN: 0029-599X. DOI: [10.1007/s00211-019-01048-4](https://doi.org/10.1007/s00211-019-01048-4). URL: <https://doi.org/10.1007/s00211-019-01048-4>.
-  — (2022). “Fast and accurate approximations to fractional powers of operators”. In: *IMA J. Numer. Anal.* 42.2, pp. 1598–1622. ISSN: 0272-4979. DOI: [10.1093/imanum/drab002](https://doi.org/10.1093/imanum/drab002). URL: <https://doi.org/10.1093/imanum/drab002>.





# Bibliography II

---

-  Andreu-Vaillo, F. et al. (2010). *Nonlocal diffusion problems*. Vol. 165. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI; Real Sociedad Matemática Española, Madrid, pp. xvi+256. ISBN: 978-0-8218-5230-9. DOI: [10.1090/surv/165](https://doi.org/10.1090/surv/165). URL: <https://doi.org/10.1090/surv/165>.
-  Benzi, M., D. Bertaccini, et al. (2020). “Non-local network dynamics via fractional graph Laplacians”. In: *J. Complex Netw.* 8.3, cnaa017, 29. ISSN: 2051-1310. DOI: [10.1093/comnet/cnaa017](https://doi.org/10.1093/comnet/cnaa017). URL: <https://doi.org/10.1093/comnet/cnaa017>.
-  Benzi, M. and I. Simunec (2022). “Rational Krylov methods for fractional diffusion problems on graphs”. In: *BIT* 62.2, pp. 357–385. ISSN: 0006-3835. DOI: [10.1007/s10543-021-00881-0](https://doi.org/10.1007/s10543-021-00881-0). URL: <https://doi.org/10.1007/s10543-021-00881-0>.
-  Berljafa, M. and S. Güttel (2017). “The RKFIT algorithm for nonlinear rational approximation”. In: *SIAM J. Sci. Comput.* 39.5, A2049–A2071. ISSN: 1064-8275. DOI: [10.1137/15M1025426](https://doi.org/10.1137/15M1025426). URL: <https://doi.org/10.1137/15M1025426>.






# Bibliography III

---

-  Bhatia, R. (1997). *Matrix analysis*. Vol. 169. Graduate Texts in Mathematics. Springer-Verlag, New York, pp. xii+347. ISBN: 0-387-94846-5. DOI: [10.1007/978-1-4612-0653-8](https://doi.org/10.1007/978-1-4612-0653-8). URL: <https://doi.org/10.1007/978-1-4612-0653-8>.
-  Bonito, A. and J. E. Pasciak (2015). “Numerical approximation of fractional powers of elliptic operators”. In: *Math. Comp.* 84.295, pp. 2083–2110. ISSN: 0025-5718. DOI: [10.1090/S0025-5718-2015-02937-8](https://doi.org/10.1090/S0025-5718-2015-02937-8). URL: <https://doi.org/10.1090/S0025-5718-2015-02937-8>.
-  Braess, D. (1986). *Nonlinear approximation theory*. Vol. 7. Springer Series in Computational Mathematics. Springer-Verlag, Berlin, pp. xiv+290. ISBN: 3-540-13625-8. DOI: [10.1007/978-3-642-61609-9](https://doi.org/10.1007/978-3-642-61609-9). URL: <https://doi.org/10.1007/978-3-642-61609-9>.
-  Brezinski, C. (1980). *Padé-type approximation and general orthogonal polynomials*. Vol. 50. International Series of Numerical Mathematics. Birkhäuser Verlag, Basel-Boston, Mass., p. 250. ISBN: 3-7643-1100-2.






# Bibliography IV

---

-  Davis, P. J. and P. Rabinowitz (1984). *Methods of numerical integration*. Second. Computer Science and Applied Mathematics. Academic Press, Inc., Orlando, FL, pp. xiv+612. ISBN: 0-12-206360-0.
-  Frommer, A., S. Güttel, and M. Schweitzer (2014). “Efficient and stable Arnoldi restarts for matrix functions based on quadrature”. In: *SIAM J. Matrix Anal. Appl.* 35.2, pp. 661–683. ISSN: 0895-4798. DOI: [10.1137/13093491X](https://doi.org/10.1137/13093491X). URL: <https://doi.org/10.1137/13093491X>.
-  Harizanov, S. et al. (2020). “Analysis of numerical methods for spectral fractional elliptic equations based on the best uniform rational approximation”. In: *J. Comput. Phys.* 408, pp. 109285, 21. ISSN: 0021-9991. DOI: [10.1016/j.jcp.2020.109285](https://doi.org/10.1016/j.jcp.2020.109285). URL: <https://doi.org/10.1016/j.jcp.2020.109285>.
-  Hofreither, C. (2021). “An algorithm for best rational approximation based on barycentric rational interpolation”. In: *Numer. Algorithms* 88.1, pp. 365–388. ISSN: 1017-1398. DOI: [10.1007/s11075-020-01042-0](https://doi.org/10.1007/s11075-020-01042-0). URL: <https://doi.org/10.1007/s11075-020-01042-0>.
-  Ilic, M. et al. (2005). “Numerical approximation of a fractional-in-space diffusion equation. I”. In: *Fract. Calc. Appl. Anal.* 8.3, pp. 323–341. ISSN: 1311-0454.




# Bibliography V

---

-  Ilic, M. et al. (2006). “Numerical approximation of a fractional-in-space diffusion equation. II. With nonhomogeneous boundary conditions”. In: *Fract. Calc. Appl. Anal.* 9.4, pp. 333–349. ISSN: 1311-0454.
-  Kwaśnicki, M. (2017). “Ten equivalent definitions of the fractional Laplace operator”. In: *Fract. Calc. Appl. Anal.* 20.1, pp. 7–51. ISSN: 1311-0454. DOI: [10.1515/fca-2017-0002](https://doi.org/10.1515/fca-2017-0002). URL: <https://doi.org/10.1515/fca-2017-0002>.
-  Lischke, A. et al. (2020). “What is the fractional Laplacian? A comparative review with new results”. In: *J. Comput. Phys.* 404, pp. 109009, 62. ISSN: 0021-9991. DOI: [10.1016/j.jcp.2019.109009](https://doi.org/10.1016/j.jcp.2019.109009). URL: <https://doi.org/10.1016/j.jcp.2019.109009>.
-  Massei, S. and L. Robol (2021). “Rational Krylov for Stieltjes matrix functions: convergence and pole selection”. In: *BIT* 61.1, pp. 237–273. ISSN: 0006-3835. DOI: [10.1007/s10543-020-00826-z](https://doi.org/10.1007/s10543-020-00826-z). URL: <https://doi.org/10.1007/s10543-020-00826-z>.
-  Musina, R. and A. I. Nazarov (2014). “On fractional Laplacians”. In: *Comm. Partial Differential Equations* 39.9, pp. 1780–1790. ISSN: 0360-5302. DOI: [10.1080/03605302.2013.864304](https://doi.org/10.1080/03605302.2013.864304). URL: <https://doi.org/10.1080/03605302.2013.864304>.

# Bibliography VI

---

-  Nakatsukasa, Y., O. Sète, and L. N. Trefethen (2018). “The AAA algorithm for rational approximation”. In: *SIAM J. Sci. Comput.* 40.3, A1494–A1522. ISSN: 1064-8275. DOI: [10.1137/16M1106122](https://doi.org/10.1137/16M1106122). URL: <https://doi.org/10.1137/16M1106122>.
-  Riascos, A. and J. Mateos (2014). “Fractional dynamics on networks: Emergence of anomalous diffusion and Lévy flights”. In: *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics* 90.3. cited By 49. DOI: [10.1103/PhysRevE.90.032809](https://doi.org/10.1103/PhysRevE.90.032809). URL: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-84907266357&doi=10.1103%2fPhysRevE.90.032809&partnerID=40&md5=be06b3148ba7bc17a50f52854beb9fac>.
-  Stahl, H. R. (2003). “Best uniform rational approximation of  $x^\alpha$  on  $[0, 1]$ ”. In: *Acta Math.* 190.2, pp. 241–306. ISSN: 0001-5962. DOI: [10.1007/BF02392691](https://doi.org/10.1007/BF02392691). URL: <https://doi.org/10.1007/BF02392691>.