

Proof that the Variance Estimator using $\frac{1}{N}$ is Biased

Proof

Given a random sample X_1, X_2, \dots, X_N drawn from a population with mean μ and variance σ^2 , we consider the biased variance estimator:

$$\hat{\sigma}_X^2 = \frac{1}{N} \sum_{i=1}^N (X_i - \hat{\mu}_X)^2.$$

where the sample mean is given by:

$$\hat{\mu}_X = \frac{1}{N} \sum_{i=1}^N X_i.$$

Our goal is to compute $\mathbb{E}[\hat{\sigma}_X^2]$ and compare it to σ^2 .

Step 1: Expanding the Squared Differences

Expanding the squared term:

$$(X_i - \hat{\mu}_X)^2 = X_i^2 - 2X_i\hat{\mu}_X + \hat{\mu}_X^2.$$

Summing over all i :

$$\sum_{i=1}^N (X_i - \hat{\mu}_X)^2 = \sum_{i=1}^N X_i^2 - 2\hat{\mu}_X \sum_{i=1}^N X_i + N\hat{\mu}_X^2.$$

Since $\sum_{i=1}^N X_i = N\hat{\mu}_X$, substitute and obtain:

$$\sum_{i=1}^N (X_i - \hat{\mu}_X)^2 = \sum_{i=1}^N X_i^2 - N\hat{\mu}_X^2.$$

Dividing by N to obtain $\hat{\sigma}_X^2$:

$$\hat{\sigma}_X^2 = \frac{1}{N} \left(\sum_{i=1}^N X_i^2 - N\hat{\mu}_X^2 \right).$$

Step 2: Taking Expectation

Taking expectation on both sides:

$$\mathbb{E}[\hat{\sigma}_X^2] = \frac{1}{N} \left(\sum_{i=1}^N \mathbb{E}[X_i^2] - N\mathbb{E}[\hat{\mu}_X^2] \right).$$

Since each X_i has mean μ and variance σ^2 , we use:

$$\mathbb{E}[X_i^2] = \sigma^2 + \mu^2.$$

This is because $\text{Var}(X_i) = \mathbb{E}(X_i^2) - (\mathbb{E}(X_i))^2$ —a property we have proven earlier about the variance (see lecture 8).

Summing over all N :

$$\sum_{i=1}^N \mathbb{E}[X_i^2] = N(\sigma^2 + \mu^2).$$

For $\mathbb{E}[\hat{\mu}_X^2]$, we use:

$$\mathbb{E}[\hat{\mu}_X^2] = \text{Var}(\hat{\mu}_X) + (\mathbb{E}[\hat{\mu}_X])^2 = \frac{\sigma^2}{N} + \mu^2.$$

Substituting these into our expectation:

$$\mathbb{E}[\hat{\sigma}_X^2] = \frac{1}{N} \left(N(\sigma^2 + \mu^2) - N \left(\frac{\sigma^2}{N} + \mu^2 \right) \right).$$

Expanding:

$$\begin{aligned} \mathbb{E}[\hat{\sigma}_X^2] &= \frac{1}{N} (N\sigma^2 + N\mu^2 - \sigma^2 - N\mu^2) \\ &= \frac{1}{N} (N\sigma^2 - \sigma^2) \\ &= \sigma^2 \left(1 - \frac{1}{N} \right). \end{aligned}$$

Step 3: Computing the Bias

Since the true variance is σ^2 , the bias of $\hat{\sigma}_X^2$ is:

$$\begin{aligned} \mathbb{E}[\hat{\sigma}_X^2] - \sigma^2 &= \sigma^2 \left(1 - \frac{1}{N} \right) - \sigma^2 \\ &= -\frac{\sigma^2}{N}. \end{aligned}$$

Since this bias is negative, $\hat{\sigma}_X^2$ underestimates the true variance, proving that it is biased.

Conclusion

To obtain an unbiased estimator, we must instead use:

$$\bar{\sigma}_X^2 = \frac{1}{N-1} \sum_{i=1}^N (X_i - \hat{\mu}_X)^2.$$

In this way:

$$\bar{\sigma}_X^2 = \frac{N}{N-1} \cdot \frac{1}{N} \sum_{i=1}^N (X_i - \hat{\mu}_X)^2 = \frac{N}{N-1} \hat{\sigma}_X^2.$$

and therefore

$$\mathbb{E}(\bar{\sigma}_X^2) \tag{1}$$

$$= \mathbb{E}\left(\frac{N}{N-1} \hat{\sigma}_X^2\right) \tag{2}$$

$$= \frac{N}{N-1} \mathbb{E}(\hat{\sigma}_X^2) \tag{3}$$

$$= \frac{N}{N-1} \sigma^2 \left(1 - \frac{1}{N}\right) \tag{4}$$

$$= \sigma^2. \tag{5}$$

which is unbiased.

This correction compensates for the additional variability introduced by using $\hat{\mu}_X$ as an estimate of μ .