Lesson 3.6 - How to Find the Weighted Mean


> **Video:** How To Find The Weighted Mean and Weighted Average In Statistics by The Organic Chemistry Tutor **Chapter:** 3 - Central Tendency

Formula

$$\bar{x}_w = \frac{\sum wx}{\sum w} = \frac{w_1x_1 + w_2x_2 + \cdots + w_nx_n}{w_1 + w_2 + \cdots + w_n}$$

- x = the values you want to average.
- w = the **weight** of each value: a frequency, a count of people, a percentage, credit hours, a volume, ...
- If the weights are percentages that add to 100% (or decimals that add to 1), the denominator is 1 and you only need the numerator.

> **How to decide which is x and which is w:** x is the thing you want the *average of*. w is *how much each one counts*.



Picture each value as a weight on a beam. The weighted mean is where the beam balances, so it moves toward the heavier weights: 84.8 instead of 86 for the class averages, and 40% instead of 35% for the antifreeze mixture.

Example 1 - a weighted mean with frequencies as weights

Data (10 values): one 10, one 12, four 16s, three 20s, one 24.

Ordinary mean: $\frac{170}{10} = 17$

Weighted mean, using frequencies as weights:

$$\bar{x}_w = \frac{1(10) + 1(12) + 4(16) + 3(20) + 1(24)}{1 + 1 + 4 + 3 + 1} = \frac{170}{10} = 17$$

The two give the same answer: a mean computed from a frequency table **is** a weighted mean.

Example 2 - combining group averages

In a class of 20: 8 students averaged 86, 7 averaged 74, and 5 averaged 98. What is the class average?

$$\bar{x}_w = \frac{8(86) + 7(74) + 5(98)}{8 + 7 + 5} = \frac{688 + 518 + 490}{20} = \frac{1696}{20} = 84.8$$

> Averaging 86, 74 and 98 directly gives 86, which is wrong because the groups are different sizes.

Example 3 - percentages as weights

At a college, 20% of students average 140 lb, 35% average 160 lb, 30% average 175 lb, and 15% average 195 lb.

$$\bar{x}_w = 0.20(140) + 0.35(160) + 0.30(175) + 0.15(195) = 28 + 56 + 52.5 + 29.25 = 165.75 \text{ lb}$$

(The weights add to 1, so there is nothing to divide by.)

Example 4 - final semester grade

Category	Weight	John	Kelly
Homework	15%	92	100
Quizzes	10%	74	82
Lab	20%	83	95
Tests	25%	76	70
Final exam	30%	88	76

John: $0.15(92) + 0.10(74) + 0.20(83) + 0.25(76) + 0.30(88) = 13.8 + 7.4 + 16.6 + 19 + 26.4 = \mathbf{83.2}$

Kelly: $0.15(100) + 0.10(82) + 0.20(95) + 0.25(70) + 0.30(76) = 15 + 8.2 + 19 + 17.5 + 22.8 = \mathbf{82.5}$

Kelly did better on homework, quizzes and lab, but those are only **45%** of the grade. John did better on tests and the final, which are **55%**, so his weighted grade is slightly higher.

Example 5 - grade point average (GPA)

Points: A = 4, B = 3, C = 2. Weights = credit hours.

Course	Credit hours w	Grade	Points x	$w \cdot x$
Chemistry	3	B	3	9
Physics	3	C	2	6
Lab	1	A	4	4
Calculus	4	A	4	16
English	3	B	3	9
Total	**14**			**44**

$$\text{GPA} = \frac{44}{14} \approx 3.14$$

Example 6 - mixing solutions

Rachel mixes **5 gal of 20%** antifreeze with **10 gal of 50%** antifreeze. Is the mixture closer to 20% or 50%?

- An equal-weight average would be 35%.
- The 50% solution has more volume, so it carries more weight. The answer must be between 35% and 50%.

$$C = \frac{5(20\%) + 10(50\%)}{5 + 10} = \frac{100 + 500}{15} = \frac{600}{15} = 40\%$$

The mixture is **40%**, closer to 50% as predicted. The gallons cancel, leaving a percentage.

Key takeaways

- Weighted mean = $\sum wx / \sum w$.
- Use it whenever the values **don't count equally**: group averages, course grades, GPA, mixtures.
- The weighted mean is pulled toward the values with the **largest weights**.