A Cellulose degradation

Substrate: F=13.39; p=6.0e-05; q=2.4e-04
Inoculum: F=2.71; p=0.0817; q=0.1089
Interaction: F=0.98; p=0.4303; q=0.5419

B Hemicellulose degradation

Substrate: F=13.43; p=5.8e-05; q=2.4e-04
Inoculum: F=2.15; p=0.1333; q=0.1599
Interaction: F=0.97; p=0.4378; q=0.5419

C Pectin degradation

Substrate: F=6.89; p=0.0036; q=0.0087
Inoculum: F=4.43; p=0.0212; q=0.0423
Interaction: F=1.33; p=0.2831; q=0.5419

D Alpha-glucan/fructan processing

Substrate: F=16.24; p=1.4e-05; q=1.6e-04
Inoculum: F=9.95; p=4.4e-04; q=0.0026
Interaction: F=1.08; p=0.3826; q=0.5419

E Oligo-/disaccharide utilization

Substrate: F=6.22; p=0.0057; q=0.0115
Inoculum: F=6.77; p=0.0039; q=0.0157
Interaction: F=0.95; p=0.4515; q=0.5419

F Cutin degradation

Substrate: F=3.36; p=0.0489; q=0.0652
Inoculum: F=5.09; p=0.0128; q=0.0385
Interaction: F=1.07; p=0.3880; q=0.5419

G Photoreactivation and NER

Substrate: F=8.62; p=0.0013; q=0.0040
Inoculum: F=19.65; p=6.0e-06; q=7.2e-05
Interaction: F=4.77; p=0.0049; q=0.0588

H DNA repair and damage tolerance

Substrate: F=2.29; p=0.1179; q=0.1287
Inoculum: F=1.48; p=0.2425; q=0.2645
Interaction: F=0.23; p=0.9193; q=0.9193

I Enzymatic ROS defense

Substrate: F=3.50; p=0.0438; q=0.0652
Inoculum: F=3.61; p=0.0401; q=0.0602
Interaction: F=2.22; p=0.0922; q=0.2767

J Carotenoid biosynthesis

Substrate: F=2.85; p=0.0728; q=0.0874
Inoculum: F=1.16; p=0.3259; q=0.3259
Interaction: F=2.54; p=0.0593; q=0.2372

K Sensory photoreception

Substrate: F=0.68; p=0.5127; q=0.5127
Inoculum: F=4.49; p=0.0191; q=0.0423
Interaction: F=0.60; p=0.6626; q=0.7228

L Aromatic-compound catabolism

Substrate: F=3.98; p=0.0299; q=0.0513
Inoculum: F=3.89; p=0.0322; q=0.0552
Interaction: F=3.38; p=0.0221; q=0.1327

Relative abundance (%)

Full sunlight | Reduced UV | Reduced UV and blue-green

Microcosm substrate: Full sunlight, Reduced UV, Reduced UV and blue-green

Inoculum origin: Full sunlight, Reduced UV, Reduced UV and blue-green

Faint points: samples; large points: estimated inoculum-substrate means with pointwise 95% CIs.
q: BH adjustment across 12 groups separately for each model term.