

POLITECNICO DI TORINO

Corso di Laurea Magistrale
in Ingegneria Matematica

Tesi di Laurea Magistrale

Some recent results on the norm of localization operators



Relatori

Prof. Fabio Nicola
firma dei relatori

.....

Candidato

Federico Riccardi
firma del candidato

.....

Anno Accademico 2022-2023

Indice

1	Introduction	3
2	Basics of functional analysis	4
3	Short-Time Fourier Transform	5
3.1	STFT	5
3.1.1	Properties of STFT	5
3.2	Fock Space and Bargmann Transform	5
3.3	Faber-Krahn Inequality for the STFT	5
4	Localization Operators	6
4.1	Definition and properties	6
4.2	Eigenvalues and eigenfunctions	6
5	Recent results from Nicola-Tilli	7
5.1	Case $q = +\infty$	7
5.2	Generic case	7

Sommario

Capitolo 1

Introduction

Capitolo 2

Basics of functional analysis

Capitolo 3

Short-Time Fourier Transform

3.1 STFT

3.1.1 Properties of STFT

3.2 Fock Space and Bargmann Transform

3.3 Faber-Krahn Inequality for the STFT

Theorem from [nicolatilli_fk]

Theorem 3.1. *For every $f \in L^2(\mathbb{R}^d)$ such that $\|f\|_{L^2} = 1$ and every measurable subset $\Omega \subset \mathbb{R}^{2d}$ with finite measure we have*

$$\int_{\Omega} |\mathcal{V}f(x, \omega)|^2 dx d\omega \leq G(|\Omega|)$$

where $G(s)$ is given by

$$G(s) := \int_0^s e^{(-d! \tau)^{1/d}} d\tau \tag{3.1}$$

Capitolo 4

Localization Operators

4.1 Definition and properties

4.2 Eigenvalues and eigenfunctions

Capitolo 5

Recent results from Nicola-Tilli

5.1 Case $q = +\infty$

5.2 Generic case

Let's now consider the case where both p and q are neither 1 or $+\infty$. The result presented in [nicolatilli_norm] include the case ...

$$\|L_F\|_{L_2 \rightarrow L_2} \leq \min\{\kappa_p^{d\kappa_p} A, \kappa_q^{d\kappa_q} B\}$$

Suppose that the minimum is given by $\kappa_p^{d\kappa_p} A$, therefore

$$\kappa_p^{d\kappa_p} A \leq \kappa_q^{d\kappa_q} B \iff \frac{B}{A} \geq \left(\frac{\kappa_p^{\kappa_p}}{\kappa_q^{\kappa_q}}\right)^d$$

We can check if the solution of the problem with just the L^p bound solves also the problem with both bounds, that is $F\|_{L^q} \leq B$, where F is given by ...

$$\|F\|_{L^q}^q = \int_{\mathbb{R}^{2d}} |F(z)|^q dz = \dots = \lambda^q \left(\frac{p-1}{q}\right)^d$$

Since we want F to satisfy the L^q constraint we should have

$$\frac{B}{A} \geq \kappa_p^{d(\frac{1}{q}-\frac{1}{p})} \left(\frac{p}{q}\right)^{\frac{d}{q}}$$

It would be nice if this bound was less restrictive than the first one. Unfortunately that's not the case, in fact it's always true that

$$\left(\frac{p'}{q'}\right)^{\frac{1}{q'}} \left(\frac{p}{q}\right)^{\frac{1}{q}} \geq 1$$

Following the path in `[nicolatilli_norm]` we obtain ...

$$G'(u(t)) = \lambda_1 t^{p-1} + \lambda_2 t^{q-1} \implies u(t) = \frac{1}{d!} \left[-\log(\lambda_1 t^{p-1} + \lambda_2 t^{q-1}) \right]^d, \quad t \in (0, M)$$

Our main goal now is to show that multipliers λ_1, λ_2 are unique and both positive. The easiest fact to prove is that both multipliers are not 0. In fact if one, say λ_2 , was 0, we would obtain that the solution of our problem is the same as the one with just the L^p bound. But we already know that this function does not satisfy the L^q constraint hence it is impossible that $\lambda_2 = 0$.

Suppose now that one of the multipliers, say always λ_2 , is negative. Consider an interval $[a, b] \subset (0, M)$ and a variation $\eta \in L^\infty(0, M)$ supported in $[a, b]$. Thanks to the Gram-Schmidt process we can construct a variation orthogonal to t^{p-1} . Since η is arbitrary we can suppose that it is not orthogonal to t^{q-1} , in particular we can suppose that $\int_a^b t^{q-1} \eta(t) dt < 0$. Therefore the directional derivative of G along η is:

$$\begin{aligned} \int_a^b G'(u(t)) \eta(t) dt &= \int_a^b (\lambda_1 t^{p-1} + \lambda_2 t^{q-1}) \eta(t) dt = \\ &= \lambda_2 \int_a^b t^{q-1} \eta(t) dt > 0 \end{aligned}$$

which contradicts the fact that u is a maximizer.

PROOF OF CONTINUITY

Lastly we shall prove that multipliers λ_1, λ_2 , and hence maximizer, are unique. For this proof it is convenient to express u in a slightly different way

$$u(t) = \frac{1}{d!} \left[\text{Log}_- \left((c_1 t)^{p-1} + (c_2 t)^{q-1} \right) \right]^d$$

To emphasize that u is parametrized by c_1, c_2 we may write $u(t; c_1, c_2)$. Now we define

$$f(c_1, c_2) = p \int_0^T t^{p-1} u(t; c_1, c_2) dt, \quad g(c_1, c_2) = q \int_0^T t^{q-1} u(t; c_1, c_2) dt$$

We want to highlight that, even if not explicit, also T depends on c_1 and c_2 . Nevertheless these functions are differentiable since both T and u are differentiable with respect to (c_1, c_2) , functions $t^{p-1}u$ and $t^{q-1}u$ and their derivatives are bounded in $(0, T)$. Our maximizer u satisfies the constraints only if $f(c_1, c_2) = A^p$, $g(c_1, c_2) = B^q$. Therefore to prove uniqueness of the maximizer we need to show that level sets $\{f = A^p\}$ and $\{g = B^q\}$ intersect in only a point.

First of all we are studying endpoints. For example, if $c_2 = 0$:

$$\begin{aligned} f(c_1, 0) &= p \int_0^{1/c_1} t^{p-1} \frac{1}{d!} \left[-\log(c_1 t)^{p-1} \right]^d dt \stackrel{\tau = c_1 t}{=} \\ &= \frac{p(p-1)^d}{c_1^p d!} \int_0^1 \tau^{p-1} [-\log(\tau)]^d d\tau = \frac{\kappa_p^d}{c_1^p} = A^p \implies c_{1,f} = \frac{\kappa_p^{d/p}}{A} \end{aligned}$$

The same can be done for g and setting $c_1 = 0$ thus we obtain four points

$$c_{1,f} = \frac{\kappa_p^{d/p}}{A}, \quad c_{1,g} = \left(\frac{p-1}{q} \right)^{d/q} \frac{1}{B}, \quad c_{2,f} = \left(\frac{q-1}{p} \right)^{d/p} \frac{1}{A}, \quad c_{2,g} = \frac{\kappa_q^{d/q}}{B}$$

In the regime we are considering one has that $c_{1,f} < c_{1,g}$ and $c_{2,f} > c_{2,g}$, indeed

$$\begin{aligned} c_{1,f} < c_{1,g} &\iff \frac{\kappa_p^{d/p}}{A} < \left(\frac{p-1}{q}\right)^{d/q} \frac{1}{B} \iff \frac{B}{A} < \kappa_p^{d(\frac{1}{q}-\frac{1}{p})} \left(\frac{p}{q}\right)^{d/q} \\ c_{2,f} > c_{2,g} &\iff \left(\frac{q-1}{p}\right)^{d/p} \frac{1}{A} > \frac{\kappa_q^{d/q}}{B} \iff \frac{B}{A} > \kappa_q^{d(\frac{1}{p}-\frac{1}{q})} \left(\frac{q}{p}\right)^{d/p} \end{aligned}$$

Since there is this dispositions of these points we expect there is an intersection between the level sets. Firstly we notice that, for every $c_1 \in (0, c_{1,f})$ there exist a unique value of c_2 for which $f(c_1, c_2) = A^p$. Indeed, from previous computations we notice that $f(c_1, 0)$ is a decreasing function hence $f(c_1, 0) > A^p$, while $\lim_{c_2 \rightarrow +\infty} f(c_1, c_2) = 0$. The uniqueness of this value follows from strict monotonicity of $f(c_1, \cdot)$, in fact:

$$\frac{\partial f}{\partial c_1}(c_1, c_2) = -\frac{p(p-1)}{(d-1)!} c_1^{p-2} \int_0^T \frac{t^{2(p-1)}}{(c_1 t)^{p-1} + (c_2 t)^{q-1}} \left[-\log((c_1 t)^{p-1} + (c_2 t)^{q-1}) \right]^{d-1} dt \quad (5.1)$$

is always strictly negative. We point out that the term $\frac{\partial T}{\partial c_1}(c_1, c_2)u(T; c_1, c_2)$ is zero because u is 0 in T . The same is true for g , therefore on the interval $(0, c_{1,f})$ the level sets of f and g can be seen as the graph of two functions φ, γ . Since $\frac{\partial f}{\partial c_2}, \frac{\partial g}{\partial c_2} < 0$ for every (c_1, c_2) from the implicit function theorem we have that φ and γ are differentiable with respect to c_1 .

Possibile disegno??

If we manage to prove that $(\varphi - \gamma)' < 0$ when $\{f = A^p\}$ and $\{g = B^q\}$ intersect we have uniqueness of the intersection point. Indeed, since $c_{2,f} = \varphi(0) > \gamma(0) = c_{2,g}$, if there were more intersection point for at least one of them it has to be $(\varphi - \gamma)' \geq 0$. Still from the implicit function theorem we have

$$\begin{aligned} \frac{d}{dc_1}(\varphi - \gamma)(c_1) &= -\frac{\frac{\partial f}{\partial c_1}(c_1, \varphi(c_1))}{\frac{\partial f}{\partial c_2}(c_1, \varphi(c_1))} + \frac{\frac{\partial g}{\partial c_1}(c_1, \gamma(c_1))}{\frac{\partial g}{\partial c_2}(c_1, \gamma(c_1))} < 0 \iff \\ \frac{\partial f}{\partial c_1}(c_1, \varphi(c_1)) \frac{\partial g}{\partial c_2}(c_1, \gamma(c_1)) - \frac{\partial f}{\partial c_2}(c_1, \varphi(c_1)) \frac{\partial g}{\partial c_1}(c_1, \gamma(c_1)) &> 0 \end{aligned}$$

As for (5.1) the other derivatives are computed. To simplify the notation we define $h(t; c_1, c_2) = \frac{1}{(d-1)!} \frac{1}{(c_1 t)^{p-1} + (c_2 t)^{q-1}} \left[-\log((c_1 t)^{p-1} + (c_2 t)^{q-1}) \right]^{d-1}$. From Fubini's theorem we can write the product of the integrals as a double integral

$$\begin{aligned} p(p-1)q(q-1)c_1^{p-2}\gamma(c_1)^{q-2} \iint_{[0,T]^2} h(t; c_1, \varphi(c_2))h(s; c_1, \gamma(c_2))t^{2(p-1)}s^{2(q-1)}dtds - \\ p(q-1)q(p-1)c_1^{p-2}\varphi(c_1)^{q-2} \iint_{[0,T]^2} h(t; c_1, \varphi(c_2))h(s; c_1, \gamma(c_2))t^{p+q-2}s^{p+q-2}dtds \end{aligned}$$

At the intersection point $\varphi(c_1) = \gamma(c_1)$ hence the sign of the previous expression depends

only on the sign of

$$\begin{aligned} & \iint_{[0,T]^2} h(t; c_1, \varphi(c_1)) h(s; c_1, \gamma(c_1)) \left(t^{2(p-1)} s^{2(q-1)} - t^{p+q-2} s^{p+q-2} \right) dt ds = \\ & = \iint_{[0,T]^2} h(t; c_1, \varphi(c_1)) h(s; c_1, \gamma(c_1)) t^{p-2} s^{q-2} (t^p s^q - t^q s^p) dt ds \end{aligned}$$

In order to simplify the notation once again we set $H(t, s; c_1) = h(t; c_1, \varphi(c_1)) h(s; c_1, \gamma(c_1))$. Let $T_1 = [0, T]^2 \cap \{t > s\}$ and $T_2 = [0, T]^2 \cap \{t < s\}$. We can split the above integral in two parts

$$\iint_{T_1} H(t, s; c_1) t^{p-2} s^{q-2} (t^p s^q - t^q s^p) dt ds + \iint_{T_2} H(t, s; c_1) t^{p-2} s^{q-2} (t^p s^q - t^q s^p) dt ds$$

We can exchange t with s in the second integral. With this change of variables the domain of integration becomes T_1 and since H is symmetric in t and s we have that the previous quantity is equal to

$$\begin{aligned} & \iint_{T_1} H(t, s; c_1) \left(t^{p-2} s^{q-2} - t^{q-2} s^{p-2} \right) (t^p s^q - t^q s^p) dt ds = \\ & \iint_{T_1} H(t, s; c_1) \frac{1}{t^2 s^2} (t^p s^q - t^q s^p)^2 dt ds \end{aligned}$$

which is strictly positive.