Supplementary Material for *Data-driven regional traffic simulation from heterogeneous mobility sources*

Federico Bellisardi, Chiara Mizzi, Alessandro Fabbri and Armando Bazzani

This document gives further detail on the data layers, the simulation model, the route-choice variants and the detector correction described in the main manuscript. It also reports additional results and lists the commands that reproduce them.

S1 Data layers

Table S1: Data layers and their role in the reconstruction.

Layer	Content	Role	Period
Mobile-phone OD (TIM)	hourly device movements between census areas, aggregated to municipalities and provinces	demand	13 January–23 February 2020
Road graph	3,925 directed links, 2,548 nodes, turn restrictions, derived from OpenStreetMap	network	–
OSM road classes	motorway to unclassified, nearest ways within 30 m	route-choice parameters	extract of October 2026
MTS loop detectors	15-min counts per station and direction	demand scale and demand settings (half of the detectors of February 2020); local correction; volume comparison	Feb 2020, Jun–Jul 2021
Viasat GPS, link counts	hourly probes per matched link, Forlì–Cesena	profile comparison	10–17/02/2020, 28/06–04/07/2021
Viasat GPS, raw traces	private-vehicle fixes in Forlì–Cesena	speeds and class usage for route choice	same periods

Demand is generated in five classes, after removing the cells with the code 000000000. Agents have an observed source and destination between the 50 most active municipalities (23% of the agents on a Monday) or between them and the provinces of neighbouring regions (5%), an observed source and a random destination (20%), a random source and an observed destination (20%), or a random source and destination (31%). This last class represents the activity not attributed to the selected relations, including trips within municipalities outside the 50. Agents with a random end therefore account for 72% of the

1.99 million agents generated on a Monday, after the province factors. The hour convention of the OD tables is discussed in Section S7.

We obtained the MTS counts of 2021 from the monthly open-data files of Regione Emilia-Romagna (dataset DS-1944, CC BY), summing all vehicle classes. The time labels of these files are in local time; converted to UTC, they reproduce the counts of the February 2020 archive used for 2020 (ratio of totals 0.995 on 12 February 2020).

S2 Demand scale

The configuration generator inherited from the operational platform rescaled individual source–destination pairs with ratios between observed and simulated counts at detectors. Because these ratios used the counts of the simulated day itself, the simulation was not independent of the detectors. We switched this rescaling off and instead multiply the demand by a factor k for each day type (main text, Methods). Earlier scans and a fit by demand class, made on a previous version of the engine and demand, are not reported.

The same generator computed the relations with neighbouring provinces from all OD cells, while the classes with random ends were computed from the same cells without removing them, so those trips were generated twice: 63,532 agents on a Monday (4.4%) and 68,388 on a Saturday (5.0%), before scaling. We corrected this and repeated all simulations and selections, with the decision to report the corrected model fixed before the runs. All results in this paper use the corrected model. On the previous version, the selected radius and departure shift were the same, k was 1.4 and 1.2, and the volume ratio, Spearman correlation and dispersion on the detectors of 2021 were 1.10, 0.66 and 0.20, as in the corrected model.

The grid of k was evaluated on the calibration detectors at the 25-km radius, without province factors, which leave k unchanged (Fig. S1). The selected values, 1.5 on weekdays and 1.3 on weekends, are the minima of the loss and give volume ratios of 1.07 and 1.10. The minima are flat: the neighbours cannot be told apart on the loss ($k = 1.4$ $[-0.002, +0.004]$ and $k = 1.6$ $[-0.001, +0.003]$ on weekdays; $k = 1.2$ $[-0.003, +0.003]$ and $k = 1.4$ $[-0.003, +0.004]$ on weekends). The selected value is the best in 52% of the bootstrap resamples on weekdays and 20% on weekends, where $k = 1.2$ (44%) and $k = 1.4$ (33%) are as often or more often the best. The intervals exclude $k \leq 1.1$ on weekdays and $k \leq 1.0$ and $k \geq 1.6$ on weekends.




Figure S1: **Choice of the demand scale k.** Calibration detectors, 25-km radius, without province factors; 83 detectors on weekdays (Monday run) and 95 on weekends (Saturday run). **(a)** Mean squared $\log_{10}$(simulated/observed) of daily volumes; large open markers: selected k. **(b)** Ratio of total simulated to total observed volume. **(c)** Difference in loss from the selected k, with 95% paired bootstrap intervals over detectors (2,000 resamples, seed 20261047).

Table S2 lists the changes tested on the calibration detectors, each against the previous configuration. Each rule was fixed before its runs and required the 95% interval to lie below zero. The first two rows and the trips within the 50 municipalities were structural choices tested on an earlier configuration, with the code 000000000 still in the data, and were not reopened; the others come from the final configuration.

Table S2: Changes tested on the calibration detectors (the first two rows and the intra-municipal row: earlier configuration). Difference in the standard deviation of $\log_{10}$(simulated/observed), new minus reference; for the departure shift, difference in the pooled hourly error (pp). Intervals: 95% paired bootstrap over detectors.

Change	Difference	Interval	Rule met	Kept
OD destination drawn around the declared one (correction of a defect)	+0.0047	[−0.0037, +0.0123]	no	yes
Origins spread around the source point	−0.0187	[−0.0421, +0.0059]	no	yes
Departures 45 min earlier, Monday (pp)	−4.05	[−4.99, −3.17]	yes	yes
Departures 45 min earlier, Saturday (pp)	+0.01	[−0.87, +0.96]	no	–
Intra-municipal trips added	+0.0105	[−0.0011, +0.0221]	no	no
k by day type at 20 km, against common $k = 1.5$	−0.0029	[−0.0061, +0.0002]	no	yes
Radius 25 km against 20 km	−0.0006	[−0.0096, +0.0086]	no	yes

The destination change corrects a verified defect, so we kept it regardless of the result. The origin spreading has no demonstrated effect. The factors by day type bring the level of both day types close to one without measurably changing the dispersion. The radius and k follow the minimum of the loss, so the 25-km radius is the selected value of that rule, without a demonstrated gain over 20 km.

Province factors. The factors were set on the calibration half in two rounds. Dispersion and Spearman correlation are 0.222 and 0.66 without them, 0.193 and 0.72 after the first round, and 0.185 and 0.74 after the second. The factors are 1.74 (Piacenza), 1.42 (Parma), 0.51 (Reggio Emilia), 1.19 (Modena), 0.90 (Bologna), 1.48 (Ferrara), 0.82 (Ravenna), 0.70 (Forlì–Cesena) and 0.55 (Rimini). The evaluation uses the rule stated in the main text, which is met on the test half and in 2021; the dispersion there includes a detector without simulated traffic.

S3 Simulation model and route-choice variants

Agents move at the speed of their link, v_i, which is reduced linearly with density ρ_i below the critical density and set to a critical speed above it. Link speeds are read in km/h from the cartography and assigned by road class from GPS (Table S3). Agents with a random end have a trip-time budget drawn uniformly between 900 and 5,600 s and choose each next link from the transition weights, so a known end sets where they start and does not steer them, while source–destination agents follow the best-path matrix of their type. When the turn-restricted search finds no feasible next node for the cost of their type, these agents use the shortest-distance path matrix. The simulator is deterministic (fixed random seed).

Table S4 lists the variants simulated for 10 February 2020. They were run with the demand of the earlier generator, before the demand scale was revised; the selection uses only GPS counts in Forlì–Cesena. Speeds alone (second to fourth rows) do not change the distribution of traffic over road classes, because most agents follow transition probabilities rather than best paths. Transition weights by road class (last three rows) raise the share of main roads and the agreement with GPS counts in Forlì–Cesena, and the selected variant combines the usage cost of Eq. (1) with these weights. Table S5 compares shortest-distance and GPS-informed route choice over the full campaigns with the scaled demand, and Table S6 shows that route choice hardly changes the hourly profiles.

The variants of Table S4 correspond, in Eq. (1), to shortest distance $(\alpha, \alpha_s) = (1, 0)$, travel time $(0, 1)$, mixed $(0.5, 0.5)$ and usage cost $(0.5, 0)$. The final configuration uses $(0.5, 0.25)$ for all agents.

The cost weights were set in a further step. We simulated a grid of (α, α_s) with all else fixed and compared it with $(0.5, 0)$ on the 100 calibration detectors, for 10 and 15 February 2020 (Table S7). Under the rule fixed before the results were seen, a variant replaces the reference if the 95% paired bootstrap interval of the change in dispersion lies below zero. $(0.5, 0.25)$ met it, in one of seven variants and without correction for multiplicity; on held-out detectors the interval reached zero in all four comparisons. We

Table S3: Road-class properties from private-vehicle GPS traces in Forlì–Cesena, February 2020. Median off-peak speed of links with at least 50 matched points (interquartile range in brackets) and median daily probes per covered link, both applied to all links of each class in the region.

Road class	Links (speed)	Speed (km/h)	Links (usage)	Probes per day
Motorway	44	123 [120–124]	61	130
Trunk	75	83 [76–95]	76	79
Primary	151	55 [46–64]	160	109
Secondary	245	51 [46–60]	248	36
Tertiary	193	53 [43–59]	227	21
Unclassified	42	49 [36–54]	57	14

applied the rule and repeated all later steps with this cost. The earlier selection on GPS is repeated on the final model in Section S8.

Table S4: Route-choice variants simulated for 10 February 2020. Calibration: Spearman correlation between simulated daily link volumes and GPS counts on Forlì–Cesena links. Detectors: Spearman correlation with the daily volumes of the detectors outside Forlì–Cesena complete on that day (not used for selection). Main roads: share of simulated vehicle-kilometres on motorway, trunk and primary links.

Variant	GPS links	Calibration ρ_s	Detectors	Detector ρ_s	Main roads
Shortest distance, default speeds	814	0.474	53	0.198	0.39
Shortest distance, GPS speeds	814	0.478	53	0.190	0.38
Travel time, GPS speeds	814	0.487	53	0.167	0.38
Mixed distance/time, GPS speeds	814	0.484	53	0.175	0.37
Travel time + class transition weights	814	0.537	53	0.109	0.47
Travel time + damped class weights	814	0.532	53	0.105	0.42
Usage cost + class transition weights (selected)	814	0.542	53	0.176	0.48

Table S5: Daily volumes of the simulation before detector correction at the detectors outside Forlì–Cesena: Spearman correlation with the observed volumes for shortest-distance and GPS-informed route choice, paired bootstrap difference with 95% interval (5,000 resamples of detectors), and median absolute $\log_{10}$ ratio of simulated to observed volume.

Campaign	Detectors	ρ_s distance	ρ_s GPS	Difference	$\lvert\log_{10}\rvert$ distance	$\lvert\log_{10}\rvert$ GPS
2020	136	0.64	0.72	0.08 [0.02, 0.15]	0.14	0.13
2021	128	0.54	0.64	0.10 [0.03, 0.18]	0.15	0.14

S4 Detector correction

The correction starts from the increment of Eq. (2) on each detector link and hour. The simulated count on a link is the number of distinct agents present on it during the hour, while the detector counts passages at a section; the two coincide for agents that traverse the whole link within the hour. Barrier counts do not enter the correction. Two pairs of detectors (404 and 680; 503 and 685) lie on the same link, and their observations are averaged before the increment is computed. Downstream, the increment of a link is passed to each admissible outgoing link, multiplied by that link's share of the simulated outgoing counts at the node; upstream, the same rule applies to incoming links. Each branch carries its own increment and its own path length. The spreading skips other detector links, which receive their own increment, and stops when the increment is below one vehicle, after three links along the path, at a forbidden turn or

Table S6: Median hourly-share MAE (pp) of the simulation before detector correction on GPS links of Forlì–Cesena, with shortest-distance and GPS-informed route choice.

Campaign	Route choice	Links	MAE (pp)
2020	shortest distance	442	1.06
2020	GPS-informed	442	1.08
2021	shortest distance	439	1.20
2021	GPS-informed	439	1.21

Table S7: Grid of cost weights on the 100 calibration detectors, 10 and 15 February 2020, without the code 000000000 and before the final choice of cost. Dispersion and Spearman correlation of daily volumes over detector-days; 95% paired bootstrap interval over detectors of the change in dispersion from the reference.

α	α_s	Dispersion	Spearman	Change in dispersion
0.50	0.00	0.189	0.736	reference
1.00	0.00	0.187	0.738	$[-0.011, +0.008]$
0.75	0.25	0.187	0.739	$[-0.009, +0.007]$
0.50	0.50	0.186	0.741	$[-0.010, +0.005]$
0.25	0.75	0.184	0.739	$[-0.013, +0.003]$
0.00	1.00	0.183	0.741	$[-0.017, +0.007]$
0.25	0.25	0.187	0.736	$[-0.007, +0.002]$
0.50	0.25	0.184	0.746	$[-0.008, -0.001]$

at a closed link. Every directed link of the raw simulation is kept in the output, and the hours of the observed table are aligned with the local-time hours of the simulation. In the leave-one-detector-out and depth experiments, all detectors on the left-out link are removed together, so that a co-located detector cannot reproduce the left-out count.

Table S8 summarizes the leave-one-detector-out test and Table S9 the dependence on the depth of the spreading.

Table S8: Leave-one-detector-out test. Errors are $\sum |\tilde{u} - c| / \sum c$ in percent of the observed volume, pooled over detectors or as the median over detectors; reached: share of left-out detectors whose link is modified by the spreading from other detectors; improved/worse: share with lower/higher error than the simulation before correction; ρ_s: Spearman correlation of daily volumes with the observed ones before correction and with the detector left-out.

Year	Detectors	Detector-days	Pooled raw	Pooled LOO	Median raw	Median LOO	Reached (%)	Improved (%)	Worse (%)	ρ_s raw	ρ_s LOO
2020	152	614	42.2	39.8	35.9	33.3	86	60	26	0.72	0.72
2021	141	572	47.4	41.4	39.4	34.8	85	62	23	0.66	0.69

S5 Additional results

Table S10 reports the profile error by number of probes per link. The error decreases with the number of probes for both the reconstruction and the mean detector profile, as expected if part of the difference is sampling noise in the probe counts. Matched links with zero simulated or GPS total are excluded, and link-hour records absent from the GPS extract are counted as zero.

The ratio between probes ψ_i and simulated vehicles ϕ_i on link i,

$$\eta_i = \frac{\sum_{d,h} \psi_i(d,h)}{\sum_{d,h} \phi_i(d,h)}, \tag{1}$$

describes how GPS samples traffic (Fig. S2). Figure S3 shows the daily volumes at left-out detectors.

Table S9: Reduction of the pooled error (percentage points of observed volume) on left-out detectors, 5-fold cross-validation over the 152 detectors of 2020 complete on each day, with bootstrap 95% intervals over detectors; difference: paired with the selected depth 3, the smallest within one standard error (1.5 pp) of the best depth. Raw error: 42.2%.

Depth (links)	Error reduction (pp)	95% interval	Difference (pp)	95% interval
1	0.9	[-0.0, 2.2]	-1.4	[-3.3, +0.2]
2	1.2	[-0.3, 2.9]	-1.1	[-2.2, -0.3]
3	2.3	[0.4, 4.7]	–	–
4	2.5	[0.1, 5.1]	+0.1	[-0.6, +0.8]
5	3.0	[0.5, 5.9]	+0.7	[-0.2, +1.5]
6	3.2	[0.5, 6.2]	+0.8	[-0.2, +1.8]

Table S10: Hourly-share MAE (pp) on GPS links of Forlì–Cesena, by number of GPS probes per link and campaign: reconstruction, mean detector profile and hourly correlation of the reconstruction (medians).

Campaign	Probes	Links	Reconstruction	Mean detector	Correlation
2020	1-49	56	4.49	4.49	0.46
2020	50-199	79	1.86	1.82	0.75
2020	200+	307	0.85	0.81	0.93
2021	1-49	64	3.86	3.89	0.35
2021	50-199	110	1.81	1.81	0.72
2021	200+	265	0.99	0.86	0.89




Figure S2: **GPS coverage. (a,b)** Usable GPS-matched links of Forlì–Cesena (number in parentheses) over the regional network, 10–17 February 2020 and 28 June–4 July 2021. **(c)** Distribution of the link ratio η_i between probe counts and simulated vehicles, Eq. (1). Road network © OpenStreetMap contributors.




Figure S3: **Daily volumes at left-out detectors.** Observed daily volume of each detector (mean over complete days, both directions) against the reconstruction on its link when the detector is left-out and all others are used, **(a)** 10–17 February 2020 and **(b)** 28 June–4 July 2021. Detectors reached by the spreading from other detectors are distinguished from those that are not; ρ_s, Spearman correlation. Dashed line: equality; dotted lines: factor of two.




Figure S4: **Timing on weekdays and weekends.** Error of the hourly shares on the GPS-matched links of Forlì–Cesena, as in Fig. 5, for weekdays and weekends separately: reconstruction and mean detector profile. Each curve shows the fraction of links whose error is below the value on the horizontal axis. Weekend curves lie to the right, so errors are larger on weekends. Curves continue beyond the 6 pp limit of the axis.

Table S11: Separating the OD relations from the province factors with existing simulations. Differences in Spearman correlation and dispersion of daily volumes, with 95% paired bootstrap intervals over detectors (2,000 resamples, seed 20261049). The last block repeats the comparison that also removes the regional pattern of the factors, with a new bootstrap, so its intervals can differ slightly from those of Section S5. Exploratory: added after the results. Dispersion excludes detectors with no simulated traffic.

Comparison	Detectors	Day	n	Spearman	Dispersion
Relations, no factors	Test half, 2020	Monday	82	+0.031 [-0.024, +0.085]	+0.032 [+0.014, +0.051]
Relations, no factors	Test half, 2020	Saturday	93	+0.036 [-0.015, +0.093]	+0.051 [+0.021, +0.080]
Relations, no factors	2021	Monday	76	+0.064 [-0.030, +0.170]	+0.041 [+0.013, +0.070]
Relations, no factors	2021	Saturday	120	+0.044 [-0.010, +0.103]	+0.037 [+0.017, +0.057]
Factors, with relations	Test half, 2020	Monday	82	+0.050 [-0.004, +0.115]	-0.024 [-0.041, -0.007]
Factors, with relations	Test half, 2020	Saturday	93	+0.021 [-0.027, +0.075]	-0.021 [-0.037, -0.007]
Factors, with relations	2021	Monday	76	+0.033 [-0.059, +0.122]	-0.031 [-0.048, -0.014]
Factors, with relations	2021	Saturday	120	+0.003 [-0.041, +0.049]	-0.027 [-0.042, -0.014]
Relations and factors	Test half, 2020	Monday	82	+0.080 [+0.015, +0.147]	+0.008 [-0.011, +0.027]
Relations and factors	Test half, 2020	Saturday	93	+0.058 [+0.001, +0.119]	+0.029 [-0.001, +0.059]
Relations and factors	2021	Monday	76	+0.102 [+0.010, +0.206]	+0.009 [-0.013, +0.031]
Relations and factors	2021	Saturday	120	+0.052 [-0.003, +0.115]	+0.010 [-0.007, +0.027]

Further numbers behind the Results of the main text. *Timing on GPS links of Forlì–Cesena.* All days, 2020 and 2021: raw minus mean detector profile +0.08 [0.06, 0.09] and +0.06 [0.04, 0.08] pp; corrected minus mean detector profile +0.01 [0.00, 0.03] and +0.02 [0.01, 0.04] pp; raw minus OD profile −0.06 [−0.07, −0.05] and −0.06 [−0.08, −0.04] pp, with the simulation better on weekdays (−0.10 and −0.10 pp) and worse on weekends (+0.04 and +0.04 pp). The correction lowers the error by a median of 0.01 pp in 2020 and does not change it in 2021. Because links are spatially correlated, these intervals are optimistic. *Correction.* The reduction at left-out detectors is 2.1 [−0.6, 4.8] and 3.1 [1.5, 4.9] pp on weekdays and weekends in 2020 and 5.9 [1.7, 11.2] and 6.2 [3.7, 9.2] pp in 2021, on different sets of detectors. Half of the detectors have a corrected error between 23% and 49% in 2020 and between 26% and 55% in 2021. Cross-validation over 2020 gives 0.9, 1.2, 2.3, 2.5, 3.0 and 3.2 pp at depths 1 to 6; the best depth is the edge of the grid. Detector 349 recorded 19,599 vehicles on 10 February 2020 and the reconstruction with that detector left out gives 20,219 (hourly correlation 0.99). *OD relations.* Removing the relations from the final model, which also removes the regional pattern of the factors, gives Spearman differences of 0.08 [0.02, 0.15] and 0.06 [0.00, 0.12] on the Monday and Saturday of the test half and 0.10 [0.01, 0.20] and 0.05 [−0.00, 0.11] in 2021, and dispersion differences of +0.008 [−0.012, 0.027], +0.029 [−0.000, 0.059], +0.009 [−0.013, 0.031] and +0.010 [−0.007, 0.028]. Spearman correlations with GPS link counts of Forlì–Cesena are 0.69–0.73 with and without relations, with all intervals including zero. *Single roads.* The day-to-day deviation of a detector's count from its own mean has a median of 0.017 in $\log_{10}$ units, against a dispersion of 0.167 on the Monday and 0.201 on the Saturday of the calibration half. Without the province factors, province explains 38% of the variance of the error on the Monday, road class 9% and observed volume 3% (48% together), and detectors less than 5 km apart have correlated errors (0.69, 20 pairs). With the factors, these shares are 9%, 12% and 6% (26% together), and the correlation is 0.29 below 5 km and −0.02 at 10–20 km. The doubtful detectors are two on closed links, one 870 m from its link and two on a road with a different reference; the median distance from the link is 11 m, and on the 193 evaluated detectors not flagged, dispersion remains 0.255 (without province factors, all days).

S6 Final configuration and reproducibility

The code repository contains the simulator and all analysis scripts, and it explains how to build the simulator and set up the Python environment. The script `hpc/run_all.sh` runs the simulations and analyses of the final configuration in order; the protocols written before the experiments are in `model/results/experiments`. The results are reproduced in the following order:

1. check the hour convention of the OD tables;

Table S12: Final configuration of the model. Every value is fixed before the final simulations.

Setting	Value
Demand	mean OD of 13 January–23 February 2020 by day of the week, without the code 000000000 and without trips within the 50 municipalities; 50 municipalities; relations with neighbouring provinces counted once
Scale k	1.5 on weekdays, 1.3 on Saturdays and Sundays
Province factors	as in Section S2, two rounds
Random ends and departures	radius 25 km; origins spread around the source; departures 45 min earlier than the OD label
Route choice	$\alpha = 0.5$, $\alpha_s = 0.25$, GPS class speeds, usage and transition weights by class
Network	3,925 links, 122 closed, unchanged
Local correction	depth 3, without travel time
Detector split	seed 20261028; 100 calibration, 97 test

2. build the road geometry, the road classes and the GPS link speeds;
3. simulate and select the route-choice variants;
4. build the 2021 detector table from the open-data files;
5. build the demand, run the tests of Table S2, choose k and the province factors, and run the simulations of the fifteen validation days;
6. run the correction, the GPS comparison and the leave-one-out test;
7. run the volume tests, the comparison of route choices, the depth selection, the predictors without simulation, the tests of the OD relations and the final tests;
8. produce the figures and tables.

Steps 1–7 require the proprietary inputs (mobile-phone OD tables, Viasat link counts and traces), which are not distributed. A simulated day takes 10–30 minutes on an 8-core workstation, depending on the demand factor and on the number of days run in parallel. In a single measured run of the final Monday on a 4-core desktop computer (AMD Ryzen 3 3200G), the simulation took 7.1 minutes of wall-clock time and 2.5 GB of memory and reproduced the link flows of the final run byte for byte.

S7 Further details of data and methods

Road classes. Each link was assigned the OpenStreetMap class of the ways nearest to three points along it (within 30 m; Geofabrik extract of October 2026). This gives 709 km of motorways, 326 km of trunk roads, 2,629 km of primary, 4,673 km of secondary and 2,863 km of tertiary roads.

Clock of the OD tables. The hourly OD labels refer to local time and to the start of the hour. On the days of 10–14 February 2020, the weekday OD profile correlates best with the mean profile of the 90 detectors complete on those days at zero lag (0.96, against 0.94 with the OD profile shifted one hour earlier and 0.74 one hour later), although the mean absolute difference does not separate zero lag from one hour earlier (0.60 and 0.60 pp).

Detector data. Time stamps are in UTC and are converted to local time; the 2021 open-data release, in local time, reproduces the archive for February 2020. A detector-day is used when both directions are complete for all 96 intervals. The archive contains no zero counts, and more than 95% of the isolated missing intervals fall before 06:00, which suggests that empty intervals are recorded as missing. The

filter therefore favours busy roads (median daily volume of about 17,000 vehicles, against about 8,000 for detector-days with isolated gaps in 2020) and retains fewer detectors on weekdays, whose nights are quieter. Reading isolated gaps (at most 20 intervals, none longer than 4) as zero extends 2020 from 152 to 227 detectors and hardly changes the raw error (39.6% to 41.1%, both directions together) or the volume ratio (0.98 to 1.03). 2021 behaves likewise (44.5% to 44.5%, ratio 1.11 to 1.11). We did not repeat the leave-one-out test on the extended set.

GPS matching. GPS fixes with engine on and speed of at least 5 km/h were matched to the nearest link within 20 m whose bearing agrees with the heading within 30°. Off-peak speeds are for weekdays, 10:00–16:00. The fallback to the shortest-distance path described in Section S3 is rare: 51 next-node decisions in one of the eight simulations, none in the others.

Splits, profiles and bootstrap. The detector split uses seed 20261028. In the comparison of hourly profiles we exclude links with zero simulated or GPS total and keep only links whose geometry is identical in the graph used for map matching; the mean detector profile is the mean normalized profile of the detector directions complete on all days of the subset. Comparisons between route choices use 2,000 paired bootstrap resamples of detectors (seed 20261029; 5,000 in Table S5, on all complete detectors). Intervals of the predictors without simulation use 2,000 resamples (seed 20261041). The depth of the correction is selected with seed 20261033, and the effect of the correction on GPS links is summarized by the minimum number of links between a link and any detector link.

More OD relations. On an earlier configuration, without province factors and with the code 000000000 in the data, $N = 100$ raised the Spearman correlation on the calibration detectors from 0.669 to 0.726 (difference interval $[0.017, 0.104]$), but dispersion went from 0.215 to 0.232 ($[-0.003, 0.035]$), so it did not beat the model. The result is hard to interpret, because the share of agents with a random end falls as N grows, which does not separate the information in the relations from the amount of random traffic. Section S8 repeats the test on the final model.

Value of the OD relations, further results. Table S11 separates the relations from the province factors using runs without the regional pattern; its last block also removes that pattern, and Section S8 gives the complete design. Without factors, the Spearman differences are 0.03–0.07 with intervals that include zero, and dispersion is larger with relations, with all intervals above zero.

Direction of travel. For each detector we computed the correlation over the 24 hours between the observed and the simulated difference of the two directions. The test is exploratory and changes no setting. Its rule required, on the test half, that the 95% interval of the difference in median correlation between $N = 100$ and the model lie above zero; it is not met on the final model (Section S8). An error in the mapping between detector and link directions, which we could not check independently, would produce the same weak signal. Comparisons of volumes in this paper therefore sum the two directions.

S8 Experiments on the final model

These experiments followed protocols written before their runs, in the code repository. They change no setting of the model and are exploratory, because the data had been seen. A run with the same engine and configuration reproduced the final Monday simulation byte for byte.

How much of the OD table is kept. On a Monday the table has 3.15 million trips per day. 17.8% have the code 000000000 at one or both ends and are removed from the model; the meaning of this code is not documented in the data we received. 43.1% are within a municipality: 38.0% within the 50 municipalities, left out, and 5.1% within other municipalities, kept in the class with two random ends, of which they are more than a third. The remaining 39.2% are between municipalities with known ends, on 7,711 pairs: 28% of these trips keep their relation (476 pairs with agents after rounding) and 72% have a known relation that is discarded. The relations kept are only 6% of the pairs, because trips concentrate on few pairs. They are short: mean 13 km, and 87% lie below 20 km on a Monday. Saturday is similar.

Relations and province factors. Figure S5 shows the complete design. Table S20 gives the differences. The interaction in dispersion is negative in all four comparisons ($[-0.022, -0.000]$, $[-0.031, -0.008]$, $[-0.024, -0.004]$, $[-0.025, -0.006]$), so the factors reduce the spread more with relations. In the simulation without relations the regional pattern acts only through random ends.




Figure S5: **OD relations and province factors, 2×2 design.** Dispersion (top) and Spearman correlation (bottom) of daily volumes in the four simulations. Parallel lines mean no interaction. Dispersion excludes detectors with no simulated traffic.

OD pairings. Table S13 gives the results. Origins, destinations and the total are identical to the real arm in every interval, and null D keeps the distance classes exactly. Null M raises the mean trip to 66–68 km at 50 municipalities (real 13 km) and 84–86 km at 347 (real 12 km). Null D moves 51% of agents at 50 municipalities and 67% at 347; more swaps add little, because the constraints leave many pairs no alternative. The six runs of null M at 347 municipalities (about 88,000 sources) ended with a segmentation fault in an earlier configuration, and these runs were not repeated. The probable cause, not confirmed for these runs, is a defect in a diagnostic output of the simulator that reads an uninitialised value for sources without a location; other runs with many sources failed there, and we corrected this for the tests in the paragraph on routing and trip length below. The 347 arm has a larger dispersion than the 50 arm (0.33–0.47 against 0.19–0.26), so it is used only for this comparison: nearly all its demand is assigned to specific pairs, and only 5% of its agents on a Monday have a random end, against 72% in the model, so some detectors away from those pairs have little traffic. The twin test takes a real run as the truth and asks whether a real run with another seed beats the nulls; it has no model error, so it overstates the power. Spearman correlations with GPS link counts of Forlì–Cesena differ from the real arm by at most 0.008 at 50 municipalities and 0.003 at 347.

Route choice on the final model. The selected form stands: no other has an interval above zero on both days (Table S14). No other cost weight replaces $(0.5, 0.25)$ on the calibration detectors (Table S16). Table S15 separates speeds and usage weights at detectors.

Scale k with province factors. The selected values remain the minima of the grid (ratios 1.01 and 1.04). The neighbours cannot be told apart on the loss: $k = 1.4$ $[-0.001, +0.004]$ and $k = 1.6$ $[-0.001, +0.003]$ on weekdays, $k = 1.2$ $[-0.002, +0.004]$ and $k = 1.4$ $[-0.002, +0.004]$ on weekends.

How many detectors. With 200 random subsets of the calibration half, 25 detectors choose the selected k or a neighbour in 72% of cases on weekdays and 80% on weekends, and 50 detectors in 96% and 99%; the mean loss on the test half changes by at most 0.001 between 25 and 50 detectors. Province

Table S13: OD pairings: real against null M and D, at 50 or 347 municipalities. Differences: real minus mean of three rearrangements; 95% paired bootstrap over detectors (2,000 resamples, seed 20261061). Sens.: median $|\log_{10}|$(real/null) of daily volume at detectors. Twin: difference in dispersion with a real run as the truth.

Null	Detectors	Day	n	Dispersion: real, null, difference			Spearman: real, null, difference			Sens.	Twin, disp.
M, 50	Test half, 2020	Monday	82	0.217	0.241	[-0.049, -0.000]	0.670	0.654	[-0.025, +0.061]	0.022	[-0.112, -0.050]
M, 50	Test half, 2020	Saturday	93	0.264	0.281	[-0.040, -0.000]	0.663	0.658	[-0.030, +0.038]	0.021	[-0.103, -0.046]
M, 50	2021	Monday	76	0.186	0.225	[-0.069, -0.011]	0.640	0.684	[-0.128, +0.032]	0.043	[-0.154, -0.085]
M, 50	2021	Saturday	120	0.214	0.234	[-0.040, -0.002]	0.638	0.666	[-0.072, +0.012]	0.022	[-0.124, -0.065]
D, 50	Test half, 2020	Monday	82	0.217	0.218	[-0.002, +0.000]	0.670	0.668	[-0.004, +0.012]	0.005	[-0.003, +0.001]
D, 50	Test half, 2020	Saturday	93	0.264	0.266	[-0.006, +0.000]	0.663	0.661	[-0.009, +0.012]	0.007	[-0.004, +0.001]
D, 50	2021	Monday	76	0.186	0.186	[-0.002, +0.002]	0.640	0.654	[-0.040, +0.003]	0.005	[-0.005, -0.001]
D, 50	2021	Saturday	120	0.214	0.214	[-0.002, +0.002]	0.638	0.642	[-0.016, +0.005]	0.006	[-0.004, -0.001]
D, 347	Test half, 2020	Monday	82	0.428	0.427	[-0.007, +0.010]	0.652	0.641	[-0.003, +0.030]	0.014	[-0.044, -0.014]
D, 347	Test half, 2020	Saturday	93	0.469	0.475	[-0.012, -0.000]	0.643	0.643	[-0.017, +0.016]	0.013	[-0.036, -0.008]
D, 347	2021	Monday	76	0.327	0.325	[-0.006, +0.010]	0.689	0.703	[-0.039, +0.008]	0.011	[-0.022, -0.004]
D, 347	2021	Saturday	120	0.414	0.418	[-0.011, +0.003]	0.656	0.656	[-0.018, +0.017]	0.012	[-0.024, -0.012]

Table S14: Route-choice variants on the final model: Spearman correlation with GPS counts in Forlì–Cesena, and 95% interval of the difference from the selected variant, paired over links.

Variant	Monday	minus selected	Saturday	minus selected
Shortest distance, default speeds	0.650	[-0.074, +0.002]	0.654	[-0.088, -0.022]
Shortest distance, GPS speeds	0.663	[-0.062, +0.012]	0.661	[-0.081, -0.015]
Travel time, GPS speeds	0.668	[-0.044, +0.009]	0.682	[-0.050, -0.005]
Mixed distance/time, GPS speeds	0.678	[-0.040, +0.024]	0.685	[-0.052, +0.005]
Travel time + class weights	0.690	[-0.010, +0.015]	0.715	[-0.005, +0.016]
Travel time + damped class weights	0.680	[-0.024, +0.010]	0.702	[-0.022, +0.007]
Usage cost + class weights, default speeds	0.676	[-0.016, -0.005]	0.701	[-0.014, -0.004]
Usage cost + class weights (selected)	0.687	–	0.710	–

Table S15: Speeds and usage weights at detectors outside Forlì–Cesena, final model: Spearman correlation and 95% paired intervals. V0: shortest distance, default speeds. V1: shortest distance, GPS speeds. D1: usage cost and class weights, default speeds. V6: usage cost and class weights, GPS speeds.

Detectors	Day	n	V0	V1	D1	V6	V1−V0	D1−V0	V6−D1
Test half, 2020	Monday	73	0.60	0.64	0.66	0.68	[+0.01, +0.07]	[-0.04, +0.16]	[-0.01, +0.05]
Test half, 2020	Saturday	84	0.56	0.58	0.66	0.68	[+0.00, +0.04]	[+0.01, +0.19]	[-0.01, +0.05]
2021	Monday	69	0.62	0.63	0.63	0.65	[-0.01, +0.04]	[-0.10, +0.13]	[-0.01, +0.05]
2021	Saturday	109	0.48	0.50	0.62	0.61	[-0.01, +0.04]	[+0.03, +0.24]	[-0.03, +0.02]

Table S16: Cost weights on the calibration detectors, final model (selected: dispersion 0.185, Spearman 0.736). Weights: 100α and $100\alpha_s$. Interval of the change in dispersion from the selected weights, 95% paired.

Weights	Dispersion	Spearman	Change in dispersion
w100, s000	0.190	0.729	[-0.004, +0.013]
w075, s025	0.188	0.735	[-0.003, +0.008]
w050, s050	0.188	0.734	[-0.002, +0.008]
w025, s075	0.184	0.735	[-0.007, +0.006]
w000, s100	0.184	0.734	[-0.013, +0.012]
w025, s025	0.189	0.729	[-0.002, +0.010]
w050, s000	0.190	0.726	[+0.001, +0.009]

factors from 25 calibration detectors give a larger dispersion than those from 100 in 9 of 12 comparisons over three subsets, and those from 50 in 5 of 12. The values for a single province vary strongly between subsets: Modena from 0.50 to 1.75 and Piacenza from 1.44 to 2.50, against 1.19 and 1.74 with all 100 detectors. Table S17 gives the correction with subsets of detectors. It uses a fixed 20% of detectors held out on single days, so it is not comparable in size with the leave-one-out test.

Table S17: Correction with subsets of detectors, depth 3, five repeats, on a fixed 20% of complete detectors held out. Hourly error of the held-out detectors.

Day	Share (%)	Detectors	Raw (%)	Corrected (%)	Gain (pp)	Range over repeats
2020-02-10	25	46	46.5	45.1	+1.4	[-0.1, +3.7]
2020-02-10	50	91	46.5	44.0	+2.5	[+0.3, +3.6]
2020-02-10	75	136	46.5	43.6	+2.9	[+0.6, +4.0]
2020-02-10	100	182	46.5	43.4	+3.1	[+2.9, +3.7]
2021-06-28	25	41	42.6	42.7	-0.1	[-1.6, +1.0]
2021-06-28	50	82	42.6	41.8	+0.8	[-1.4, +3.7]
2021-06-28	75	123	42.6	41.5	+1.1	[-1.5, +3.8]
2021-06-28	100	164	42.6	41.3	+1.4	[-1.3, +4.3]

More relations with province factors. $N = 100$, with a multiplier of 1.15 on both day types, does not beat the model on the calibration half: dispersion 0.200 against 0.185 ($[-0.002, 0.030]$), Spearman 0.781 against 0.736 ($[0.004, 0.096]$); the rule requires lower dispersion. The 2021 detectors were not used. The direction of travel does not meet its rule on the test half (median signal 0.046 against 0.054, $[-0.051, 0.048]$).

2021 without calibration detectors. The 141 detectors of 2021 include 60 of the calibration half. All detectors, and in brackets the 81 others: volume ratio 1.10 (1.11, interval [0.99, 1.24]), Spearman correlation 0.66 (0.65, [0.50, 0.76]), median $|\log_{10}|$ ratio 0.142 (0.146, [0.124, 0.200]), within a factor of two 84% (81%, [73, 90]).

Variability between runs. Between four seeds, the ranges are at most 0.002 for the volume ratio, 0.002 for the Spearman correlation, 0.006 for dispersion and 0.011 for the median error. These ranges may understate the variation between runs, because choices made while agents move use a random generator that does not change with the seed.

What explains the error at single roads. This diagnostic followed a protocol in the code repository, written before it was computed, and uses no simulations. The outcome is $\log_{10}$(simulated/observed) of

daily volume and its absolute value, raw simulation. Table S18 gives the out-of-fold R^2 lost when a group of predictors is removed from an ordinary least squares model (5 folds, 100 repeats), with permutation p values Holm-adjusted within outcome. By its rule, a group is relevant if on the Monday it loses at least 0.05 with adjusted $p < 0.05$, and at least 0.02 in 2021. Only road class meets it, on the signed error: primary roads are overestimated (median +0.051, 94 detectors) and secondary roads underestimated (−0.036, 53), while trunk roads have an unstable value (−0.237 with 12 detectors, −0.044 with 6 in 2021). This diagnostic uses the 164 detectors with weekday data on the Monday, whereas Fig. 7 of the main text uses the 152 with complete days of the leave-one-out test. The full model has an out-of-fold R^2 of 0.09 on the Monday, 0.12 on the Saturday and −0.10 in 2021, and −0.33 on the Monday when one province is left out in turn; a random forest gives 0.04 out of bag on the Monday. No group predicts the absolute error under the rule. The measure of omitted roads counts their length within 1 km, and does not show whether a given road takes traffic from the detector's road. 2021 is not a replication, since detectors and demand are the same.

Table S18: Out-of-fold R^2 lost when a group of predictors is removed, signed error, with Holm-adjusted permutation p. Omitted: OpenStreetMap roads farther than 30 m from any model link, within 1 km. Parallel: model links within 1 km with a similar bearing. Activity: population and OpenStreetMap points of activity within 1 and 5 km.

Group	Monday (164)	p	2021 (122)	p
Road class	0.194	0.007	0.090	0.042
Province	0.067	0.066	−0.055	0.78
Length and centrality	−0.020	1.00	−0.013	0.78
Parallel roads	−0.025	1.00	−0.023	0.78
Omitted roads	−0.007	0.70	−0.059	0.78
Activity	0.007	1.00	−0.008	0.48
Distance from the regional boundary	−0.013	1.00	−0.007	0.78

Routing and trip length. On a Monday, trips between municipalities average 12.0 km in a straight line between centres (median 10.4 km, 96.5% below 25 km). The simulation gives about 40 km per agent: 42–45 km for agents with a random end and 27 km for agents with a known relation, estimated from hourly positions. These further tests followed a protocol in the code repository, written before their runs, and are exploratory. They used the detectors of the calibration half and half of the test half, by province, in 2020 and 2021; the other half was set aside and not used, and the model was not changed. With 347 municipalities, the share of detectors more than a factor of two below, after removing the level of each arm, is 17.9% on average in 2020 and 6.5% in the model. Four route types per relation give 16.8%, and a probability of 0.10 or 0.30 of leaving the best path at each node gives 17.2% and 11.9%, so no arm reaches the threshold of 10%. Four route types raise the Spearman correlation in 2021 to 0.70, against 0.66 with one route and 0.62 in the model. With the model of 50 municipalities, shorter walks of 900–1,800 s and k selected again at 2.5 on both day types, with province factors estimated again, give a dispersion of 0.272 on the test detectors against 0.222 in the model ($[0.017, 0.091]$ and $[0.021, 0.073]$ on the two days), and a Spearman correlation of 0.73 against 0.72, and of 0.66 against 0.62 in 2021, with intervals that include zero. Walks of exactly 900 s, with or without a radius of 10 km for known relations, and with the factors of the model, give dispersions of 0.28–0.30 and Spearman correlations of 0.75–0.76. By the rule fixed before the runs, these results do not replace the model. The tests with four route types and those on trip length used a simulator with the diagnostic output corrected; on the final Monday it reproduces the link flows byte for byte.

Route weights by link. Following a protocol written before the runs, 398 links of Forlì–Cesena with at least 50 probes in February 2020 received a weight from their own GPS count, on the scale of the class weights, with everything else unchanged, for one Monday and one Saturday. The simulation moves towards GPS on those links (Spearman from 0.66 to 0.77, in sample). At the detectors of the province, which did

not enter the weights, the Spearman differences have intervals that include zero, and the dispersion falls on the Monday of February 2020 (0.228 to 0.176, $[-0.091, -0.016]$, 18 detectors) but not on the Saturday (0.314 to 0.326, $[-0.054, +0.059]$, 19) or in 2021 (Monday 0.177 to 0.190, $[-0.032, +0.053]$, 7; Saturday 0.224 to 0.200, $[-0.072, +0.035]$, 11). The rule is not met, and the model keeps the class weights. With so few detectors the test is inconclusive.

S9 Value of the OD relations

To measure the value of the OD relations, we ran one simulation per day type without them. We kept the number of agents per interval and k, but drew all origins and destinations from population. The province factors act through sources with a location, so this simulation loses them too. Under a rule fixed before the runs, the relations are supported if, on the test half of 2020 or on the detectors of 2021, the 95% interval of the Spearman difference is above zero or that of the dispersion difference is below zero. We also compared with GPS link counts, and we repeated the runs with further seeds to measure the variability between them. We also tried more relations: on the calibration detectors only, we let the relations among the 100 most active municipalities keep both ends, with a new multiplier of k (Section S8).

Further experiments followed protocols written before their runs (Section S8). They are exploratory, because the data had already been seen, and they do not change the model. To separate relations from factors, we ran the simulation without relations with node scores multiplied by the square root of the province factor, which gives a trip the geometric mean of the factors of its ends, and we completed a 2×2 design. To isolate the pairings, we rearranged the destinations of agents with located relations within each 15-min interval, three times per case, at 50 and 347 municipalities. Origins, destinations and the total are unchanged. In null M the rearrangement is random, while in null D each agent stays in the distance class of its real trip (0–5, 5–10, 10–20, 20–50, above 50 km). Real pairs are supported if they beat null D in at least two of four comparisons on the same metric.

We now ask what the OD relations themselves add. Only part of the table can answer: on a Monday, 28% of the trips between municipalities with known ends keep their relation, 11% of the table (Section S8). Removing the relations from the final model takes away the regional pattern of the factors as well. That comparison raises the Spearman correlation on the test half (0.08 [0.02, 0.15] on the Monday and 0.06 [0.00, 0.12] on the Saturday) and on the Monday of 2021 (0.10 [0.01, 0.20]), which meets the rule fixed before the runs, while the Saturday interval of 2021 reaches zero and dispersion does not improve (Section S5). When the same pattern is kept in the simulation without relations, the Spearman differences stay positive with intervals that include zero, and the model with relations has a larger spread of errors in all four comparisons, with intervals that exclude zero in three (Fig. S6, top; Table S20). The factors reduce the spread more when relations are present (Section S8). Chance between runs is small compared with these differences.

Removing relations also changes how many trips start and end in each place. To isolate the pairings, we kept these numbers exact in each 15-min interval and rearranged only who goes where. Random pairs make the trips far too long, and they raise dispersion, with intervals that exclude zero in all four comparisons, two of them narrowly (Fig. S6, middle); agreement in ranking does not change measurably. When distances are kept, real and rearranged pairs can hardly be told apart (Fig. S6, bottom). At 50 municipalities no dispersion interval excludes zero and at 347 one does, so the rule fixed before the runs (two of four) is not met. Agreement with GPS link counts does not change either. These tests have limited power: rearranging with distances kept changes the volume at the median detector by only 1–3%, and a test with a simulated run as the truth shows that the expected effect at 50 municipalities is about as small as the intervals (Section S8). Extending the relations to 100 municipalities with province factors does not beat the model under its rule (Section S8).

The simulation without relations keeps the agents per interval, the daily rhythm and the population, so it is not a random traffic pattern. In an analysis made after the results above, without detector 153, the larger spread with relations comes mainly from the low tail in 2020: 10–14% of detectors are more than a factor of two below with relations, against 5% without. With a spread measured from the quartiles no interval excludes zero, and the Spearman correlation is higher with relations in all four comparisons, with intervals that include zero. The median error differs in one comparison of four, in favour of the simulation without relations. So these detectors do not show that random ends are better than relations.

Null D with its five distance classes can still change trip length within a class, and it could not be read as a test of pairings alone. We therefore registered a further test, null D2, with a measured power rule: each radius is classified only if the power of the rule reaches 80%, and otherwise it is inconclusive whatever the interval. The protocol asked for distance bins of 2 km. These failed its check on the share of agents moved, so before any D2 run an amendment set them to 4 km; the width was chosen knowing that share (51–52% of agents moved, mean distance within 0.3%), and it clears the threshold narrowly. Each arm uses two real runs and three rearrangements at 347 municipalities, and pools 368 rows on 168 detectors. The estimand is the slope b of the observed log difference from the null on the simulated log difference between real and null, estimated with a cross-fitted instrumental variable, with $b = 1$ expected if real pairs act as simulated and $b = 0$ if they add nothing. Table S19 gives the results. Both radii are inconclusive, with a tau/sigma ratio of 0.01 and 0.06 against the 0.3 needed for 80% power. At 25 km the estimate has no meaning, because the signal is near zero. At 10 km, the primary radius of the protocol, tested only here, the interval excludes both 0 and 1 and the volume is 40–45% too low, with k not retuned; under the simulated power analysis an interval excludes zero in 6% of cases if $b = 0$ and in 14% if $b = 1$. So this estimate cannot be read as the value of the pairings. The arm also differs from the reported model, as described in Section S8. The value of specific origin–destination pairings could not be determined from link-count observations with the statistical power available in this study.

Table S19: Null D2 test, 347 municipalities, 168 detectors, 368 rows. Interval: 95% bootstrap over detectors. τ/σ: measured signal over residual spread. Power: of the registered rule, at the measured τ/σ. Exploratory, because the data had been seen.

Radius	b	Interval	τ/σ	Power	Classification
25 km, as in the model	not interpretable	$[-233, 195]$	0.011	0.00	inconclusive
10 km	4.58	$[2.39, 7.37]$	0.063	0.11	inconclusive

Winter 2023. Following a protocol written before any simulated value was compared with a 2023 count, we compared the model with mean detector counts of winter 2023, outside the calibration half. This is not a replication, because the simulations are the same and the detectors overlap with the earlier sets, and the dates and holidays of the period are not documented in the data we received. Set A (all detectors outside the calibration half) gives a Spearman correlation of 0.74 [0.65, 0.81] on the Monday (122 detectors) and 0.73 [0.63, 0.80] on the Saturday (140); set B (those also in the 2020 or 2021 sets) gives 0.70 [0.57, 0.79] (89) and 0.67 [0.53, 0.78] (99); set C (the others) gives 0.85 [0.72, 0.91] (33) and 0.85 [0.72, 0.91] (41). Most of the apparent gain comes from which detectors are in the set: on common detectors the year changes little (Monday test half 0.63 in 2020 and 0.63 in 2023; Saturday test half 0.62 and 0.62; Monday 2021 0.64 and 0.66; Saturday 2021 0.64 and 0.69). The means over many days also lower the noise, and set C is not untouched, because earlier rounds scanned k on means of 2022–24. The volume is 39% above the 2023 counts on the Monday and 12% above on the Saturday; we report this as a description only. Null D2 on set A gives the same classification at both radii.

Table S20: OD relations and province factors in a complete 2×2 design: difference from removing the relations with the factors kept in the random ends, and from adding the factors to the simulation without relations. Positive Spearman differences and negative dispersion differences mean a gain from adding the source. Intervals: 95% paired bootstrap over detectors. Exploratory.

		OD relations, with factors		Factors, no relations
Detectors	Day	Spearman	Dispersion	Dispersion
Test half, 2020	Monday	+0.024 [-0.027, +0.077]	+0.021 [+0.002, +0.038]	-0.013 [-0.026, +0.001]
Test half, 2020	Saturday	+0.034 [-0.016, +0.085]	+0.037 [+0.011, +0.063]	-0.008 [-0.021, +0.006]
2021	Monday	+0.068 [-0.011, +0.152]	+0.021 [-0.002, +0.045]	-0.011 [-0.026, +0.003]
2021	Saturday	+0.034 [-0.021, +0.094]	+0.022 [+0.003, +0.040]	-0.012 [-0.024, -0.000]




Figure S6: **What the OD relations add.** Each row compares the model with real relations with a version without them: the marker is the estimated change and the bar its 95% bootstrap interval, so a bar crossing the dashed line means that the data cannot establish a change. Blue circles: test half, 2020; orange squares: all detectors, 2021; Monday and Saturday simulations. **Relations removed, factors kept**: no located relations, with the regional pattern of the factors kept in the random ends. **Null M**: destinations rearranged at random within each 15-min interval, so that origins and destinations by municipality are exact and distances are lost. **Null D**: the same, with each agent kept in the distance class of its real trip. The null rows are the mean of three rearrangements. **(a)** Spread of the error between roads. **(b)** Ranking of roads. Real relations tend to have a lower spread than random pairs, and cannot be told apart from pairs with the same distances.