

UNIVERSITA' DI BOLOGNA

FACOLTA' DI SCIENZE MATEMATICHE FISICHE E NATURALI

CORSO DI LAUREA MAGISTRALE IN SCIENZE INFORMATICHE

Tesi di laurea

---

# Multi $\pi$ calcolo

---


*Candidato:*

Federico VISCOMI

*Tutore*

Prof. Roberto GORRIERI

.....



---

ANNO ACCADEMICO 20011/2012



## 0.1 Abstract

Il  $\pi$  calcolo e' un formalismo che descrive e analizza le proprieta' del calcolo concorrente. Nasce come proseguio del lavoro gia' svolto sul CCS (Calculus of Communicating Systems). L'aspetto appetibile del  $\pi$  calcolo rispetto ai formalismi precedenti e' l'essere in grado di descrivere la computazione concorrente in sistemi la cui configurazione puo' cambiare nel tempo. Nel CCS e nel  $\pi$  calcolo manca la possibilita' di modellare sequenze atomiche di azioni e di modellare la sincronizzazione multiparte. Il Multi CCS [3] estende il CCS con un'operatore di strong prefixing proprio per colmare tale vuoto. In questa tesi si cerca di trasportare per analogia le soluzioni introdotte dal Multi CCS verso il  $\pi$  calcolo. Il risultato finale e' un linguaggio chiamato Multi  $\pi$  calcolo.

In particolare il Multi  $\pi$  calcolo permette la sincronizzazione transazionale e la sincronizzazione multiparte. aggiungere una sintesi brevissima dei risultati ottenuti sul Multi  $\pi$  calcolo.



# Contents

0.1	Abstract . . . . .	3
<b>1</b>	<b>TODO</b>	<b>7</b>
<b>2</b>	<b><math>\Pi</math> calculus</b>	<b>9</b>
2.1	Syntax . . . . .	9
2.2	Operational Semantic(without structural congruence) . . . . .	12
2.2.1	Early operational semantic(without structural congruence) . . . . .	12
2.2.2	Late operational semantic(without structural congruence) . . . . .	13
2.2.3	Distinction between late and early semantics . . . . .	14
2.3	Structural congruence . . . . .	14
2.4	Operational semantic with structural congruence . . . . .	24
2.4.1	Early semantic with $\alpha$ conversion only . . . . .	24
2.4.2	Early semantic with structural congruence . . . . .	25
2.4.3	Late semantic with structural congruence . . . . .	25
2.5	Equivalence of the semantics . . . . .	26
2.5.1	Equivalence of the early semantics . . . . .	26
2.5.2	Equivalence of the late semantics . . . . .	37
2.6	Bisimilarity, congruence and equivalence . . . . .	37
2.6.1	Late bisimilarity . . . . .	37
2.6.2	Early bisimilarity . . . . .	38
2.6.3	Congruence . . . . .	38
2.6.4	Open bisimilarity . . . . .	38
<b>3</b>	<b>Multi <math>\pi</math> calculus with strong output</b>	<b>39</b>
3.1	Syntax . . . . .	39
3.2	Operational semantic . . . . .	39
3.2.1	Early operational semantic with structural congruence . . . . .	39
3.2.2	Low level semantic . . . . .	40
3.2.3	Early operational semantic without structural congruence . . . . .	49
3.3	Strong bisimilarity and equivalence . . . . .	51
3.3.1	Strong bisimilarity . . . . .	51
3.3.2	Open bisimilarity . . . . .	53
<b>4</b>	<b>Multi <math>\pi</math> calculus with strong input</b>	<b>55</b>
4.1	Syntax . . . . .	55
4.2	Operational semantic . . . . .	55
4.2.1	Early operational semantic with structural congruence . . . . .	55
4.2.2	Late operational semantic with structural congruence . . . . .	56
4.2.3	Low level semantic . . . . .	57
4.2.4	Late operational semantic without structural congruence . . . . .	62

<b>5</b>	<b>Multi <math>\pi</math> calculus with strong input and output</b>	<b>65</b>
5.1	Syntax . . . . .	65
5.2	Operational semantic . . . . .	65
5.2.1	Early operational semantic with structural congruence . . . . .	65
5.2.2	Late operational semantic with structural congruence . . . . .	67
5.2.3	Low level semantic . . . . .	70

# Chapter 1

## TODO

- dimostrare(o negare) l'equivalenza del pi calcolo con e senza congruenza strutturale
- nel multi pi calcolo con strong prefixing solo su input o solo su output: definire una semantica di basso livello sulla falsariga di quell'articolo
- fare un quadro generale sulle equivalenze nel pi calcolo
- scegliere una equivalenza(forse la open va bene) per multi pi calcolo che sia una congruenza per input(ma non lo sarà per il parallelo)
- trovare equivalenza che sia una congruenza(es: open step) per tutti gli operatori
- trovare la congruenza coarsest contenuta nella bisimulazione scelta in precedenza





## Chapter 2

# $\Pi$ calculus

The  $\pi$  calculus is a mathematical model of processes whose interconnections change as they interact. The basic computational step is the transfer of a communications link between two processes. The idea that the names of the links belong to the same category as the transferred objects is one of the cornerstone of the calculus. The  $\pi$  calculus allows channel names to be communicated along the channels themselves, and in this way it is able to describe concurrent computations whose network configuration may change during the computation.

A coverage of  $\pi$  calculus is on [4], [5] and [7]

### 2.1 Syntax

We suppose that we have a countable set of names  $\mathbb{N}$ , ranged over by lower case letters  $a, b, \dots, z$ . This names are used for communication channels and values. Furthermore we have a set of identifiers, ranged over by  $A$ . We represent the agents or processes by upper case letters  $P, Q, \dots$ . A process can perform the following actions:

$$\pi ::= \bar{x}y \mid x(z) \mid \tau$$

The process are defined by the following grammar:

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A$$

and they have the following intuitive meaning:

$0$  is the empty process which cannot perform any actions

$\pi.P$  is an action prefixing, this process can perform action  $\pi$  e then behave like  $P$ , the action can be:

$\bar{x}y$  is an output action, this sends the name  $y$  along the name  $x$ . We can think about  $x$  as a channel or a port, and about  $y$  as an output datum sent over the channel

$x(z)$  is an input action, this receives a name along the name  $x$ .  $z$  is a variable which stores the received data.

$\tau$  is a silent or invisible action, this means that a process can evolve to  $P$  without interaction with the environment

for any action which is not a  $\tau$ , the first name that appears in the action is called subject of the action and the second name is called object of the action.

$P + Q$  is the sum, this process can enact either  $P$  or  $Q$

$P|Q$  is the parallel composition,  $P$  and  $Q$  can execute concurrently and also synchronize with each other

---

$B(0, I) = \emptyset$	$B(Q + R, I) = B(Q, I) \cup B(R, I)$
$B(\bar{x}y.Q, I) = B(Q, I)$	$B(Q R, I) = B(Q, I) \cup B(R, I)$
$B(x(y).Q, I) = \{y, \bar{y}\} \cup B(Q, I)$	$B((\nu x)Q, I) = \{x, \bar{x}\} \cup B(Q, I)$
$B(\tau.Q, I) = B(Q, I)$	
$B(A(\tilde{x}), I) = \begin{cases} B(Q, I \cup \{A\}) \text{ where } A(\tilde{x}) \stackrel{\text{def}}{=} Q & \text{if } A \notin I \\ \emptyset & \text{if } A \in I \end{cases}$	

---

Table 2.1: Bound occurrences

---

$fn(\bar{x}y.Q) = \{x, \bar{x}, y, \bar{y}\} \cup fn(Q)$	$fn(Q + R) = fn(Q) \cup fn(R)$	$fn(0) = \emptyset$
$fn(x(y).Q) = \{x, \bar{x}\} \cup (fn(Q) - \{y, \bar{y}\})$	$fn(Q R) = fn(Q) \cup fn(R)$	
$fn((\nu x)Q) = fn(Q) - \{x, \bar{x}\}$	$fn(\tau.Q) = fn(Q)$	$fn(A(\tilde{x})) = \{\tilde{x}\}$

---

Table 2.2: Free occurrences

$(\nu z)P$  is the scope restriction. This process behave as  $P$  but the name  $z$  is local. This process cannot use the name  $z$  to interact with other processes.

$A$  is an identifier. Every identifier has a definition

$$A(x_1, \dots, x_n) = P$$

the  $x_i$ s must be pairwise different. The intuition is that we can substitute for some of the  $x_i$ s in  $P$  to get a  $\pi$  calculus process.

To resolve ambiguity we can use parenthesis and observe the conventions that prefixing and restriction bind more tightly than composition and prefixing binds more tightly than sum.

**Definition 2.1.1.** We say that the input prefix  $x(z).P$  binds  $z$  in  $P$  or is a *binder* for  $z$  in  $P$ . We also say that  $P$  is the *scope* of the binder and that any occurrence of  $z$  in  $P$  are *bound* by the binder. Also the restriction operator  $(\nu z)P$  is a binder for  $z$  in  $P$ .

**Definition 2.1.2.**  $bn(P)$  is the set of names that have a bound occurrence in  $P$  and is defined as  $B(P, \emptyset)$ , where  $B(P, I)$ , with  $I$  a set of identifiers, is defined in table 2.1

**Definition 2.1.3.** We say that a name  $x$  is *free* in  $P$  if  $P$  contains a non bound occurrence of  $x$ . We write  $fn(P)$  for the set of names with a free occurrence in  $P$ .  $fn(P)$  is defined in table 2.2

**Definition 2.1.4.**  $n(P)$  which is the set of all names in  $P$  and is defined in the following way:

$$n(P) = fn(P) \cup bn(P)$$

**Definition 2.1.5.** We say that  $\tau$  and actions which does not have any binder  $xy, \bar{x}y$  are *free* actions. Whether the other actions are *bound* actions.

In a definition

$$A(x_1, \dots, x_n) = P$$

---

$0\{b/a\} = 0$
$(\bar{x}y.Q)\{b/a\} = \bar{x}\{b/a\}y\{b/a\}.Q\{b/a\}$
$(x(y).Q)\{b/a\} = x\{b/a\}(y).Q\{b/a\}$ if $y \neq a$ and $y \neq b$
$(x(a).Q)\{b/a\} = x\{b/a\}(a).Q$
$(x(b).Q)\{b/a\} = x\{b/a\}(c).((Q\{c/b\})\{b/a\})$ where $c \notin n(Q)$
$(\tau.Q)\{b/a\} = \tau.Q\{b/a\}$
if $a \in \{x_1, \dots, x_n\}$ then
$(A(x_1, \dots, x_n \mid y_1, \dots, y_m))\{b/a\} = \begin{cases} A(x_1\{b/a\}, \dots, x_n\{b/a\} \mid y_1, \dots, y_m) & \text{if } b \notin \{y_1, \dots, y_m\} \\ A(x_1\{b/a\}, \dots, x_n\{b/a\} \mid y_1, \dots, y_{i-1}, c, y_{i+1}, \dots, y_m) & \text{if } b = y_i \\ \text{where } c \text{ is fresh} \end{cases}$
if $a \notin \{x_1, \dots, x_n\}$ then
$(A(x_1, \dots, x_n \mid y_1, \dots, y_m))\{b/a\} = A(x_1, \dots, x_n \mid y_1, \dots, y_m)$
$(Q + R)\{b/a\} = Q\{b/a\} + R\{b/a\}$
$(Q R)\{b/a\} = Q\{b/a\} R\{b/a\}$
$((\nu y)Q)\{b/a\} = (\nu y)Q\{b/a\}$ if $y \neq a$ and $y \neq b$
$((\nu a)Q)\{b/a\} = (\nu a)Q$
$((\nu b)Q)\{b/a\} = (\nu c)((Q\{c/b\})\{b/a\})$ where $c \notin n(Q)$ if $a \in fn(Q)$
$((\nu b)Q)\{b/a\} = (\nu b)Q$ if $a \notin fn(Q)$

---

Table 2.3: Syntactic substitution

the  $x_1, \dots, x_n$  are all the free names contained in  $P$ , specifically

$$fn(P) \subseteq \{x_1, \dots, x_n\}$$

If we look at the definitions of  $bn$  and of  $fn$  we notice that if  $P$  contains another identifier whose definition is:

$$B(z_1, \dots, z_h) = Q$$

then we have

$$fn(Q) \subseteq \{x_1, \dots, x_n\}$$

**Definition 2.1.6.**  $P\{b/a\}$  is the syntactic substitution of name  $b$  for a different name  $a$  inside a  $\pi$  calculus process, and it consists in replacing every free occurrences of  $a$  with  $b$ . If  $b$  is a bound name in  $P$ , in order to avoid name capture we perform an appropriate  $\alpha$  conversion.  $P\{b/a\}$  is defined in table 2.3. There is the following short notation

$$\{\tilde{x}/\tilde{y}\} \text{ means } \{x_1/y_1, \dots, x_n/y_n\}$$

---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(y).P \xrightarrow{xz} P\{z/y\}}$
<b>SumR</b> $\frac{Q \xrightarrow{\alpha} Q'}{P + Q \xrightarrow{\alpha} Q'}$	<b>ParR</b> $\frac{Q \xrightarrow{\alpha} Q' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P Q'}$
<b>SumL</b> $\frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'}$	<b>ParL</b> $\frac{P \xrightarrow{\alpha} P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P' Q}$
<b>Res</b> $\frac{P \xrightarrow{\alpha} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha} (\nu z)P'}$	<b>ResAlp</b> $\frac{(\nu w)P\{w/z\} \xrightarrow{xz} P' \quad w \notin n(P)}{(\nu z)P \xrightarrow{xz} P'}$
<b>EComR</b> $\frac{P \xrightarrow{\bar{x}y} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>ClsL</b> $\frac{P \xrightarrow{\bar{x}(z)} P' \quad Q \xrightarrow{xz} Q' \quad z \notin fn(Q)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$
<b>EComL</b> $\frac{P \xrightarrow{xy} P' \quad Q \xrightarrow{\bar{x}y} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>ClsR</b> $\frac{P \xrightarrow{xz} P' \quad Q \xrightarrow{\bar{x}(z)} Q' \quad z \notin fn(P)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$
<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$	<b>Cns</b> $\frac{A(\tilde{x}) \stackrel{def}{=} P \quad P\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha} P'}{A(\tilde{x})\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha} P'}$
<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$	<b>OpnAlp</b> $\frac{(\nu w)P\{w/z\} \xrightarrow{\bar{x}(w)} P' \quad w \notin n(P) \quad x \neq w \neq z.}{(\nu z)P \xrightarrow{\bar{x}(w)} P'}$

---

Table 2.4: Early transition relation without structural congruence

## 2.2 Operational Semantic(without structural congruence)

### 2.2.1 Early operational semantic(without structural congruence)

The semantic of a  $\pi$  calculus process is a labeled transition system such that:

- the nodes are  $\pi$  calculus process. The set of node is  $\mathbb{P}$
- the actions can be:
  - unbound input  $xy$
  - unbound output  $\bar{x}y$
  - the silent action  $\tau$
  - bound output  $\bar{x}(y)$

The set of actions is  $\mathbb{A}$ , we use  $\alpha$  to range over the set of actions.

- the transition relations is  $\rightarrow \subseteq \mathbb{P} \times \mathbb{A} \times \mathbb{P}$

In the following section we present the early semantic without structural congruence and without *alpha* conversion.

**Definition 2.2.1.** The *early transition relation*  $\rightarrow \subseteq \mathbb{P} \times \mathbb{A} \times \mathbb{P}$  is the smallest relation induced by the rules in table 2.4. Where with  $\tilde{x}$  we mean a sequence of names  $x_1, \dots, x_n$ .

**Example** We show now an example of the so called scope extrusion, in particular we prove that

$$a(x).P \mid (\nu b)\bar{a}b.Q \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)$$

where we suppose that  $b \notin fn(P)$ . In this example the scope of  $(\nu b)$  moves from the right hand component to the left hand.

$$\text{CLOSER} \frac{\text{EINP} \frac{}{a(x).P \xrightarrow{ab} P\{b/x\}} \quad \text{OPN} \frac{\text{OUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q} \quad a \neq b}{(\nu b)\bar{a}b.Q \xrightarrow{\bar{a}(b)} Q} \quad b \notin fn((\nu b)\bar{a}b.Q)}{a(x).P \mid (\nu b)\bar{a}b.Q \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)}$$

**Example** We want to prove now that:

$$((\nu b)a(x).P) \mid \bar{a}b.Q \xrightarrow{\tau} ((\nu c)(P\{c/b\}\{b/x\})) \mid Q$$

where  $b \notin bn(P)$

$$\text{RESALP} \frac{\text{RES} \frac{\text{EINP} \frac{}{(a(x).P)\{c/b\} \xrightarrow{ab} P\{c/b\}\{b/x\}} \quad c \notin n(a(b))}{(\nu c)((a(x).P)\{c/b\}) \xrightarrow{ab} (\nu c)(P\{c/b\}\{b/x\})} \quad b \notin n((a(x).P)\{c/b\})}{(\nu b)a(x).P \xrightarrow{ab} (\nu c)P\{c/b\}\{b/x\}}$$

$$\text{EComL} \frac{(\nu b)a(x).P \xrightarrow{ab} (\nu c)P\{c/b\}\{b/x\} \quad \text{EOUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q}}{((\nu b)a(x).P) \mid \bar{a}b.Q \xrightarrow{\tau} ((\nu c)(P\{c/b\}\{b/x\})) \mid Q}$$

**Example** We have to spend some time to deal with the change of bound names in an identifier. Suppose we have

$$A(x) \stackrel{def}{=} \underbrace{x(y).x(a).0}_P$$

From the definition of substitution it follows that

$$A(x)\{y/x\} = A(y)$$

The identifier  $A(y)$  is expected to behave consistently with

$$P\{y/x\} = y(z).y(a).0$$

so we have to prove

$$A(y) \xrightarrow{yw} y(a).0$$

We can prove this in the following way:

$$\text{CNS} \frac{A(x) \stackrel{def}{=} P \quad \text{EINP} \frac{}{P\{y/x\} \xrightarrow{yw} y(a).0}}{A(y) \xrightarrow{yw} y(a).0}$$

### 2.2.2 Late operational semantic(without structural congruence)

In this case the set of actions  $\mathbb{A}$  contains

- bound input  $x(y)$
- unbound output  $\bar{x}y$
- the silent action  $\tau$
- bound output  $\bar{x}(y)$

**Definition 2.2.2.** The *late transition relation without structural congruence*  $\rightarrow_{\subseteq} \mathbb{P} \times \mathbb{A} \times \mathbb{P}$  is the smallest relation induced by the rules in table 2.5. TUTTE LE SEMANTICHE LATE DEL PI CALCOLO SONO DA AGGIORNARE!!!! !!! !! !

---

<b>LInp</b> $\frac{z \notin fn(P)}{x(y).P \xrightarrow{x(z)} P\{z/y\}}$	<b>Res</b> $\frac{P \xrightarrow{\alpha} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha} (\nu z)P'}$
<b>SumL</b> $\frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'}$	<b>SumR</b> $\frac{Q \xrightarrow{\alpha} Q'}{P + Q \xrightarrow{\alpha} Q'}$
<b>ParL</b> $\frac{P \xrightarrow{\alpha} P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P' Q}$	<b>ParR</b> $\frac{Q \xrightarrow{\alpha} Q' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P Q'}$
<b>ComL</b> $\frac{P \xrightarrow{x(y)} P' \quad Q \xrightarrow{\bar{x}(z)} Q'}{P Q \xrightarrow{\tau} P'\{z/y\} Q'}$	<b>ComR</b> $\frac{P \xrightarrow{\bar{x}(z)} P' \quad Q \xrightarrow{x(y)} Q'}{P Q \xrightarrow{\tau} P' Q'\{z/y\}}$
<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$	<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$
<b>ClsL</b> $\frac{P \xrightarrow{\bar{x}(z)} P' \quad Q \xrightarrow{xz} Q' \quad z \notin fn(Q)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$	<b>ClsR</b> $\frac{P \xrightarrow{xz} P' \quad Q \xrightarrow{\bar{x}(z)} Q' \quad z \notin fn(P)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$
<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$	<b>Cns</b> $\frac{A(\tilde{x}) \stackrel{def}{=} P \quad P\{\tilde{y}/\tilde{x}\} \xrightarrow{\alpha} P'}{A(\tilde{y}) \xrightarrow{\alpha} P'}$

---

Table 2.5: Late semantic without structural congruence

### 2.2.3 Distinction between late and early semantics

There are some differences between late and early semantics:

**Communication** da scrivere

**Input** da scrivere

**Parallel composition** the side condition in the rule *Par* for the late semantic is important because:

$$(x(z).P|Q)|\bar{x}y.R \xrightarrow{\tau} (P\{w/z\}|Q)\{y/w\}|R$$

da scrivere

## 2.3 Structural congruence

Structural congruences are a set of equations defining equality and congruence relations on process. They can be used in combination with an SOS semantic for languages. In some cases structural congruences help simplifying the SOS rules: for example they can capture inherent properties of composition operators (e.g. commutativity, associativity and zero element). Also, in process calculi, structural congruences let processes interact even in case they are not adjacent in the syntax. There is a possible trade off between what to include in the structural congruence and what to include in the transition rules: for example in the case of the commutativity of the sum operator. It is worth noticing that in most process calculi every structurally congruent processes should never be distinguished and thus any semantic must assign them the same behaviour.

**Definition 2.3.1.** A *change of bound names* in a process  $P$  is the replacement of a subterm  $x(z).Q$  of  $P$  by  $x(w).Q\{w/z\}$  or the replacement of a subterm  $(\nu z)Q$  of  $P$  by  $(\nu w)Q\{w/z\}$  where in each case  $w$  does not occur in  $Q$ .

**Definition 2.3.2.** A *context*  $C[\cdot]$  is a process with a placeholder. If  $C[\cdot]$  is a context and we replace the placeholder with  $P$ , then we obtain  $C[P]$ . In doing so, we make no  $\alpha$  conversions.

---

$\text{ALPSUM} \frac{P_1 \equiv_\alpha Q_1 \quad P_2 \equiv_\alpha Q_2}{P_1 + P_2 \equiv_\alpha Q_1 + Q_2}$	$\text{ALPTAU} \frac{P \equiv_\alpha Q}{\tau.P \equiv_\alpha \tau.Q}$
$\text{ALPRES1} \frac{P\{y/x\} \equiv_\alpha Q \quad x \neq y \quad y \notin \text{fn}(P)}{(\nu x)P \equiv_\alpha (\nu y)Q}$	$\text{ALPRES} \frac{P \equiv_\alpha Q}{(\nu x)P \equiv_\alpha (\nu x)Q}$
$\text{ALPINP1} \frac{P\{y/x\} \equiv_\alpha Q \quad x \neq y \quad y \notin \text{fn}(P)}{z(x).P \equiv_\alpha z(y).Q}$	$\text{ALPINP} \frac{P \equiv_\alpha Q}{x(y).P \equiv_\alpha x(y).Q}$
$\text{ALPPAR} \frac{P_1 \equiv_\alpha Q_1 \quad P_2 \equiv_\alpha Q_2}{P_1   P_2 \equiv_\alpha Q_1   Q_2}$	$\text{ALPOUT} \frac{P \equiv_\alpha Q}{\bar{x}y.P \equiv_\alpha \bar{x}y.Q}$
$\text{ALPIDE} \frac{}{A(\tilde{x} \tilde{y}) \equiv_\alpha A(\tilde{x} \tilde{y})}$	$\text{ALPZERO} \frac{}{0 \equiv_\alpha 0}$

---

Table 2.6:  $\alpha$  equivalence laws

**Definition 2.3.3.** A *congruence* is a binary relation on processes such that:

- $S$  is an equivalence relation
- $S$  is preserved by substitution in contexts: for each pair of processes  $(P, Q)$  and for each context  $C[\cdot]$

$$(P, Q) \in S \Rightarrow (C[P], C[Q]) \in S$$

**Definition 2.3.4.** Processes  $P$  and  $Q$  are  $\alpha$  convertible or  $\alpha$  equivalent if  $Q$  can be obtained from  $P$  by a finite number of changes of bound names. If  $P$  and  $Q$  are  $\alpha$  equivalent then we write  $P \equiv_\alpha Q$ . Specifically the  $\alpha$  equivalence is the smallest binary relation on processes that satisfies the laws in table 2.6

It remains the problem of proving that  $\alpha$  equivalence is well defined, i.e. if we change only some bound names in a process  $P$  then we get a process  $\alpha$  equivalent to  $P$ .

**Lemma 2.3.1.** Inversion lemma for  $\alpha$  equivalence

- If  $P \equiv_\alpha 0$  then  $P$  is also the null process  $0$
- If  $P \equiv_\alpha \tau.Q_1$  then  $P = \tau.P_1$  for some  $P_1$  such that  $P_1 \equiv_\alpha Q_1$
- If  $P \equiv_\alpha \bar{x}y.Q_1$  then  $P = \bar{x}y.P_1$  for some  $P_1$  such that  $P_1 \equiv_\alpha Q_1$
- If  $P \equiv_\alpha z(y).Q_1$  then one and only one of the following cases holds:
  - $P = z(x).P_1$  for some  $P_1$  such that  $P_1\{y/x\} \equiv_\alpha Q_1$
  - $P = z(y).P_1$  for some  $P_1$  such that  $P_1 \equiv_\alpha Q_1$
- If  $P \equiv_\alpha Q_1 + Q_2$  then  $P = P_1 + P_2$  for some  $P_1$  and  $P_2$  such that  $P_1 \equiv_\alpha Q_1$  and  $P_2 \equiv_\alpha Q_2$ .
- If  $P \equiv_\alpha Q_1 | Q_2$  then  $P = P_1 | P_2$  for some  $P_1$  and  $P_2$  such that  $P_1 \equiv_\alpha Q_1$  and  $P_2 \equiv_\alpha Q_2$ .
- If  $P \equiv_\alpha (\nu y)Q_1$  then one and only one of the following cases holds:
  - $P = (\nu x)P_1$  such that  $P_1\{y/x\} \equiv_\alpha Q_1$
  - $P = (\nu y).P_1$  for some  $P_1$  such that  $P_1 \equiv_\alpha Q_1$
- If  $P \equiv_\alpha A(\tilde{x})$  then  $P$  is  $Q$ .

---

SC-ALP	$\frac{P \equiv_\alpha Q}{P \equiv Q}$	$\alpha$ conversion
abelian monoid laws for sum:		
SC-SUM-ASC	$M_1 + (M_2 + M_3) \equiv (M_1 + M_2) + M_3$	associativity
SC-SUM-COM	$M_1 + M_2 \equiv M_2 + M_1$	commutativity
SC-SUM-INC	$M + 0 \equiv M$	zero element
abelian monoid laws for parallel:		
SC-COM-ASC	$P_1 (P_2 P_3) \equiv (P_1 P_2) P_3$	associativity
SC-COM-COM	$P_1 P_2 \equiv P_2 P_1$	commutativity
SC-COM-INC	$P 0 \equiv P$	zero element
scope extension laws:		
SC-RES	$(\nu z)(\nu w)P \equiv (\nu w)(\nu z)P$	
SC-RES-INC	$(\nu z)0 \equiv 0$	
SC-RES-COM	$(\nu z)(P_1 P_2) \equiv P_1 (\nu z)P_2$ if $z \notin fn(P_1)$	
SC-RES-SUM	$(\nu z)(P_1 + P_2) \equiv P_1 + (\nu z)P_2$ if $z \notin fn(P_1)$	
unfolding law:		
SC-IDE	$A(\tilde{w}) \equiv P\{\tilde{w}/\tilde{x}\}$	if $A(\tilde{x}) \stackrel{def}{=} P$

---

Table 2.7: Structural congruence axioms

*Proof.* This lemma works because given  $Q$  we know which rules must be at the end of any proof tree of  $P \equiv_\alpha Q$ .  $\square$

**Lemma 2.3.2.** Let  $P$  be a process and  $y, w, z$  names such that  $w = z$  or  $w \notin fn(P)$  then  $P\{w/z\}\{y/w\} \equiv_\alpha P$  non ho una dimostrazione ma lo da per scontato in [2] paragrafo 1.3.1

**Definition 2.3.5.** We define a *structural congruence*  $\equiv$  as the smallest congruence on processes that satisfies the axioms in table 2.7

We can make some clarification on the axioms of the structural congruence:

*unfolding* this just helps replace an identifier by its definition, with the appropriate parameter instantiation. The alternative is to use the rule *Cns* in table 2.4.

*$\alpha$  conversion* is the  $\alpha$  conversion, i.e., the choice of bound names, it identifies agents like  $x(y).\bar{z}y$  and  $x(w).\bar{z}w$ . In the semantic of  $\pi$  calculus we can use the structural congruence with the rule SC-ALP or we can embed the  $\alpha$  conversion in the SOS rules. In the early case, the rule for input and the rules *ResAlp*, *OpnAlp*, *Cns* take care of  $\alpha$  conversion, whether in the late case the rule for communication and the rules *ResAlp*, *OpnAlp*, *Cns* are in charge for  $\alpha$  conversion.

*abelian monoidal properties of some operators* We can deal with associativity and commutativity properties of sum and parallel composition by using SOS rules or by axiom of the structural congruence. For example the commutativity of the sum can be expressed by the following two rules:

$$\text{SumL} \frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'} \quad \text{SumR} \frac{Q \xrightarrow{\alpha} Q'}{P + Q \xrightarrow{\alpha} Q'}$$

or by the following rule and axiom:

$$\text{Sum} \frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'} \quad \text{SC-SUM} \quad P + Q \equiv Q + P$$



and the rule *Str*

*scope extension* We can use the scope extension laws in table 2.7 or the rules *Opn* and *Cls* in table 2.4 to deal with the scope extension.

**Lemma 2.3.3.**

$$a \in fn(Q) \Rightarrow fn(Q\{b/a\}) = (fn(Q) - \{a\}) \cup \{b\}$$

*Proof.* □

**Lemma 2.3.4.**  $P \equiv_{\alpha} Q \Rightarrow fn(P) = fn(Q)$

*Proof.* The proof goes by induction on rules

*AlpZero* the lemma holds because  $P$  and  $Q$  are the same process.

*AlpTau* :

$$\begin{array}{ll} P \equiv_{\alpha} Q & \text{rule premise} \\ \Rightarrow fn(P) = fn(Q) & \text{inductive hypothesis} \\ \Rightarrow fn(\tau.P) = fn(\tau.Q) & \text{definition of } fn \end{array}$$

*AlpOut* :

$$\begin{array}{ll} P \equiv_{\alpha} Q & \text{rule premise} \\ \Rightarrow fn(P) = fn(Q) & \text{inductive hypothesis} \\ \Rightarrow fn(P) \cup \{x, y\} = fn(Q) \cup \{x, y\} & \text{definition of } fn \\ \Rightarrow fn(\bar{x}y.P) = fn(\bar{x}y.Q) & \end{array}$$

*AlpRes1* : we consider two cases:

$x \notin fn(P)$  :

$$\begin{array}{ll} P\{y/x\} \equiv_{\alpha} Q \text{ and } y \notin fn(P) & \text{rule premises} \\ \Rightarrow fn(P\{y/x\}) = fn(Q) & \text{inductive hypothesis} \\ \Rightarrow fn(P) = fn(Q) & x \notin fn(P) \text{ and def of substitution} \\ \Rightarrow fn(P) = fn(Q) \text{ and } y \notin fn(Q) & y \notin fn(P) \end{array}$$

Since  $x \notin fn(P)$  then  $fn(P) = fn(P) - \{x\}$ . Since  $y \notin fn(Q)$  then  $fn(Q) = fn(Q) - \{y\}$ . From  $fn(P) = fn(P) - \{x\}$ ,  $fn(Q) = fn(Q) - \{y\}$ ,  $fn(P) = fn(Q)$  and the definition of substitution it follows that  $fn((\nu x)P) = fn((\nu y)Q)$

$x \in fn(P)$  :

$$\begin{array}{ll} P\{y/x\} \equiv_{\alpha} Q & \text{rule premise} \\ \Rightarrow fn(P\{y/x\}) = fn(Q) & \text{inductive hypothesis} \\ \Rightarrow fn(P\{y/x\}) - \{y\} = fn(Q) - \{y\} & \\ \Rightarrow (fn(P) - \{x\} \cup \{y\}) - \{y\} = fn(Q) - \{y\} & \\ \Rightarrow fn(P) - \{x\} = fn(Q) - \{y\} & \text{definition of } fn \\ \Rightarrow fn((\nu x)P) = fn((\nu y)Q) & \end{array}$$

*AlpInp1* : we consider two cases:

$x \notin fn(P)$  :

$$\begin{array}{ll} P\{y/x\} \equiv_{\alpha} Q \text{ and } y \notin fn(P) & \text{rule premises} \\ \Rightarrow fn(P\{y/x\}) = fn(Q) & \text{inductive hypothesis} \\ \Rightarrow fn(P) = fn(Q) & x \notin fn(P) \text{ and def of substitution} \\ \Rightarrow fn(P) = fn(Q) \text{ and } y \notin fn(Q) & y \notin fn(P) \end{array}$$

Since  $x \notin fn(P)$  then  $fn(P) = fn(P) - \{x\}$ . Since  $y \notin fn(Q)$  then  $fn(Q) = fn(Q) - \{y\}$ . From  $fn(P) = fn(P) - \{x\}$ ,  $fn(Q) = fn(Q) - \{y\}$  and  $fn(P) = fn(Q)$  it follows that  $fn(P) - \{x\} = fn(Q) - \{y\}$  and so  $(fn(P) - \{x\}) \cup \{z\} = (fn(Q) - \{y\}) \cup \{z\}$  which gives  $fn(z(x).P) = fn(z(y).Q)$ .

$x \in fn(P) :$

$$\begin{array}{ll}
P\{y/x\} \equiv_{\alpha} Q & \text{rule premise} \\
\Rightarrow fn(P\{y/x\}) = fn(Q) & \text{inductive hypothesis} \\
\Rightarrow fn(P) - \{y\} = fn(Q) - \{y\} & \\
\Rightarrow (fn(P) - \{x\} \cup \{y\}) - \{y\} = fn(Q) - \{y\} & \text{lemma 2.3.3} \\
\Rightarrow fn(P) - \{x\} = fn(Q) - \{y\} & \\
\Rightarrow (fn(P) - \{x\}) \cup \{z\} = (fn(Q) - \{y\}) \cup \{z\} & \text{definition of } fn \\
\Rightarrow fn(z(x).P) = fn(z(y).Q) & 
\end{array}$$

*AlpSum* :

$$\begin{array}{ll}
P_1 \equiv_{\alpha} Q_1 \text{ and } P_2 \equiv_{\alpha} Q_2 & \text{rule premises} \\
\Rightarrow fn(P_1) = fn(Q_1) \text{ and } fn(P_2) = fn(Q_2) & \text{inductive hypothesis} \\
\Rightarrow fn(P_1) \cup fn(P_2) = fn(Q_1) \cap fn(Q_2) & \text{definition of } fn \\
\Rightarrow fn(P_1 + P_2) = fn(Q_1 + Q_2) & 
\end{array}$$

*AlpPar* :

$$\begin{array}{ll}
P_1 \equiv_{\alpha} Q_1 \text{ and } P_2 \equiv_{\alpha} Q_2 & \text{rule premises} \\
\Rightarrow fn(P_1) = fn(Q_1) \text{ and } fn(P_2) = fn(Q_2) & \text{inductive hypothesis} \\
\Rightarrow fn(P_1) \cup fn(P_2) = fn(Q_1) \cap fn(Q_2) & \text{definition of } fn \\
\Rightarrow fn(P_1 | P_2) = fn(Q_1 | Q_2) & 
\end{array}$$

*AlpRes* :

$$\begin{array}{ll}
P \equiv_{\alpha} Q & \text{rule premise} \\
\Rightarrow fn(P) = fn(Q) & \text{inductive hypothesis} \\
\Rightarrow fn(P) - \{x\} = fn(Q) - \{x\} & \text{definition of } fn \\
\Rightarrow fn((\nu x)P) = fn((\nu x)Q) & 
\end{array}$$

*AlpInp* :

$$\begin{array}{ll}
P \equiv_{\alpha} Q\{x/y\} & \text{rule premise} \\
\Rightarrow fn(P) = fn(Q) & \text{inductive hypothesis} \\
\Rightarrow (fn(P) - \{y\}) \cup \{x\} = (fn(Q) - \{y\}) \cup \{x\} & \text{definition of } fn \\
\Rightarrow fn(x(y).P) = fn(x(y).Q) & 
\end{array}$$

*AlpIde* the lemma holds because  $P$  and  $Q$  are the same process.

□

**Lemma 2.3.5.**  $x \notin fn(P) \Rightarrow P\{x/y\}\{b/a\} \equiv_{\alpha} P\{b/a\}\{x/y\}$

**Lemma 2.3.6.**  $\alpha$  equivalence is invariant with respect to substitution. In other words

$$\begin{array}{l}
P \equiv_{\alpha} Q \\
b \notin fn(P) \quad \Rightarrow \quad P\{b/a\} \equiv_{\alpha} Q\{b/a\} \\
b \notin fn(Q)
\end{array}$$

*Proof.* : If  $a$  and  $b$  are the same name then the substitution has no effect and the lemma holds. Otherwise:

$$\begin{aligned}
 P &\equiv_{\alpha} Q && \text{lemma hypothesis} \\
 \Rightarrow fn(P) &= fn(Q) && \text{lemma 2.3.4} \\
 \Rightarrow a \notin fn(P) \wedge a \notin fn(Q) &\text{ or } a \in fn(P) \wedge a \in fn(Q)
 \end{aligned}$$

In the former case  $a$  is not a free name in  $P$  and  $Q$  so the substitutions have no effects and the lemma holds. In the latter case  $a$  is a free names in both processes: the proof goes by induction on the length of the proof tree of  $P \equiv_{\alpha} Q$  and then by cases on the last rule of the proof tree. Let  $x, y, a$  and  $b$  be pairwise different.

*base case* The length of the proof is one and the rule used can be only: *AlpZero* or *AlpIde*: the lemma holds because  $P$  and  $Q$  are syntactically the same process.

*inductive case* :

*AlpTau* :

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 && \text{rule premise} \\
 \Rightarrow P_1\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} && \text{inductive hypothesis} \\
 \Rightarrow \tau.(P_1\{b/a\}) &\equiv_{\alpha} \tau.(Q_1\{b/a\}) && \text{rule AlpTau} \\
 \Rightarrow (\tau.P_1)\{b/a\} &\equiv_{\alpha} (\tau.Q_1)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

*AlpSum* :

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 \text{ and } P_2 \equiv_{\alpha} Q_2 && \text{rule premises} \\
 \Rightarrow P_1\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} \text{ and } P_2\{b/a\} \equiv_{\alpha} Q_2\{b/a\} && \text{inductive hypothesis} \\
 \Rightarrow P_1\{b/a\} + P_2\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} + Q_2\{b/a\} && \text{rule AlpSum} \\
 \Rightarrow (P_1 + P_2)\{b/a\} &\equiv_{\alpha} (Q_1 + Q_2)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

*AlpPar* : this case is very similar to the previous one.

*AlpOut* :

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 && \text{rule premise} \\
 \Rightarrow P_1\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} && \text{inductive hypothesis} \\
 \Rightarrow \bar{x}\{b/a\}y\{b/a\}.P_1\{b/a\} &\equiv_{\alpha} \bar{x}\{b/a\}y\{b/a\}.Q_1\{b/a\} && \text{rule AlpOut} \\
 \Rightarrow (\bar{x}y.P_1)\{b/a\} &\equiv_{\alpha} (\bar{x}y.Q_1)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

*AlpInp* :

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 && \text{rule premise} \\
 \Rightarrow P_1\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} && \text{inductive hypothesis} \\
 \Rightarrow x\{b/a\}(y).P_1\{b/a\} &\equiv_{\alpha} x\{b/a\}(y).Q_1\{b/a\} && \text{rule AlpInp} \\
 \Rightarrow (x(y).P_1)\{b/a\} &\equiv_{\alpha} (x(y).Q_1)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 && \text{rule premise} \\
 \Rightarrow b(a).P_1 &\equiv_{\alpha} b(a).Q_1 && \text{rule AlpIn} \\
 \Rightarrow a\{b/a\}(a).P_1 &\equiv_{\alpha} a\{b/a\}(a).Q_1 && \text{definition of substitution} \\
 \Rightarrow (a(a).P_1)\{b/a\} &\equiv_{\alpha} (a(a).Q_1)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

$$\begin{aligned}
 P_1 &\equiv_{\alpha} Q_1 && \text{rule premise} \\
 \Rightarrow P_1\{b/a\} &\equiv_{\alpha} Q_1\{b/a\} && \text{inductive hypothesis} \\
 \Rightarrow b\{b/a\}(x).(P_1\{b/a\}) &\equiv_{\alpha} b\{b/a\}(x).(Q_1\{b/a\}) && \text{rule AlpIn} \\
 \Rightarrow (b(x).P_1)\{b/a\} &\equiv_{\alpha} (b(x).Q_1)\{b/a\} && \text{definition of substitution}
 \end{aligned}$$

*AlpInp1* : we have various cases:

- the last part of the proof tree of  $P \equiv_\alpha Q$  is

$$\text{ALPINP1} \frac{P_1 \equiv_\alpha Q_1\{x/y\} \quad x \neq y \quad x \notin \text{fn}(Q_1)}{\underbrace{z(x).P_1}_P \equiv_\alpha \underbrace{z(y).Q_1}_Q}$$

$$\begin{array}{ll} P_1 \equiv_\alpha Q_1\{x/y\} \text{ and } x \notin \text{fn}(Q_1) & \text{rule premise} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{x/y\}\{b/a\} & \text{inductive hypothesis} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{x/y\} & \text{transitivity and lemma 2.3.5} \\ \Rightarrow z(x).(P_1\{b/a\}) \equiv_\alpha z(y).(Q_1\{b/a\}) & \text{rule } \textit{AlpInp1} \\ \Rightarrow (z(x).P_1)\{b/a\} \equiv_\alpha (z(y).Q_1)\{b/a\} & \text{definition of substitution} \end{array}$$

- the last part of the proof tree of  $P \equiv_\alpha Q$  is

$$\text{ALPINP1} \frac{P_1 \equiv_\alpha Q_1\{x/y\} \quad x \neq y \quad x \notin \text{fn}(Q_1)}{\underbrace{b(x).P_1}_P \equiv_\alpha \underbrace{b(y).Q_1}_Q}$$

$$\begin{array}{ll} P_1 \equiv_\alpha Q_1\{x/y\} \text{ and } x \notin \text{fn}(Q_1) & \text{rule premise} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{x/y\}\{b/a\} & \text{inductive hypothesis} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{x/y\} & \text{transitivity and lemma 2.3.5} \\ \Rightarrow b(x).(P_1\{b/a\}) \equiv_\alpha b(y).(Q_1\{b/a\}) & \text{rule } \textit{AlpInp1} \\ \Rightarrow (b(x).P_1)\{b/a\} \equiv_\alpha (b(y).Q_1)\{b/a\} & \text{definition of substitution} \end{array}$$

- the last part of the proof tree of  $P \equiv_\alpha Q$  is

$$\text{ALPINP1} \frac{P_1 \equiv_\alpha Q_1\{x/y\} \quad x \neq y \quad x \notin \text{fn}(Q_1)}{\underbrace{a(x).P_1}_P \equiv_\alpha \underbrace{a(y).Q_1}_Q}$$

$$\begin{array}{ll} P_1 \equiv_\alpha Q_1\{x/y\} \text{ and } x \notin \text{fn}(Q_1) & \text{rule premise} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{x/y\}\{b/a\} & \text{inductive hypothesis} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{x/y\} & \text{transitivity and lemma 2.3.5} \\ \Rightarrow a(x).(P_1\{b/a\}) \equiv_\alpha a(y).(Q_1\{b/a\}) & \text{rule } \textit{AlpInp1} \\ \Rightarrow (a(x).P_1)\{b/a\} \equiv_\alpha (a(y).Q_1)\{b/a\} & \text{definition of substitution} \end{array}$$

- the last part of the proof tree of  $P \equiv_\alpha Q$  is

$$\text{ALPINP1} \frac{P_1 \equiv_\alpha Q_1\{a/y\} \quad a \neq y \quad a \notin \text{fn}(Q_1)}{\underbrace{a(a).P_1}_P \equiv_\alpha \underbrace{a(y).Q_1}_Q}$$

$$\begin{array}{ll} P_1 \equiv_\alpha Q_1\{a/y\} \text{ and } x \notin \text{fn}(Q_1) & \text{rule premise} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{a/y\}\{b/a\} & \text{inductive hypothesis} \\ \Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{a/y\} & \text{transitivity and lemma 2.3.5} \\ \Rightarrow a(a).(P_1\{b/a\}) \equiv_\alpha a(y).(Q_1\{b/a\}) & \text{rule } \textit{AlpInp1} \\ \Rightarrow (a(a).P_1)\{b/a\} \equiv_\alpha (a(y).Q_1)\{b/a\} & \text{definition of substitution} \end{array}$$

- the last part of the proof tree of  $P \equiv_\alpha Q$  is

$$\text{ALPINP1} \frac{P_1 \equiv_\alpha Q_1\{x/a\} \quad x \neq a \quad x \notin \text{fn}(Q_1)}{\underbrace{a(x).P_1}_P \equiv_\alpha \underbrace{a(a).Q_1}_Q}$$

$P_1 \equiv_\alpha Q_1\{x/a\}$ and $x \notin fn(Q_1)$	rule premise
$\Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{x/a\}\{b/a\}$	inductive hypothesis
$\Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{x/a\}$	transitivity and lemma 2.3.5
$\Rightarrow a(x).(P_1\{b/a\}) \equiv_\alpha a(a).(Q_1\{b/a\})$	rule <i>AlpInp1</i>
$\Rightarrow (a(x).P_1)\{b/a\} \equiv_\alpha (a(a).Q_1)\{b/a\}$	definition of substitution
• mancano x x y e x y x	

*AlpRes* :

$P_1 \equiv_\alpha Q_1$	rule premise
$\Rightarrow P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}$	inductive hypothesis
$\Rightarrow (\nu x)(P_1\{b/a\}) \equiv_\alpha (\nu x)(Q_1\{b/a\})$	rule <i>AlpRes</i>
$\Rightarrow ((\nu x)P_1)\{b/a\} \equiv_\alpha ((\nu x)Q_1)\{b/a\}$	definition of substitution

*AlpRes1* :

$$\text{ALPRES1} \frac{P_1 \equiv_\alpha Q_1\{x/y\} \quad x \neq y \quad x \notin fn(Q_1)}{\underbrace{(\nu x)P_1}_P \equiv_\alpha \underbrace{(\nu y)Q_1}_Q}$$

$P_1 \equiv_\alpha Q_1\{x/y\}$ and $x \neq y$ and $x \notin fn(Q_1)$	rule premises
$P_1\{b/a\} \equiv_\alpha Q_1\{x/y\}\{b/a\}$	inductive hypothesis
$P_1\{b/a\} \equiv_\alpha Q_1\{b/a\}\{x/y\}$	lemma 2.3.5 and transitivity
$(\nu x)(P_1\{b/a\}) \equiv_\alpha (\nu y)(Q_1\{b/a\})$	rule <i>AlpRes1</i>
$((\nu x)P_1)\{b/a\} \equiv_\alpha ((\nu y)Q_1)\{b/a\}$	definition of substitution

□

**Lemma 2.3.7.**

$$P \equiv_\alpha P\{x/y\}\{y/x\}$$

esistono delle precondizioni per le quali il lemma e' vero? esistono delle precondizioni per le quali si puo' addirittura avere l'uguaglianza sintattica?

In the proof of equivalence of the semantics in the next section we need the following lemmas

**Lemma 2.3.8.**  $P\{x/y\} \equiv_\alpha Q$  if and only if  $P \equiv_\alpha Q\{y/x\}$ .

NON FUNZIONA LA DIMOSTRAZIONE! staro' forse esagerando?

*Proof.* The proof is an induction on the length of the proof tree of  $P\{x/y\} \equiv_\alpha Q$  and then by cases on the last rule:

**base case** the last rule can be

*AlpZero* in this case both  $P$  and  $Q$  are the null process 0 so the thesis holds.

*AlpIde* for this rule to apply  $P\{x/y\}$  and  $Q$  must be some identifier  $A$  with the same variable.

Suppose that  $P = A(\tilde{a}|\tilde{b})$  There can be some different cases:

$y \in \tilde{a}$  we can suppose that  $\tilde{a} = y, \tilde{c}$  then

$x \in \tilde{b}$  we can suppose that  $\tilde{b} = x, \tilde{d}$ , then

$$Q = P\{x/y\} = A(x, \tilde{c}|z, \tilde{d})$$

where  $z$  is a fresh name. We need now the identifier equal to  $Q\{y/x\} = A(x, \tilde{c}|z, \tilde{d})\{y/x\}$  so we have to distinguish two cases:

$x \in \text{tilded}$

$x \notin \text{tilded}$

$$Q\{y/x\} = A(x, \tilde{c}|z, \tilde{d})\{y/x\} = A(y, \tilde{c}|z, \tilde{d})$$

$y \notin \tilde{y}$  in this case there is no need to change bound names so

$$Q\{y/x\} = A(y, \tilde{z}|\tilde{y})$$

$x \notin \tilde{x}$  then

$$Q\{y/x\} = Q = A(\tilde{x}|\tilde{y})$$

□

**Lemma 2.3.9.** The  $\alpha$  equivalence is an equivalence relation.

*Proof.* :

**reflexivity** We prove  $P \equiv_\alpha P$  by structural induction on  $P$ :

0 :

$$\text{ALPZERO} \frac{}{0 \equiv_\alpha 0}$$

$\tau.P_1$  : for induction  $P_1 \equiv_\alpha P_1$  so

$$\text{ALPTAU} \frac{P_1 \equiv_\alpha P_1}{\tau.P_1 \equiv_\alpha \tau.P_1}$$

$x(y).P_1$  : for induction  $P_1 \equiv_\alpha P_1$  so

$$\text{ALPINP} \frac{P_1 \equiv_\alpha P_1}{x(y).P_1 \equiv_\alpha x(y).P_1}$$

$\bar{x}y.P_1$  : for induction  $P_1 \equiv_\alpha P_1$  so

$$\text{ALPOUT} \frac{P_1 \equiv_\alpha P_1}{\bar{x}y.P_1 \equiv_\alpha \bar{x}y.P_1}$$

$P_1 + P_2$  : for induction  $P_1 \equiv_\alpha P_1$  and  $P_2 \equiv_\alpha P_2$  so

$$\text{ALPSUM} \frac{P_1 \equiv_\alpha P_1 \quad P_2 \equiv_\alpha P_2}{P_1 + P_2 \equiv_\alpha P_1 + P_2}$$

$P_1|P_2$  : for induction  $P_1 \equiv_\alpha P_1$  and  $P_2 \equiv_\alpha P_2$  so

$$\text{ALPPAR} \frac{P_1 \equiv_\alpha P_1 \quad P_2 \equiv_\alpha P_2}{P_1|P_2 \equiv_\alpha P_1|P_2}$$

$(\nu x)P_1$  : for induction  $P_1 \equiv_\alpha P_1$  so

$$\text{ALPRES} \frac{P_1 \equiv_\alpha P_1}{(\nu x)P_1 \equiv_\alpha (\nu x)P_1}$$

$A(\tilde{x}|\tilde{y})$  :

$$\text{ALPIDE} \frac{}{A(\tilde{x}|\tilde{y}) \equiv_\alpha A(\tilde{x}|\tilde{y})}$$

**symmetry** A proof of

$$P \equiv_\alpha Q \Rightarrow Q \equiv_\alpha P$$

can go by induction on the length of the proof tree of  $P \equiv_\alpha Q$  and then by cases on the last rule used. Nevertheless we notice that the base case rules *AlpZero* and *AlpIde* are symmetric and the inductive case rules are symmetric except for *AlpRes1* and *AlpInp1*. So we provide with the cases for those last two rules:

*AlpRes1* the last part of the proof tree is

$$\text{ALPRES1} \frac{P\{y/x\} \equiv_{\alpha} Q \quad x \neq y \quad y \notin \text{fn}(P)}{(\nu x)P \equiv_{\alpha} (\nu y)Q}$$

we apply the inductive hypothesis on  $P\{y/x\} \equiv_{\alpha} Q$  and get  $Q \equiv_{\alpha} P\{y/x\}$  which implies  $Q\{x/y\} \equiv_{\alpha} P$

DA DIMOSTRARE  $Q \equiv_{\alpha} P\{y/x\}$  and  $y \notin \text{fn}(P)$  implies  $Q\{x/y\} \equiv_{\alpha} P$  and  $x \notin \text{fn}(Q)$

so an application of the same rule yields:

$$\text{ALPRES1} \frac{Q\{x/y\} \equiv_{\alpha} P \quad x \neq y \quad x \notin \text{fn}(Q)}{(\nu y)QP \equiv_{\alpha} (\nu x)Q}$$

*AlpInp1* this is very similar to the previous.

**transitivity** suppose

$$P \equiv_{\alpha} Q \text{ and } Q \equiv_{\alpha} R$$

we prove the thesis  $P \equiv_{\alpha} R$  by induction on the length of the proof tree of  $P \equiv_{\alpha} Q$ . If the tree has only one node then the rule used must be *AlpZero* or *AlpIde*. In the former case both  $P$  and  $Q$  are 0 and so  $0 \equiv_{\alpha} R$ . For symmetry and the inversion lemma then  $R$  is also 0. In the latter case a similar argument applies. If the proof tree has more than one node then we proceed by cases on the last rule

*AlpInp* : In this case  $P = x(y).P_1$ ,  $Q = x(y).Q_1$  and  $P_1 \equiv_{\alpha} Q_1$  and  $x(y).Q_1 \equiv_{\alpha} R$  which implies for symmetry and the inversion lemma that one of the following cases holds:

- $R = x(y).R_1$  and  $Q_1 \equiv_{\alpha} R_1$ :
 

$P_1 \equiv_{\alpha} Q_1$ and $Q_1 \equiv_{\alpha} R_1$	inductive hypothesis
$\Rightarrow P_1 \equiv_{\alpha} R_1$	rule <i>AlpInp</i>
$\Rightarrow x(y).P_1 \equiv_{\alpha} x(y).R_1$	
- $R = x(z).R_1$  and  $Q_1\{y/z\} \equiv_{\alpha} R_1$ :
 

$P_1 \equiv_{\alpha} Q_1$	lemma 2.3.6
$\Rightarrow P_1\{y/z\} \equiv_{\alpha} Q_1\{y/z\}$	inductive hypothesis
$\Rightarrow P_1\{y/z\} \equiv_{\alpha} R_1$	rule <i>AlpInp1</i>
$\Rightarrow x(y).P_1 \equiv_{\alpha} x(z).R_1$	

*AlpRes* :

*AlpInp1* :

*AlpRes1* :

*AlpSum* :

*AlpPar* :

*AlpSum* :

*AlpTau* :

*AlpOut* :

□

**Lemma 2.3.10.** E' FALSO!!!! !!!!! !! !:

- If  $P \equiv \tau.P_1$  then  $P = \tau.P_1$  for some  $P_1$  such that  $P_1 \equiv Q$
- If  $P \equiv \bar{x}y.P_1$  then  $P = \bar{x}y.P_1$  for some  $P_1$  such that  $P_1 \equiv Q$
- If  $P \equiv x(y).Q$  then one and only one of the following cases holds:

---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(y).P \xrightarrow{xz} P\{z/y\}}$
<b>ParL</b> $\frac{P \xrightarrow{\alpha} P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P' Q}$	<b>ParR</b> $\frac{Q \xrightarrow{\alpha} Q' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P Q'}$
<b>SumL</b> $\frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'}$	<b>SumR</b> $\frac{Q \xrightarrow{\alpha} Q'}{P + Q \xrightarrow{\alpha} Q'}$
<b>Res</b> $\frac{P \xrightarrow{\alpha} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha} (\nu z)P'}$	<b>Alp</b> $\frac{P \equiv_{\alpha} Q \quad P \xrightarrow{\alpha} P'}{Q \xrightarrow{\alpha} P'}$
<b>EComL</b> $\frac{P \xrightarrow{xy} P' \quad Q \xrightarrow{\bar{x}y} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>EComR</b> $\frac{P \xrightarrow{\bar{x}y} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\tau} P' Q'}$
<b>ClsL</b> $\frac{P \xrightarrow{\bar{x}(z)} P' \quad Q \xrightarrow{xz} Q' \quad z \notin fn(Q)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$	<b>ClsR</b> $\frac{P \xrightarrow{xz} P' \quad Q \xrightarrow{\bar{x}(z)} Q' \quad z \notin fn(P)}{P Q \xrightarrow{\tau} (\nu z)(P' Q')}$
<b>Cns</b> $\frac{A(\tilde{x} \tilde{y}) \stackrel{def}{=} P \quad P\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha} P'}{A(\tilde{x} \tilde{y})\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha} P'}$	<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$
<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$	

---

Table 2.8: Early transition relation with  $\alpha$  conversion

- $P = x(z).P_1$  for some  $P_1$  such that  $P_1\{z/y\} \equiv Q$
- $P = x(y).P_1$  for some  $P_1$  such that  $P_1 \equiv Q$
- If  $P \equiv Q_1 + Q_2$  then  $P = P_1 + P_2$  for some  $P_1$  and  $P_2$  such that  $P_1 \equiv Q_1$  and  $P_2 \equiv Q_2$ .
- If  $P \equiv Q_1|Q_2$  then  $P = P_1|P_2$  for some  $P_1$  and  $P_2$  such that  $P_1 \equiv Q_1$  and  $P_2 \equiv Q_2$ .
- If  $P \equiv (\nu y)Q$  then one and only one of the following cases holds:
  - $P = (\nu z)P_1$  such that  $P_1\{z/y\} \equiv Q$
  - $P = (\nu y).P_1$  for some  $P_1$  such that  $P_1 \equiv Q$
- If  $P \equiv A(\tilde{x}|\tilde{y})$  then ??? ?? ?

*Proof.*

□

## 2.4 Operational semantic with structural congruence

### 2.4.1 Early semantic with $\alpha$ conversion only

In this subsection we introduce the early operational semantic for  $\pi$  calculus with the use of a minimal structural congruence, specifically we exploit only the easy of  $\alpha$  conversion.

**Definition 2.4.1.** The *early transition relation with  $\alpha$  conversion*  $\rightarrow_{\subseteq} \mathbb{P} \times \mathbb{A} \times \mathbb{P}$  is the smallest relation induced by the rules in table 2.8.



---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(z).P \xrightarrow{xy} P\{y/z\}}$	<b>Par</b> $\frac{P \xrightarrow{\alpha} P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P' Q}$
<b>Sum</b> $\frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'}$	<b>ECom</b> $\frac{P \xrightarrow{xy} P' \quad Q \xrightarrow{\bar{x}y} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>Res</b> $\frac{P \xrightarrow{\alpha} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha} (\nu z)P'}$
<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$	<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$	<b>Str</b> $\frac{P \equiv P' \quad P \xrightarrow{\alpha} Q \quad Q \equiv Q'}{P' \xrightarrow{\alpha} Q'}$

---

Table 2.9: Early semantic with structural congruence

### 2.4.2 Early semantic with structural congruence

**Definition 2.4.2.** The *early transition relation with structural congruence*  $\rightarrow_{\subseteq} \mathbb{P} \times \mathbb{A} \times \mathbb{P}$  is the smallest relation induced by the rules in table 2.9.

**Example** We prove now that

$$a(x).P \mid (\nu b)\bar{a}b.Q \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)$$

where  $b \notin fn(P)$ . This follows from

$$a(x).P \mid (\nu b)\bar{a}b.Q \equiv (\nu b)(a(x).P \mid \bar{a}b.Q)$$

and

$$(\nu b)(a(x).P \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)$$

with the rule *Str*. We can prove the last transition in the following way:

$$\text{RES} \frac{\text{COM} \frac{\text{EINP} \frac{}{a(x).P \xrightarrow{ab} P\{b/x\}} \quad \text{OUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q}}{a(x).P \mid \bar{a}b.Q \xrightarrow{\tau} P\{b/x\} \mid Q}}{(\nu b)(a(x).P \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)}$$

**Example** We want to prove now that:

$$((\nu b)a(x).P) \mid \bar{a}b.Q \xrightarrow{\tau} (\nu c)(P\{c/b\}\{b/x\} \mid Q)$$

where the name  $c$  is not in the free names of  $Q$ . We can exploit the structural congruence and get that

$$((\nu b)a(x).P) \mid \bar{a}b.Q \equiv (\nu c)(a(x).(P\{c/b\}) \mid \bar{a}b.Q)$$

then we have

$$\text{RES} \frac{\text{COM} \frac{\text{EINP} \frac{}{a(x).P\{c/b\} \xrightarrow{ab} P\{c/b\}\{b/x\}} \quad \text{OUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q}}{a(x).(P\{c/b\}) \mid \bar{a}b.Q \xrightarrow{\tau} (P\{c/b\}\{b/x\} \mid Q)}}{(\nu c)(a(x).(P\{c/b\}) \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu c)(P\{c/b\}\{b/x\} \mid Q)}$$

Now we just apply the rule *Str* to prove the thesis.

### 2.4.3 Late semantic with structural congruence

**Definition 2.4.3.** The *late transition relation with structural congruence*  $\rightarrow_{\subseteq} \mathbb{P} \times \mathbb{A} \times \mathbb{P}$  is the smallest relation induced by the rules in table 2.10.

**Example** We prove now that

---

<b>Prf</b> $\frac{}{\alpha.P \xrightarrow{\alpha} P}$	<b>Sum</b> $\frac{P \xrightarrow{\alpha} P'}{P + Q \xrightarrow{\alpha} P'}$
<b>Par</b> $\frac{P \xrightarrow{\alpha} P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\alpha} P' Q}$	<b>Res</b> $\frac{P \xrightarrow{\alpha} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha} (\nu z)P'}$
<b>LCom</b> $\frac{P \xrightarrow{x(y)} P' \quad Q \xrightarrow{\bar{x}z} Q'}{P Q \xrightarrow{\tau} P'\{z/y\} Q'}$	<b>Str</b> $\frac{P \equiv P' \quad P \xrightarrow{\alpha} Q \quad Q \equiv Q'}{P' \xrightarrow{\alpha} Q'}$
<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$	

---

Table 2.10: Late semantic with structural congruence

$$a(x).P \mid (\nu b)\bar{a}b.Q \xrightarrow{\tau} P\{b/x\} \mid Q$$

where  $b \notin fn(P)$ . This follows from

$$a(x).P \mid (\nu b)\bar{a}b.Q \equiv (\nu b)(a(x).P \mid \bar{a}b.Q)$$

and

$$(\nu b)(a(x).P \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)$$

with the rule *Str*. We can prove the last transition in the following way:

$$\text{RES} \frac{\text{LCOM} \frac{\text{LINP} \frac{b \notin fn(P)}{a(x).P \xrightarrow{ab} P\{b/x\}} \quad \text{OUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q}}{a(x).P \mid \bar{a}b.Q \xrightarrow{\tau} P\{b/x\} \mid Q} \quad b \notin n(\tau)}{(\nu b)(a(x).P \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu b)(P\{b/x\} \mid Q)}$$

**Example** We want to prove now that:

$$((\nu b)a(x).P) \mid \bar{a}b.Q \xrightarrow{\tau} (\nu c)(P\{c/b\}\{b/x\} \mid Q)$$

where the name  $c$  is not in the free names of  $Q$  and is not in the names of  $P$ . We can exploit the structural congruence and get that

$$((\nu b)a(x).P) \mid \bar{a}b.Q \equiv (\nu c)(a(x).(P\{c/b\}) \mid \bar{a}b.Q)$$

then we have

$$\text{RES} \frac{\text{LCOM} \frac{\text{LINP} \frac{b \notin fn(P\{c/b\})}{a(x).P\{c/b\} \xrightarrow{ab} P\{c/b\}\{b/x\}} \quad \text{OUT} \frac{}{\bar{a}b.Q \xrightarrow{\bar{a}b} Q}}{a(x).(P\{c/b\}) \mid \bar{a}b.Q \xrightarrow{\tau} (P\{c/b\}\{b/x\} \mid Q)} \quad c \notin n(\tau)}{(\nu c)(a(x).(P\{c/b\}) \mid \bar{a}b.Q) \xrightarrow{\tau} (\nu c)(P\{c/b\}\{b/x\} \mid Q)}$$

Now we just apply the rule *Str* to prove the thesis.

## 2.5 Equivalence of the semantics

### 2.5.1 Equivalence of the early semantics

In this subsection we write  $\rightarrow_1$  for the early semantic without structural congruence,  $\rightarrow_2$  for the early semantic with just  $\alpha$  conversion and  $\rightarrow_3$  for the early semantic with the full structural congruence. We call  $R_1$  the set of rules for  $\rightarrow_1$ ,  $R_2$  the set of rules for  $\rightarrow_2$  and  $R_3$  the set of rules for  $\rightarrow_3$ . In the following section we will need:

**Lemma 2.5.1.**

$$P \equiv Q \Rightarrow fn(Q) = fn(P)$$

*Proof.* A proof can go by induction on the proof tree of  $P \equiv Q$  and then by cases on the last rule used in the proof tree.

**base case** The last and only rule of the proof tree can be one of the following axioms:

$$\begin{array}{ll} \text{SC-ALP} & \frac{P \equiv_{\alpha} Q}{P \equiv Q} \\ \text{SC-SUM-ASC} & M_1 + (M_2 + M_3) \equiv (M_1 + M_2) + M_3 \\ \text{SC-SUM-COM} & M_1 + M_2 \equiv M_2 + M_1 \\ \text{SC-SUM-INC} & M + 0 \equiv M \\ \text{SC-COM-ASC} & P_1|(P_2|P_3) \equiv (P_1|P_2)|P_3 \\ \text{SC-COM-COM} & P_1|P_2 \equiv P_2|P_1 \\ \text{SC-COM-INC} & P|0 \equiv P \\ \text{SC-RES} & (\nu z)(\nu w)P \equiv (\nu w)(\nu z)P \\ \text{SC-RES-INC} & (\nu z)0 \equiv 0 \\ \text{SC-RES-COM} & (\nu z)(P_1|P_2) \equiv P_1|(\nu z)P_2 \text{ if } z \notin fn(P_1) \\ \text{SC-RES-SUM} & (\nu z)(P_1 + P_2) \equiv P_1 + (\nu z)P_2 \text{ if } z \notin fn(P_1) \\ \text{SC-IDE} & A(\tilde{w}|\tilde{y}) \equiv P\{\tilde{w}/\tilde{x}\} \end{array}$$

**inductive case**

$$\text{SC-REFL} \quad P \equiv P$$

$$\text{SC-SIMM} \quad \frac{Q \equiv P}{P \equiv Q}$$

$$\text{SC-TRAN} \quad \frac{P \equiv Q \quad Q \equiv R}{P \equiv R}$$

$$\text{SC-CONG} \quad \frac{P \equiv Q}{C[P] \equiv C[Q]}$$

□

We would like to prove that  $P \xrightarrow{\alpha}_2 P' \Rightarrow P \xrightarrow{\alpha}_1 P'$  but this is false because

$$\text{ALP} \quad \frac{\overline{xy}.x(y).0 \equiv_{\alpha} \overline{xy}.x(w).0 \quad \text{OUT} \quad \overline{xy}.x(w).0 \xrightarrow{\overline{xy}}_2 x(w).0}{\overline{xy}.x(y).0 \xrightarrow{\overline{xy}}_2 x(w).0}$$

so we want to prove

$$\overline{xy}.x(y).0 \xrightarrow{\overline{xy}}_1 x(w).0$$

The head of the transition has an output prefixing at the top level so the only rule we could use is *Out*, but the application of *Out* yields

$$\overline{xy}.x(y).0 \xrightarrow{\overline{xy}}_1 x(y).0$$

which is not what we want. So we prove a weaker version

**Theorem 2.5.2.**

$$P \xrightarrow{\alpha}_2 P' \Rightarrow \exists P'' : P'' \equiv_{\alpha} P' \text{ and } P \xrightarrow{\alpha}_1 P''$$

*Proof.* The proof goes by induction on the depth of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  and then by cases on the last rule used:

**base case** If the depth of the derivation tree is one, the rule used has to be a prefix rule

$$\{Out, EInp, Tau\} \subseteq R_1 \cap R_2$$

so a derivation tree of  $P \xrightarrow{\alpha}_2 P'$  is also a derivation tree of  $P \xrightarrow{\alpha}_1 P'$

**inductive case** If the depth of the derivation tree is more than one, then we proceed by cases on the last rule  $R$ . If the rule  $R$  is not a prefix rule and it is in common between the two semantics:

$$R \in \{ParL, ParR, SumL, SumR, Res, EComL, EComR, ClsL, ClsR, Cns, Opn\}$$

then we just apply the inductive hypothesis on the premises of  $R$  and then reapply  $R$  to get the desired derivation tree. We show just the case for  $SumL$  when the end of the derivation tree is

$$\text{SUML} \frac{P_1 \xrightarrow{\alpha}_2 P'_1}{\underbrace{P_1 + P_2}_P \xrightarrow{\alpha}_2 \underbrace{P'_1}_{P'}}$$

rule premise  
inductive hypothesis  
rule  $SumL$

$$\begin{aligned} &P_1 \xrightarrow{\alpha}_2 P'_1 \\ \Rightarrow &P_1 \xrightarrow{\alpha}_1 P''_1 \text{ and } P'_1 \equiv_{\alpha} P''_1 \\ \Rightarrow &P_1 + P_2 \xrightarrow{\alpha}_1 P''_1 \end{aligned}$$

If the rule  $R$  is in

$$R_2 - R_1 = \{Alp\}$$

then the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  is

$$\text{ALP} \frac{P \equiv_{\alpha} Q \quad \text{S} \frac{\dots}{Q \xrightarrow{\alpha}_2 P'}}{P \xrightarrow{\alpha}_2 P'}$$

and the proof goes by cases on  $S$  the last rule in the proof tree of  $Q \xrightarrow{\alpha}_2 P'$ :

**Out** : If  $S = Out$  then there exists some names  $x, y$  and a process  $Q_1$  such that

$$Q = \bar{x}y.Q_1$$

and  $\alpha = \bar{x}y$ .

$$\begin{aligned} &P \equiv_{\alpha} \bar{x}y.Q_1 && \text{inversion lemma} \\ \Rightarrow &P = \bar{x}y.P_1 \text{ and } P_1 \equiv_{\alpha} Q_1 && \text{rule } Out \\ \Rightarrow &\bar{x}y.P_1 \xrightarrow{\bar{x}y}_1 P_1 \end{aligned}$$

**EInp** If  $S = EInp$  then there exists some names  $x, y, z$  and a process  $Q_1$  such that  $Q = x(y).Q_1$ ,  $\alpha = xz$  and  $P' = Q_1\{z/y\}$ . Since

$$P \equiv_{\alpha} x(y).Q_1$$

then for the inversion lemma we have two cases:

• :

$$\begin{aligned} &P = x(y).P_1 \text{ and } P_1 \equiv_{\alpha} Q_1 && \text{rule } EInp \\ \Rightarrow &x(y).P_1 \xrightarrow{xz}_1 P_1\{z/y\} \end{aligned}$$

This is what we want because for lemma 2.3.6

$$P_1 \equiv_\alpha Q_1 \Rightarrow P_1\{z/y\} \equiv_\alpha Q_1\{z/y\}$$

• :

$$\begin{aligned} P &= x(w).P_1 \text{ and } P_1\{y/w\} \equiv_\alpha Q_1 && \text{rule } EInp \\ \Rightarrow x(w).P_1 &\xrightarrow{xz}_1 P_1\{z/w\} \end{aligned}$$

This is what we want because

$$\begin{aligned} P_1\{y/w\} &\equiv_\alpha Q_1 && \text{lemma 2.3.6} \\ \Rightarrow P_1\{y/w\}\{z/y\} &\equiv_\alpha Q_1\{z/y\} \\ \Rightarrow P_1\{z/w\} &\equiv_\alpha Q_1\{z/y\} \end{aligned}$$

**Tau** If  $S = Tau$  then there exists a process  $Q_1$  such that  $Q = \tau.Q_1$  and  $\alpha = \tau$  and  $P' = Q_1$ .

$$\begin{aligned} P &\equiv_\alpha \tau.Q_1 && \text{inversion lemma} \\ \Rightarrow P &= \tau.P_1 \text{ and } P_1 \equiv_\alpha Q_1 && \text{rule } Tau \\ \Rightarrow \tau.P_1 &\xrightarrow{\tau}_1 P_1 \end{aligned}$$

**ParL** If  $S = ParL$  then there exists some processes  $Q_1, Q_2$  such that

$$Q = Q_1|Q_2$$

Since

$$P \equiv_\alpha Q_1|Q_2$$

then for the inversion lemma there exists  $P_1, P_2$  such that

$$P = P_1|P_2 \text{ and } P_1 \equiv_\alpha Q_1 \text{ and } P_2 \equiv_\alpha Q_2$$

and so the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  looks like this:

$$\text{ALP} \frac{P_1|P_2 \equiv_\alpha Q_1|Q_2 \quad \text{PARL} \frac{Q_1 \xrightarrow{\alpha}_2 Q'_1 \quad bn(\alpha) \cap fn(Q_2) = \emptyset}{Q_1|Q_2 \xrightarrow{\alpha}_2 Q'_1|Q_2}}{\underbrace{P_1|P_2}_P \xrightarrow{\alpha}_2 \underbrace{Q'_1|Q_2}_{P'}}$$

from this hypothesis we can create the following proof tree of  $P_1 \xrightarrow{\alpha}_2 Q'_1$ :

$$\text{ALP} \frac{P_1 \equiv_\alpha Q_1 \quad Q_1 \xrightarrow{\alpha}_2 Q'_1}{P_1 \xrightarrow{\alpha}_2 Q'_1}$$

this proof tree is smaller than the proof tree of  $P_1|P_2 \xrightarrow{\alpha}_2 Q'_1|Q_2$  so we can apply the inductive hypothesis and get that there exists a process  $Q'_1$  such that

$$Q'_1 \equiv Q''_1 \text{ and } P_1 \xrightarrow{\alpha}_1 Q''_1$$

then we apply again the rule *ParL* and get

$$\text{PARL} \frac{P_1 \xrightarrow{\alpha}_1 Q''_1 \quad bn(\alpha) \cap fn(P_2) = \emptyset}{\underbrace{P_1|P_2}_P \xrightarrow{\alpha}_1 \underbrace{Q''_1|P_2}_{P''}}$$

The second premise of the previous instance holds because:

$$bn(\alpha) \cap fn(Q_2) = \emptyset \text{ and } P_2 \equiv_\alpha Q_2 \Rightarrow bn(\alpha) \cap fn(P_2) = \emptyset$$

**ParR, SumL, SumR, EComL, EComR, ClsL, ClsR** This cases are similar to the previous.

**Res** If  $S = Res$  then there exists some name  $z$  and a process  $Q_1$  such that

$$Q = (\nu z)Q_1$$

and  $P' = (\nu z)Q_1'$ . Since

$$P \equiv_\alpha (\nu z)Q_1$$

then for the inversion lemma we have two cases:

- there exists some  $P_1$  such that

$$P = (\nu z)P_1 \text{ and } P_1 \equiv_\alpha Q_1$$

and so the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  looks like this:

$$\text{ALP} \frac{(\nu z)P_1 \equiv_\alpha (\nu z)Q_1 \quad \text{RES} \frac{Q_1 \xrightarrow{\alpha}_2 Q_1' \quad z \notin n(\alpha)}{(\nu z)Q_1 \xrightarrow{\alpha}_2 (\nu z)Q_1'}}{(\nu z)P_1 \xrightarrow{\alpha}_2 (\nu z)Q_1'}$$

from this we create the following proof tree of  $P_1 \xrightarrow{\alpha}_2 Q_1'$ :

$$\text{ALP} \frac{P_1 \equiv_\alpha Q_1 \quad Q_1 \xrightarrow{\alpha}_2 Q_1'}{P_1 \xrightarrow{\alpha}_2 Q_1'}$$

to which we can apply the inductive hypothesis and get that there exists a process  $Q_1''$  such that

$$P_1 \xrightarrow{\alpha}_1 Q_1'' \text{ and } Q_1'' \equiv_\alpha Q_1'$$

then we apply the rule *Res* to get

$$\text{RES} \frac{P_1 \xrightarrow{\alpha}_1 Q_1'' \quad z \notin n(\alpha)}{(\nu z)P_1 \xrightarrow{\alpha}_1 (\nu z)Q_1''}$$

this satisfies the thesis of the theorem because

$$(\nu z)Q_1'' \equiv (\nu z)Q_1'$$

- there exists some  $P_1$  such that

$$P = (\nu y)P_1 \text{ and } P_1\{z/y\} \equiv_\alpha Q_1$$

and so the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  looks like this:

$$\text{ALP} \frac{(\nu y)P_1 \equiv_\alpha (\nu z)Q_1 \quad \text{RES} \frac{Q_1 \xrightarrow{\alpha}_2 Q_1' \quad z \notin n(\alpha)}{(\nu z)Q_1 \xrightarrow{\alpha}_2 (\nu z)Q_1'}}{(\nu y)P_1 \xrightarrow{\alpha}_2 (\nu z)Q_1'}$$

from this we create the following proof tree of  $P_1\{z/y\} \xrightarrow{\alpha}_2 Q_1'$ :

$$\text{ALP} \frac{P_1\{z/y\} \equiv_\alpha Q_1 \quad Q_1 \xrightarrow{\alpha}_2 Q_1'}{P_1\{z/y\} \xrightarrow{\alpha}_2 Q_1'}$$

to which we can apply the inductive hypothesis and get that there exists a process  $Q_1''$  such that

$$P_1\{z/y\} \xrightarrow{\alpha}_1 Q_1'' \text{ and } Q_1'' \equiv_\alpha Q_1'$$

then we apply the rule *Res* and *ResALP* to get

$$\text{RESALP} \frac{\text{RES} \frac{P_1\{z/y\} \xrightarrow{\alpha}_1 Q_1'' \quad z \notin n(\alpha)}{(\nu z)P_1\{z/y\} \xrightarrow{\alpha}_1 (\nu z)Q_1''}}{(\nu y)P_1 \xrightarrow{\alpha}_1 (\nu z)Q_1''}$$

this satisfies the thesis of the theorem because

$$(\nu z)Q_1'' \equiv (\nu z)Q_1'$$

**Alp** we can assume that there are no two consecutive application of the rule *Alp* because we can merge them thanks to the transitivity of the alpha equivalence.

**Opn** If  $S = \text{Opn}$  then there exists some names  $x, z$  and a process  $Q_1$  such that

$$Q = (\nu z)Q_1$$

and  $P' = Q'_1$  and  $\alpha = \bar{x}(z)$ . Since

$$P \equiv_\alpha (\nu z)Q_1$$

then for the inversion lemma we have two cases:

- there exists some  $P_1$  such that

$$P = (\nu z)P_1 \text{ and } P_1 \equiv_\alpha Q_1$$

and so the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  looks like this:

$$\text{ALP} \frac{(\nu z)P_1 \equiv_\alpha (\nu z)Q_1 \quad \text{OPN} \frac{Q_1 \xrightarrow{\bar{x}z}_2 Q'_1 \quad z \neq x}{(\nu z)Q_1 \xrightarrow{\bar{x}(z)}_2 Q'_1}}{(\nu z)P_1 \xrightarrow{\bar{x}(z)}_2 Q'_1}$$

from this we create the following proof tree of  $P_1 \xrightarrow{\bar{x}z}_2 Q'_1$ :

$$\text{ALP} \frac{P_1 \equiv_\alpha Q_1 \quad Q_1 \xrightarrow{\bar{x}z}_2 Q'_1}{P_1 \xrightarrow{\bar{x}z}_2 Q'_1}$$

to which we can apply the inductive hypothesis and get that there exists a process  $Q''_1$  such that

$$P_1 \xrightarrow{\bar{x}z}_1 Q''_1 \text{ and } Q''_1 \equiv_\alpha Q'_1$$

then we apply the rule *Opn* to get

$$\text{OPN} \frac{P_1 \xrightarrow{\bar{x}z}_1 Q''_1 \quad z \neq x}{(\nu z)P_1 \xrightarrow{\alpha}_1 Q''_1}$$

- there exists some  $P_1$  such that

$$P = (\nu y)P_1 \text{ and } P_1\{z/y\} \equiv_\alpha Q_1$$

and so the last part of the derivation tree of  $P \xrightarrow{\alpha}_2 P'$  looks like this:

$$\text{ALP} \frac{(\nu y)P_1 \equiv_\alpha (\nu z)Q_1 \quad \text{OPN} \frac{Q_1 \xrightarrow{\bar{x}z}_2 Q'_1 \quad z \neq x}{(\nu z)Q_1 \xrightarrow{\bar{x}z}_2 Q'_1}}{(\nu y)P_1 \xrightarrow{\alpha}_2 Q'_1}$$

from this we create the following proof tree of  $P_1\{z/y\} \xrightarrow{\alpha}_2 Q'_1$ :

$$\text{ALP} \frac{P_1\{z/y\} \equiv_\alpha Q_1 \quad Q_1 \xrightarrow{\alpha}_2 Q'_1}{P_1\{z/y\} \xrightarrow{\alpha}_2 Q'_1}$$

to which we can apply the inductive hypothesis and get that there exists a process  $Q''_1$  such that

$$P_1\{z/y\} \xrightarrow{\alpha}_1 Q''_1 \text{ and } Q''_1 \equiv_\alpha Q'_1$$

then we apply the rule *Opn* and *OpnAlp* to get

$$\text{OPNALP} \frac{\text{OPN} \frac{P_1\{z/y\} \xrightarrow{\bar{x}z}_1 Q''_1 \quad z \neq x}{(\nu z)P_1\{z/y\} \xrightarrow{\bar{x}(z)}_1 Q''_1} \quad z \notin n(P) \quad x \neq y \neq z}{(\nu y)P_1 \xrightarrow{\bar{x}(z)}_1 Q''_1}$$

**Cns** Since there is no process  $\alpha$  equivalent to an identifier except for the identifier itself, the last part of the derivation tree of  $P \xrightarrow{\alpha_2} P'$  looks like this:

$$\text{ALP} \frac{A(\tilde{x}|\tilde{y})\{\tilde{w}/\tilde{x}\} \equiv_{\alpha} A(\tilde{x}|\tilde{y})\{\tilde{w}/\tilde{x}\} \quad \text{CNS} \frac{A(\tilde{x}|\tilde{y}) \stackrel{\text{def}}{=} R \quad R\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha_2} P'}{A(\tilde{x}|\tilde{y})\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha_2} P'}}{A(\tilde{x}|\tilde{y})\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha_2} P'}$$

here we can apply the inductive hypothesis on the conclusion of  $S$  and get that there exists a process  $P''$  such that  $A(\tilde{x}|\tilde{y})\{\tilde{w}/\tilde{x}\} \xrightarrow{\alpha_1} P''$  and  $P' \equiv_{\alpha} P''$

□

**Theorem 2.5.3.**  $P \xrightarrow{\alpha_1} P' \Rightarrow P \xrightarrow{\alpha_2} P'$

*Proof.* The proof can go by induction on the length of the derivation of a transaction, and then both the base case and the inductive case proceed by cases on the last rule used in the derivation. However it is not necessary to show all the details of the proof because the rules in  $R_2$  are almost the same as the rules in  $R_1$ , the only difference is that in  $R_2$  we have the rule *Alp* instead of *ResAlp* and *OpnAlp*. The rule *Alp* can mimic the rule *ResAlp* in the following way:

$$\frac{(\nu z)P \equiv_{\alpha} (\nu w)P\{w/z\} \quad w \notin n(P) \quad (\nu w)P\{w/z\} \xrightarrow{xz} P'}{(\nu z)P \xrightarrow{xz} P'}$$

And the rule *Alp* can mimic the rule *OpnAlp* in the following way:

$$\frac{(\nu z)P \equiv_{\alpha} (\nu w)P\{w/z\} \quad w \notin n(P) \quad (\nu w)P\{w/z\} \xrightarrow{\bar{x}(w)} P' \quad x \neq w \neq z}{(\nu z)P \xrightarrow{\bar{x}(w)} P'}$$

□

**Theorem 2.5.4.**  $P \xrightarrow{\alpha_2} P' \Leftrightarrow \exists P'' : P' \equiv P'' \text{ and } P \xrightarrow{\alpha_3} P''$

*Proof.*  $\Rightarrow$  First we prove  $P \xrightarrow{\alpha_2} P' \Rightarrow \exists P'' : P' \equiv P'' \text{ and } P \xrightarrow{\alpha_3} P''$ . The proof is by induction on the length of the derivation of  $P \xrightarrow{\alpha_2} P'$ , and then both the base case and the inductive case proceed by cases on the last rule used.

**base case** in this case the rule used can be one of the following *Out*, *EInp*, *Tau* which are also in  $R_3$  so a derivation of  $P \xrightarrow{\alpha_2} P'$  is also a derivation of  $P \xrightarrow{\alpha_3} P'$

**inductive case :**

- the last rule used can be one in  $R_2 \cap R_3 = \{Res, Opn\}$  and so for example we have

$$\text{RES} \frac{P \xrightarrow{\alpha_2} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha_2} (\nu z)P'}$$

we apply the inductive hypothesis on  $P \xrightarrow{\alpha_2} P'$  and get  $\exists P''$  such that  $P' \equiv P''$  and  $P \xrightarrow{\alpha_3} P''$ . The proof we want is:

$$\text{RES} \frac{P \xrightarrow{\alpha_3} P'' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\alpha_3} (\nu z)P''}$$

and  $(\nu z)P'' \equiv (\nu z)P'$

- the last rule used can be one in  $\{ParL, ParR, SumL, SumR, EComL, EComR\}$ , in this case we can proceed as in the previous case and if necessary add an application of *Str* thus exploiting the commutativity of sum or parallel composition. For example

$$\text{PARR} \frac{Q \xrightarrow{\alpha_2} Q' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P|Q \xrightarrow{\alpha_2} P|Q'}$$



now we apply the inductive hypothesis to  $Q \xrightarrow{\alpha_2} Q'$  and get  $Q \xrightarrow{\alpha_3} Q''$  for a  $Q''$  such that  $Q' \equiv Q''$ . The proof we want is

$$\text{STR} \frac{P|Q \equiv Q|P \quad \text{PAR} \frac{Q \xrightarrow{\alpha_3} Q'' \quad bn(\alpha) \cap fn(Q) = \emptyset}{Q|P \xrightarrow{\alpha_3} Q''|P}}{P|Q \xrightarrow{\alpha_3} Q''|P}$$

and  $Q''|P \equiv P|Q'$

- if the last rule used is *Cns*:

$$\text{CNS} \frac{A(\tilde{x}|\tilde{z}) \stackrel{def}{=} P \quad P\{\tilde{y}/\tilde{x}\} \xrightarrow{\alpha_2} P'}{A(\tilde{y}|\tilde{z}) \xrightarrow{\alpha_2} P'}$$

we apply the inductive hypothesis on the premise and get  $P\{\tilde{y}/\tilde{x}\} \xrightarrow{\alpha_3} P''$  such that  $P' \equiv P''$ . Now the proof we want is

$$\text{STR} \frac{A(\tilde{y}|\tilde{z}) \equiv P\{\tilde{y}/\tilde{x}\} \quad P\{\tilde{y}/\tilde{x}\} \xrightarrow{\alpha_3} P''}{A(\tilde{y}|\tilde{z}) \xrightarrow{\alpha_3} P''}$$

- if the last rule is *Alp*, then we just notice that this rule is a particular case of *Str*
- if the last rule is *ClsL* (the case for *ClsR* is symmetric) then we have

$$\text{CLS L} \frac{P \xrightarrow{\bar{x}(z)}_2 P' \quad Q \xrightarrow{xz}_2 Q' \quad z \notin fn(Q)}{P|Q \xrightarrow{\tau}_2 (\nu z)(P'|Q')}$$

there is no easy way to mimic this rule with the rules in  $R_3$ . But if in the derivation tree we have an introduction of the bound output  $\bar{x}(z)$  followed directly by an elimination of the same bound output such as:

$$\text{CLS L} \frac{\text{OPN} \frac{P \xrightarrow{\bar{x}z}_2 P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)}_2 P'} \quad Q \xrightarrow{xz}_2 Q' \quad z \notin fn(Q)}{((\nu z)P)|Q \xrightarrow{\tau}_2 (\nu z)(P'|Q')}$$

we can apply the inductive hypothesis and get that

$$P \xrightarrow{\bar{x}z}_3 P'' \text{ and } Q \xrightarrow{xz}_3 Q''$$

where  $P' \equiv P''$  and  $Q' \equiv Q''$ , so we create the needed proof in the following way

$$\text{STR} \frac{(\nu z)(P|Q) \equiv ((\nu z)P)|Q \quad \text{COM} \frac{P \xrightarrow{\bar{x}z}_3 P'' \quad Q \xrightarrow{xz}_3 Q''}{P|Q \xrightarrow{\tau}_3 P''|Q''} \quad \text{RES} \frac{(\nu z)(P|Q) \xrightarrow{\tau}_3 (\nu z)(P''|Q'')}{(\nu z)(P|Q) \xrightarrow{\tau}_3 (\nu z)(P''|Q'')}}{((\nu z)P)|Q \xrightarrow{\tau}_3 (\nu z)(P''|Q'')}$$

We can always take a derivation tree in  $R_2$  and move downward each occurrence of *Opn* until we find the appropriate occurrence of *ClsL*. In this process we might need to use the structural congruence, in particular the scope extension axioms. We can attempt to prove that in the following way:

$$P \xrightarrow{\bar{x}(z)}_2 P' \Rightarrow \exists R : (\nu z)R \equiv P$$

and if  $(\nu z)R \xrightarrow{\bar{x}(z)}_2 P'$  then there exists a derivation tree for this transition such that the last rule used is *Opn*

PRIMA DEVO DIMOSTRARE IL LEMMA DI INVERSIONE PER LA CONGRUENZA STRUTTURALE(SE E' VERO)

Secondly we prove  $P \xrightarrow{\alpha}_3 P' \Rightarrow \exists P'' : P' \equiv P''$  and  $P \xrightarrow{\alpha}_2 P''$ . The proof is by induction on the length of the derivation of  $P \xrightarrow{\alpha}_3 P'$ , and then both the base case and the inductive case proceed by cases on the last rule used.

$\Leftarrow$  **base case** in this case the rule used can be one of the following *Out*, *EInp*, *Tau* which are also in  $R_2$  so a derivation of  $P \xrightarrow{\alpha}_3 P'$  is also a derivation of  $P \xrightarrow{\alpha}_2 P'$

**inductive case :**

- the last rule used can be one in  $R_2 \cap R_3 = \{Res, Opn\}$ , this goes like in the previous proof for the opposite direction with the transition numbers swapped.
- the last rule used can be one of *Par*, *Sum* or *ECom*, in this case we apply the inductive hypothesis to the premises and then apply the appropriate rule: *ParL*, *SumL* or *EComL*. For example

$$\text{PAR} \frac{P \xrightarrow{\alpha}_3 P' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P|Q \xrightarrow{\alpha}_3 P'|Q}$$

now we apply the inductive hypothesis to  $P \xrightarrow{\alpha}_3 P'$  and get  $P \xrightarrow{\alpha}_2 P''$  for a  $P''$  such that  $P' \equiv P''$ . The proof we want is

$$\text{PARL} \frac{P \xrightarrow{\alpha}_2 P'' \quad bn(\alpha) \cap fn(Q) = \emptyset}{P|Q \xrightarrow{\alpha}_2 P|Q''}$$

and  $Q''|P \equiv P|Q'$

- if the last rule is *Str*, then we have

$$\text{STR} \frac{P \equiv Q \quad Q \xrightarrow{\alpha}_3 P'}{P \xrightarrow{\alpha}_3 P'}$$

we proceed by cases on the premise  $Q \xrightarrow{\alpha}_3 P'$ . In the cases of prefix we can just use the appropriate prefix rule of  $R_2$  and get rid of the *Str*. In the other cases we can move upward the occurrence of *Str*, after that we have one or two smaller derivation trees that are suitable to application of the inductive hypothesis and finally we apply some appropriate rules in  $R_2$ .

**Out** Since we are using the rule *Out*,  $Q = \bar{x}y.Q_1$  for some  $Q_1$ .  $Q \equiv P$  means for the inversion lemma for structural congruence that  $P = \bar{x}y.P_1$  for some  $P_1 \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{\bar{x}y.P_1 \equiv \bar{x}y.Q_1 \quad \text{OUT} \frac{\bar{x}y.Q_1 \xrightarrow{\bar{x}y}_3 Q_1}{\bar{x}y.Q_1 \xrightarrow{\bar{x}y}_3 Q_1}}{\bar{x}y.P_1 \xrightarrow{\bar{x}y}_3 Q_1}$$

So we get

$$\text{OUT} \frac{\bar{x}y.P_1 \xrightarrow{\bar{x}y}_3 Q_1}{\bar{x}y.P_1 \xrightarrow{\bar{x}y}_2 P_1}$$

where  $P_1 \equiv Q_1$

**Tau** this is very similar to the previous case

**EInp** Since we are using the rule *EInp*,  $Q = x(y).Q_1$  for some  $Q_1$ . From  $Q \equiv P$  using the inversion lemma for structural congruence we can have two cases:

- $P = x(y).P_1$  for some  $P_1 \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{x(y).P_1 \equiv x(y).Q_1 \quad \text{EINP} \frac{x(y).Q_1 \xrightarrow{xw}_3 Q_1\{w/y\}}{x(y).Q_1 \xrightarrow{xw}_3 Q_1\{w/y\}}}{x(y).P_1 \xrightarrow{xw}_3 Q_1\{w/y\}}$$

So we get

$$\text{EINP} \frac{}{x(y).P_1 \xrightarrow{2} P_1\{w/y\}}$$

where  $P_1 \equiv Q_1$  implies  $P_1\{w/y\} \equiv Q_1\{w/y\}$

- $P = x(z).P_1$  for some  $P_1 \equiv Q_1\{z/y\}$ . The last part of the derivation tree is

$$\text{STR} \frac{x(z).P_1 \equiv x(y).Q_1 \quad \text{EINP} \frac{}{x(y).Q_1 \xrightarrow{3} Q_1\{w/y\}}}{x(z).P_1 \xrightarrow{3} Q_1\{w/y\}}$$

So we get

$$\text{EINP} \frac{}{x(z).P_1 \xrightarrow{2} P_1\{w/z\}}$$

where  $P_1 \equiv Q_1\{z/y\}$  implies  $P_1\{w/z\} \equiv Q_1\{z/y\}\{w/z\} \equiv Q_1\{w/y\}$

**Par** Since we are using the rule *Par*,  $Q = Q_1|Q_2$  for some  $Q_1, Q_2$ .  $Q \equiv P$  means for the inversion lemma for structural congruence that  $P = P_1|P_2$  for some  $P_1, P_2$  such that  $P_1 \equiv Q_1$  and  $P_2 \equiv Q_2$ . The last part of the derivation tree is

$$\text{STR} \frac{P_1|P_2 \equiv Q_1|Q_2 \quad \text{PAR} \frac{Q_1 \xrightarrow{3} Q'_1 \quad bn(\alpha) \cap fn(Q_2) = \emptyset}{Q_1|Q_2 \xrightarrow{3} Q'_1|Q_2}}{P_1|P_2 \xrightarrow{3} Q'_1|Q_2}$$

the first step is the creation of this proof tree:

$$\text{STR} \frac{P_1 \equiv Q_1 \quad Q_1 \xrightarrow{3} Q'_1}{P_1 \xrightarrow{3} Q'_1}$$

which is smaller then the inductive case, so we apply the inductive hypothesis and get  $P_1 \xrightarrow{2} Q'_1$  where  $Q'_1 \equiv Q''_1$ . The last step is

$$\text{PARL} \frac{P_1 \xrightarrow{2} Q''_1 \quad bn(\alpha) \cap fn(P_2) = \emptyset}{P_1|P_2 \xrightarrow{2} Q''_1|P_2}$$

**Sum** this case is very similar to the previous.

**ECom** this case is also similar to the *Par* case.

**Res** Since we are using the rule *Res*,  $Q = (\nu z)Q_1$  for some  $Q_1$  and some  $z$ .  $(\nu z)Q_1 \equiv P$  means thanks to the inversion lemma for structural congruence that one of the following cases holds:

- $P = (\nu z)P_1$  for some  $P_1$  such that  $P_1 \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{(\nu z)P_1 \equiv (\nu z)Q_1 \quad \text{RES} \frac{Q_1 \xrightarrow{3} Q'_1 \quad z \notin n(\alpha)}{(\nu z)Q_1 \xrightarrow{3} (\nu z)Q'_1}}{(\nu z)P_1 \xrightarrow{3} (\nu z)Q'_1}$$

first we create the following proof:

$$\text{STR} \frac{P_1 \equiv Q_1 \quad Q_1 \xrightarrow{3} Q'_1}{P_1 \xrightarrow{3} Q'_1}$$

now we can apply the inductive hypothesis and get  $P_1 \xrightarrow{2} Q'_1$  where  $Q'_1 \equiv Q''_1$ . The last step is

$$\text{RES} \frac{P_1 \xrightarrow{2} Q''_1 \quad z \notin n(\alpha)}{(\nu z)P_1 \xrightarrow{2} (\nu z)Q''_1}$$

- $P = (\nu y)P_1$  for some  $P_1$  such that  $P_1\{z/y\} \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{(\nu y)P_1 \equiv (\nu z)Q_1 \quad \text{RES} \frac{Q_1 \xrightarrow{\alpha}_3 Q'_1 \quad z \notin n(\alpha)}{(\nu z)Q_1 \xrightarrow{\alpha}_3 (\nu z)Q'_1}}{(\nu y)P_1 \xrightarrow{\alpha}_3 (\nu z)Q'_1}$$

we create the following proof of  $P_1\{z/y\} \xrightarrow{\alpha}_3 Q'_1$ :

$$\text{STR} \frac{P_1\{z/y\} \equiv Q_1 \quad Q_1 \xrightarrow{\alpha}_3 Q'_1}{P_1\{z/y\} \xrightarrow{\alpha}_3 Q'_1}$$

this proof tree is shorter then the one of  $(\nu y)P_1 \xrightarrow{\alpha}_3 (\nu z)Q'_1$  so we can apply the inductive hypothesis and get that there exists a process  $Q''_1$  such that

$$P_1\{z/y\} \xrightarrow{\alpha}_2 Q''_1 \text{ and } Q''_1 \equiv Q'_1$$

now we can apply the rules *Res* and *Alp* to get the desired proof tree:

$$\text{ALP} \frac{(\nu z)P_1\{z/y\} \equiv_\alpha (\nu y)P_1 \quad \text{RES} \frac{P_1\{z/y\} \xrightarrow{\alpha}_2 Q''_1 \quad z \notin (\alpha)}{(\nu z)P_1\{z/y\} \xrightarrow{\alpha}_2 (\nu z)Q''_1}}{(\nu y)P_1 \xrightarrow{\alpha}_2 (\nu z)Q''_1}$$

**Opn** Since we are using the rule *Opn*,  $Q = (\nu z)Q_1$  for some  $Q_1$ .  $(\nu z)Q_1 \equiv P$  means for the inversion lemma for structural congruence that

- $P = (\nu z)P_1$  for some  $P_1$  such that  $P_1 \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{(\nu z)P_1 \equiv (\nu z)Q_1 \quad \text{OPN} \frac{Q_1 \xrightarrow{\bar{x}z} Q'_1 \quad z \neq x}{(\nu z)Q_1 \xrightarrow{\bar{x}(z)} Q'_1}}{(\nu z)P_1 \xrightarrow{\bar{x}(z)} Q'_1}$$

first:

$$\text{STR} \frac{P_1 \equiv Q_1 \quad Q_1 \xrightarrow{\bar{x}z}_3 Q'_1}{P_1 \xrightarrow{\bar{x}z}_3 Q'_1}$$

then we apply the inductive hypothesis and get  $P_1 \xrightarrow{\bar{x}z}_2 Q''_1$  where  $Q'_1 \equiv Q''_1$ . The last step is

$$\text{RES} \frac{P_1 \xrightarrow{\bar{x}z}_2 Q''_1 \quad z \neq x}{(\nu z)P_1 \xrightarrow{\bar{x}z}_2 Q''_1}$$

- $P = (\nu z)P_1$  for some  $P_1$  such that  $P_1 \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{(\nu z)P_1 \equiv (\nu z)Q_1 \quad \text{OPN} \frac{Q_1 \xrightarrow{\bar{x}z} Q'_1 \quad z \neq x}{(\nu z)Q_1 \xrightarrow{\bar{x}(z)} Q'_1}}{(\nu z)P_1 \xrightarrow{\bar{x}(z)} Q'_1}$$

the first step is:

$$\text{STR} \frac{P_1 \equiv Q_1 \quad Q_1 \xrightarrow{\bar{x}z}_3 Q'_1}{P_1 \xrightarrow{\bar{x}z}_3 Q'_1}$$

then we apply the inductive hypothesis and get  $P_1 \xrightarrow{\bar{x}z}_2 Q''_1$  where  $Q'_1 \equiv Q''_1$ . The last step is

$$\text{RES} \frac{P_1 \xrightarrow{\bar{x}z}_2 Q''_1 \quad z \neq x}{(\nu z)P_1 \xrightarrow{\bar{x}z}_2 Q''_1}$$

- $P = (\nu y)P_1$  for some  $P_1$  such that  $P_1\{z/y\} \equiv Q_1$ . The last part of the derivation tree is

$$\text{STR} \frac{(\nu y)P_1 \equiv (\nu z)Q_1 \quad \text{OPN} \frac{Q_1 \xrightarrow{\bar{x}z} Q'_1 \quad z \neq x}{(\nu z)Q_1 \xrightarrow{\bar{x}(z)} Q'_1}}{(\nu y)P_1 \xrightarrow{\bar{x}(z)} Q'_1}$$

we can create the following proof of  $P_1\{z/y\} \xrightarrow{\bar{x}z} Q'_1$ :

$$\text{STR} \frac{P_1\{z/y\} \equiv Q_1 \quad Q_1 \xrightarrow{\bar{x}z} Q'_1}{P_1\{z/y\} \xrightarrow{\bar{x}z} Q'_1}$$

this proof tree is shorter then the one of  $(\nu y)P_1 \xrightarrow{\bar{x}(z)} Q'_1$  so we can apply the inductive hypothesis and get that there exists a process  $Q''_1$  such that

$$Q''_1 \equiv Q'_1 \text{ and } P_1\{z/y\} \xrightarrow{\bar{x}z} Q''_1$$

so now we only need to apply the rules *Opn* and *Alp*:

$$\text{ALP} \frac{(\nu y)P_1 \equiv_\alpha (\nu z)P_1\{z/y\} \quad \text{OPN} \frac{P_1\{z/y\} \xrightarrow{\bar{x}z} Q''_1 \quad z \neq x}{(\nu z)P_1\{z/y\} \xrightarrow{\bar{x}(z)} Q''_1}}{(\nu y)P_1 \xrightarrow{\bar{x}(z)} Q''_1}$$

□

### 2.5.2 Equivalence of the late semantics

## 2.6 Bisimilarity, congruence and equivalence

We present here some behavioural equivalences and some of their properties. In the following we will use the phrase *bn*( $\alpha$ ) is fresh in a definition to mean that the name in *bn*( $\alpha$ ), if any, is different from any free name occurring in any of the agents in the definition. We write  $\rightarrow_E$  for the early semantic and  $\rightarrow_L$  for the late semantic. It's not a concern which late semantic we are talking about because we have proved them equivalent.

### 2.6.1 Late bisimilarity

**Definition 2.6.1.** A *strong late bisimulation* (according to [4]) is a binary symmetric relation **S** on processes such that for each process  $P$  and  $Q$ ,  $PSQ$  implies:

- if  $P \xrightarrow{a(x)}_L P'$  and  $x \notin \text{fn}(P) \cup \text{fn}(Q)$  then there exists a process  $Q'$  such that  $Q \xrightarrow{a(x)}_L Q'$  and for all  $u$   $P'\{u/x\} \mathbf{S} Q'\{u/x\}$
- if  $P \xrightarrow{\alpha}_L P'$ ,  $\alpha$  is not an input and  $\text{bn}(\alpha) \cap (\text{fn}(P) \cup \text{fn}(Q)) = \emptyset$  then there exists a process  $Q'$  such that  $Q \xrightarrow{\alpha}_L Q'$  and  $P' \mathbf{S} Q'$

$P$  and  $Q$  are *late bisimilar* written  $P \sim_L Q$  if there exists a strong late bisimulation **S** such that  $PSQ$ .

**Example** Strong late bisimulation is not closed under substitution in general:

$$a(u).0 | \bar{b}v.0 \sim_L a(u).\bar{b}v.0 + \bar{b}v.a(u).0$$

and the bisimulation (without the symmetric part) is the following:

$$\{(a(u).0 | \bar{b}v.0, a(u).\bar{b}v.0 + \bar{b}v.a(u).0), (a(u).0 | 0, a(u).0), (0 | 0, 0), (0 | \bar{b}v.0, \bar{b}v.0)\}$$

If we apply the substitution  $\{a/b\}$  to each process then they are not strongly bisimilar anymore because  $(a(u).0|\bar{b}v.0)\{a/b\}$  is  $a(u).0|\bar{a}v.0$  and this process can perform an invisible action whether  $(a(u).\bar{b}v.0 + \bar{b}v.a(u).0)\{a/b\}$  cannot.

We refer to strong late bisimulation as strong *ground* late bisimulation, because it is not preserved by substitution.

**Proposition 2.6.1.** If  $P \sim Q$  and  $\sigma$  is injective then  $P\sigma \sim Q\sigma$

**Proposition 2.6.2.**  $\sim_L$  is an equivalence

**Proposition 2.6.3.**  $\sim_L$  is preserved by all operators except input prefix

**Definition 2.6.2.** Two processes  $P$  and  $Q$  are *strong late equivalent* written  $P \sim_L Q$  if for each substitution  $\sigma$   $P\sigma \sim_L Q\sigma$

**Example** If  $z \notin fn(R) \cup \{x\}$  then  $x(y).R \sim_L (z)x(y).R$

## 2.6.2 Early bisimilarity

**Definition 2.6.3.** A *strong early bisimulation* (according to [4]) is a symmetric binary relation  $\mathbf{S}$  on processes such that for each process  $P$  and  $Q$ :  $PSQ$ ,  $P \xrightarrow{\alpha}_E P'$  and  $bn(\alpha) \cap (fn(P) \cup fn(Q)) = \emptyset$  implies that there exists  $Q'$  such that  $Q \xrightarrow{\alpha}_E Q'$  and  $P'SQ'$ .  $P$  and  $Q$  are *early bisimilar* written  $P \sim_E Q$  if there exists a strong early bisimulation  $\mathbf{S}$  such that  $PSQ$

**Definition 2.6.4.** Two processes  $P$  and  $Q$  are *strong early equivalent* written  $P \sim_E Q$  if for each substitution  $\sigma$   $P\sigma \sim_E Q\sigma$

## 2.6.3 Congruence

**Definition 2.6.5.** We say that two agents  $P$  and  $Q$  are *strongly congruent*, written  $P \sim Q$  if

$$P\sigma \sim Q\sigma \text{ for all substitution } \sigma$$

**Proposition 2.6.4.** Strong congruence is the largest congruence in bisimilarity.

## 2.6.4 Open bisimilarity

**Definition 2.6.6.** A *distinction* is a finite symmetric and irreflexive binary relation on names. A substitution  $\sigma$  *respects* a distinction  $D$  if for each name  $a, b$   $aDb$  implies  $\sigma(a) \neq \sigma(b)$ . We write  $D\sigma$  for the composition of the two relation.

**Definition 2.6.7.** An *strong open simulation* (according to [4]) is  $\{S_D\}_{D \in \mathbb{D}}$  a family of binary relations on processes such that for each process  $P, Q$ , for each distinction  $D \in \mathbb{D}$ , for each name substitution  $\sigma$  which respects  $D$  if  $PS_DQ$ ,  $P\sigma \xrightarrow{\alpha} P'$  and  $bn(\alpha) \cap (fn(P\sigma) \cup fn(Q\sigma)) = \emptyset$  then:

- if  $\alpha = \bar{a}(x)$  then there exists  $Q'$  such that  $Q\sigma \xrightarrow{\bar{a}(x)} Q'$  and  $P'S_{D'}Q'$  where  $D' = D\sigma \cup \{x\} \times (fn(P\sigma) \cup fn(Q\sigma)) \cup (fn(P\sigma) \cup fn(Q\sigma)) \times \{x\}$
- if  $\alpha$  is not a bound output then there exists  $Q'$  such that  $Q\sigma \xrightarrow{\alpha} Q'$  and  $P'S_{D\sigma}Q'$

$P$  and  $Q$  are *open  $D$  bisimilar*, written  $P \sim_O^D Q$  if there exists a member  $S_D$  of an open bisimulation such that  $PS_DQ$ ; they are *open bisimilar* if they are open  $\emptyset$  bisimilar, written  $P \sim_O D$ .

## Chapter 3

# Multi $\pi$ calculus with strong output

### 3.1 Syntax

As we did with  $\pi$  calculus, we suppose that we have a countable set of names  $\mathbb{N}$ , ranged over by lower case letters  $a, b, \dots, z$ . This names are used for communication channels and values. Furthermore we have a set of identifiers, ranged over by  $A$ . We represent the agents or processes by upper case letters  $P, Q, \dots$ . A multi  $\pi$  process, in addition to the same actions of a  $\pi$  process, can perform also a strong prefix output:

$$\pi ::= \bar{x}y \mid x(z) \mid \bar{x}y \mid \tau$$

The process are defined, just as original  $\pi$  calculus, by the following grammar:

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A(y_1, \dots, y_n)$$

and they have the same intuitive meaning as for the  $\pi$  calculus. The strong prefix output allows a process to make an atomic sequence of actions, so that more than one process can synchronize on this sequence. For the moment we allow the strong prefix to be on output names only. Also one can use the strong prefix only as an action prefixing for processes that can make at least a further action.

Multi  $\pi$  calculus is a conservative extension of the  $\pi$  calculus in the sense that: any  $\pi$  calculus process  $p$  is also a multi  $\pi$  calculus process and the semantic of  $p$  according to the SOS rules of  $\pi$  calculus is the same as the semantic of  $p$  according to the SOS rules of multi  $\pi$  calculus.

We have to extend the following definition to deal with the strong prefix:

$$B(\bar{x}y.Q, I) = B(Q, I) \quad F(\bar{x}y.Q, I) = \{x, \bar{x}, y, \bar{y}\} \cup F(Q, I)$$

### 3.2 Operational semantic

#### 3.2.1 Early operational semantic with structural congruence

The semantic of a multi  $\pi$  process is labeled transition system such that

- the nodes are multi  $\pi$  calculus process. The set of node is  $\mathbb{P}_m$
- the actions are multi  $\pi$  calculus actions. The set of actions is  $\mathbb{A}_m$ , we use  $\alpha, \alpha_1, \alpha_2, \dots$  to range over the set of actions, we use  $\sigma, \sigma_1, \sigma_2, \dots$  to range over the set  $\mathbb{A}_m^+ \cup \{\tau\}$ . Note that  $\sigma$  is a non empty sequence of actions.
- the transition relations is  $\rightarrow \subseteq \mathbb{P}_m \times (\mathbb{A}_m^+ \cup \{\tau\}) \times \mathbb{P}_m$

In this case, a label can be a sequence of prefixes, whether in the original  $\pi$  calculus a label can be only a prefix. We use the symbol  $\cdot$  to denote the concatenation operator.

**Definition 3.2.1.** The *early transition relation* is the smallest relation induced by the rules in table 3.1.

---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(y).P \xrightarrow{xz} P\{z/y\}}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$
<b>SOutSeq</b> $\frac{P \xrightarrow{\sigma} Q \quad  \sigma  > 1}{\bar{x}y.P \xrightarrow{\bar{x}y \cdot \sigma} Q}$	<b>SOut</b> $\frac{P \xrightarrow{\alpha} Q \quad \alpha \text{ output}}{\bar{x}y.P \xrightarrow{\bar{x}y \cdot \alpha} Q}$	<b>SOutTau</b> $\frac{P \xrightarrow{\tau} Q}{\bar{x}y.P \xrightarrow{\bar{x}y} Q}$
<b>EComSng</b> $\frac{P \xrightarrow{\bar{x}y} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>EComSeq</b> $\frac{P \xrightarrow{\bar{x}y \cdot \sigma} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\sigma} P' Q'}$	
<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\sigma} Q}{P \xrightarrow{\sigma} Q}$	<b>Par</b> $\frac{P \xrightarrow{\sigma} P'}{P Q \xrightarrow{\sigma} P' Q}$	
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\alpha)}{(\nu)zP \xrightarrow{\sigma} (\nu)zP'}$	

---

Table 3.1: Multi  $\pi$  early semantic with structural congruence

**Lemma 3.2.1.** If  $P \xrightarrow{\sigma} Q$  then only one of the following cases hold:

- $|\sigma| = 1$
- $|\sigma| > 1$  and all the actions are output.

**Example** Multi-party synchronization. We show an example of a derivation of three processes that synchronize.

$$\begin{array}{c}
 \text{Res} \frac{x \notin n(\tau) \quad \text{EComSeq} \frac{\bar{x}y.\bar{x}y.0|x(y).0 \xrightarrow{\bar{x}y} 0|0 \quad \text{Inp} \frac{}{x(y).0 \xrightarrow{xy} 0}}{((\bar{x}y.\bar{x}y.0|x(y).0)|x(y).0) \xrightarrow{\tau} ((0|0)|0)}}{(\nu x)((\bar{x}y.\bar{x}y.0|x(y).0)|x(y).0) \xrightarrow{\tau} (\nu x)((0|0)|0)} \\
 \\
 \text{EComSng} \frac{\text{SOut} \frac{\text{Out} \frac{}{\bar{x}y.0 \xrightarrow{\bar{x}y} 0}}{\bar{x}y.\bar{x}y.0 \xrightarrow{\bar{x}y \cdot \bar{x}y} 0} \quad x(y).0 \xrightarrow{xy} 0}{\bar{x}y.\bar{x}y.0|x(y).0 \xrightarrow{\bar{x}y} 0|0}}
 \end{array}$$

**Example** Transactional synchronization In this setting two process cannot synchronize on a sequence of actions with length greater than one. This is because of the rules *EComSng* and *EComSeq*.

### 3.2.2 Low level semantic

This section contains the definition of an alternative semantic for multi  $\pi$ . First we define a low level version of the multi  $\pi$  calculus (here with strong prefixing on output only), we call this language low multi  $\pi$ . The low multi  $\pi$  is the multi  $\pi$  enriched with a marked or intermediate process  $*P$ :

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A \mid *P$$

$$\pi ::= \bar{x}y \mid x(y) \mid \bar{x}y \mid \tau$$

**Definition 3.2.2.** The low level transition relation is the smallest relation induced by the rules in table 3.2 in which  $P$  stands for a process without mark,  $L$  stands for a process with mark and  $S$  can stand for both.



---

<b>Out</b> $\frac{}{\bar{x}y.P \mapsto P}$	<b>EInp</b> $\frac{}{x(y).P \mapsto P\{z/y\}}$	<b>Tau</b> $\frac{}{\tau.P \mapsto P}$
<b>SOutLow</b> $\frac{}{\bar{x}y.P \mapsto *P}$	<b>StarEps</b> $\frac{S \mapsto S'}{*S \mapsto S'}$	<b>StarOut</b> $\frac{S \mapsto S'}{*S \mapsto S'}$
<b>Com1</b> $\frac{P \mapsto P' \quad Q \mapsto Q'}{P Q \mapsto P' Q'}$		
<b>Com2L</b> $\frac{L_1 \mapsto L'_1 \quad P \mapsto Q}{L_1 P \mapsto L'_1 Q}$	<b>Com2R</b> $\frac{P \mapsto Q \quad L_1 \mapsto L'_1}{P L_1 \mapsto Q L'_1}$	
<b>Com3L</b> $\frac{P \mapsto L \quad Q \mapsto Q'}{P Q \mapsto L Q'}$	<b>Com3R</b> $\frac{P \mapsto P' \quad Q \mapsto L}{P Q \mapsto P' L}$	
<b>Com4L</b> $\frac{L \mapsto Q \quad P \mapsto P'}{L P \mapsto Q P'}$	<b>Com4R</b> $\frac{P \mapsto P' \quad L \mapsto Q}{P L \mapsto P' Q}$	
<b>Res</b> $\frac{S \mapsto S' \quad y \notin n(\gamma)}{(\nu y)S \mapsto (\nu y)S'}$	<b>Sum</b> $\frac{P \mapsto S}{P + Q \mapsto S}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \mapsto S}{P \mapsto S}$
<b>Par1L</b> $\frac{S \mapsto S'}{S Q \mapsto S' Q}$	<b>Par1R</b> $\frac{S \mapsto S'}{Q S \mapsto Q S'}$	

---

Table 3.2: Low multi  $\pi$  early semantic with structural congruence

**Lemma 3.2.2.** For all unmarked processes  $P, Q$  and marked processes  $L, L_1, L_2$ .

- if  $L_1 \xrightarrow{\alpha} L_2$  or  $P \xrightarrow{\alpha} L$  then  $\alpha$  can only be an output or an  $\epsilon$
- if  $L \xrightarrow{\alpha} P$  then  $\alpha$  can only be an output or a  $\tau$
- if  $P \xrightarrow{\alpha} Q$  then  $\alpha$  cannot be an  $\epsilon$

**Definition 3.2.3.** Let  $P, Q$  be unmarked processes and  $L_1, \dots, L_{k-1}$  marked processes. We define the derivation relation  $\rightarrow_s$  in the following way:

$$\text{Low} \frac{P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} Q \quad k \geq 1}{P \xrightarrow{\gamma_1 \cdots \gamma_k}_s Q}$$

We need to be precise about the concatenation operator  $\cdot$  since we have introduced the new label  $\epsilon$ . Let  $a$  be an action such that  $a \neq \tau$  and  $a \neq \epsilon$  then the following rules hold:

$$\begin{aligned} \epsilon \cdot a &= a \cdot \epsilon = a & \epsilon \cdot \epsilon &= \epsilon & \tau \cdot \epsilon &= \epsilon \cdot \tau = \tau \\ \tau \cdot a &= a \cdot \tau = a & \tau \cdot \tau &= \tau \end{aligned}$$

**Example** Multi-parti synchronization

$$\begin{array}{c} \text{SOutLow} \frac{}{\bar{x}a.\bar{x}b.\bar{x}c.P \xrightarrow{\bar{x}a} *\bar{x}b.\bar{x}c.P} \quad \text{Inp} \frac{}{x(d).Q \xrightarrow{xa} Q\{a/d\}} \\ \text{Com3L} \frac{}{\bar{x}a.\bar{x}b.\bar{x}c.P|x(d).Q \xrightarrow{\epsilon} *\bar{x}b.\bar{x}c.P|Q\{a/d\}} \\ \text{Par1L} \frac{}{\bar{x}a.\bar{x}b.\bar{x}c.P|x(d).Q|x(e).R \xrightarrow{\epsilon} *\bar{x}b.\bar{x}c.P|Q\{a/d\}|x(e).R} \\ \text{Par1L} \frac{}{\bar{x}a.\bar{x}b.\bar{x}c.P|x(d).Q|x(e).R|x(f).S \xrightarrow{\epsilon} *\bar{x}b.\bar{x}c.P|Q\{a/d\}|x(e).R|x(f).S} \\ \\ \text{SOutLow} \frac{}{\bar{x}b.\bar{x}c.P \xrightarrow{\bar{x}b} *\bar{x}c.P} \\ \text{StarOut} \frac{}{*\bar{x}b.\bar{x}c.P \xrightarrow{\bar{x}b} *\bar{x}c.P} \\ \text{Par1L} \frac{}{*\bar{x}b.\bar{x}c.P|Q\{a/d\} \xrightarrow{\bar{x}b} *\bar{x}c.P|Q\{a/d\}} \quad \text{EInp} \frac{}{x(e).R \xrightarrow{xb} R\{b/e\}} \\ \text{Com2L} \frac{}{*\bar{x}b.\bar{x}c.P|Q\{a/d\}|x(e).R \xrightarrow{\epsilon} *\bar{x}c.P|Q\{a/d\}|R\{b/e\}} \\ \text{Par1L} \frac{}{*\bar{x}b.\bar{x}c.P|Q\{a/d\}|x(e).R|x(f).S \xrightarrow{\epsilon} *\bar{x}c.P|Q\{a/d\}|R\{b/e\}|x(f).S} \\ \\ \text{Out} \frac{}{\bar{x}c.P \xrightarrow{\bar{x}c} P} \\ \text{StarOut} \frac{}{*\bar{x}c.P \xrightarrow{\bar{x}c} P} \\ \text{Par1L} \frac{}{*\bar{x}c.P|Q\{a/d\} \xrightarrow{\bar{x}c} P|Q\{a/d\}} \\ \text{Par1L} \frac{}{*\bar{x}c.P|Q\{a/d\}|R\{b/e\} \xrightarrow{\bar{x}c} P|Q\{a/d\}|R\{b/e\}} \quad \text{EInp} \frac{}{x(f).S \xrightarrow{xc} R\{c/f\}} \\ \text{Com4L} \frac{}{*\bar{x}c.P|Q\{a/d\}|R\{b/e\}|x(f).S \xrightarrow{\tau} P|Q\{a/d\}|R\{b/e\}|S\{c/f\}} \end{array}$$

**Proposition 3.2.3.** Let  $\rightarrow$  be the relation defined in table 3.1. If  $P \xrightarrow{\sigma} Q$  then there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

*Proof.* The proof is by induction on the depth of the derivation tree of  $P \xrightarrow{\sigma} Q$ :

**base case**

If the depth is one then the rule used has to be one of: *EInp*, *Out*, *Tau*. These rules are also in table 3.2 so we can derive  $P \xrightarrow{\sigma} Q$ .

**inductive case**

If the depth is greater than one then the last rule used in the derivation can be:

*SOutSeq* : the last part of the derivation tree looks like this:

$$\mathbf{SOutSeq} \frac{P_1 \xrightarrow{\sigma} Q \quad |\sigma| > 1}{\overline{xy}.P_1 \xrightarrow{\overline{xy}.\sigma} Q}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \dots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \dots \gamma_{k+1} = \sigma$$

then a proof of the conclusion follows from:

$$\mathbf{SOutLow} \frac{}{\overline{xy}.P_1 \xrightarrow{\overline{xy}} *P_1} \quad \mathbf{Star} \frac{P_1 \xrightarrow{\gamma_1} L_1}{*P_1 \xrightarrow{\gamma_1} L_1}$$

*SOut* : this case is similar to the previous.

*SOutTau* : this case is similar to the previous observing that  $\overline{xy} \cdot \tau = \overline{xy}$ .

*Sum* : the last part of the derivation tree looks like this:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\sigma} Q}{P_1 + P_2 \xrightarrow{\sigma} Q}$$

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \dots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \dots \gamma_{k+1} = \sigma$$

A proof of the conclusion is:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\gamma_1} L_1}{P_1 + P_2 \xrightarrow{\gamma_1} L_1}$$

*Cong* : this case is similar to the previous.

*EComSng* : the last part of the derivation tree looks like this:

$$\mathbf{Com} \frac{P_1 \xrightarrow{\overline{xy}} P'_1 \quad Q_1 \xrightarrow{xy} Q'_1}{P_1|Q_1 \xrightarrow{\tau} P'_1|Q'_1}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \dots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} P'_1 \quad \text{and} \quad \gamma_1 \dots \gamma_{k+1} = \overline{xy}$$

and there exist  $R_1, \dots, R_h$  and  $\delta_1, \dots, \delta_{h+1}$  with  $h \geq 0$  such that

$$Q_1 \xrightarrow{\delta_1} R_1 \xrightarrow{\delta_2} R_2 \dots R_{h-1} \xrightarrow{\delta_h} R_h \xrightarrow{\delta_{h+1}} Q'_1 \quad \text{and} \quad \delta_1 \dots \delta_{h+1} = xy$$

For lemma 3.2.2 there cannot be an input action in a transition involving marked processes so  $h$  must be 0 and  $Q_1 \xrightarrow{\delta_1} Q'_1$  with  $\delta_1 = xy$ . Just one of the  $\gamma$ s is  $\overline{xy}$  and the others are  $\epsilon$  or  $\tau$ . We can have three different cases now:

$\gamma_1 = \overline{xy}$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\tau} L_1|Q'_1 \xrightarrow{\epsilon} L_2|Q'_1 \dots \xrightarrow{\epsilon} L_k|Q'_1 \xrightarrow{\tau} P'_1|Q'_1$$

we derive the first transition with rule *Com3L*, whether for the other transition we use the rule *Par1L*.

$\gamma_i = \bar{x}y$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1 \xrightarrow{\epsilon} L_{i+1}|Q'_1 \cdots \xrightarrow{\epsilon} L_k|Q'_1 \xrightarrow{\tau} P'_1|Q'_1$$

we derive the transition  $L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1$  with rule *Com2L*, whether for the other transitions we use the rule *Par1L*.

$\gamma_{k+1} = \bar{x}y$  similar.

*Res* : the last part of the derivation tree looks like this:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\sigma} Q_1 \quad z \notin n(\sigma)}{(\nu z)P_1 \xrightarrow{\sigma} (\nu z)Q_1}$$

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q_1 \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \sigma$$

We can apply the rule *Res* to each of the previous transitions because

$$z \notin n(\sigma) \text{ implies } z \notin n(\gamma_i) \text{ for each } i$$

and then get a proof of the conclusion:

$$(\nu z)P_1 \xrightarrow{\gamma_1} (\nu z)L_1 \xrightarrow{\gamma_2} (\nu z)L_2 \cdots (\nu z)L_{k-1} \xrightarrow{\gamma_k} (\nu z)L_k \xrightarrow{\gamma_{k+1}} (\nu z)Q_1$$

*Par* : this case is similar to the previous.

*EComSeq* : the last part of the derivation tree looks like this:

$$\mathbf{EComSeq} \frac{P_1 \xrightarrow{\bar{x}y \cdot \sigma} P'_1 \quad Q_1 \xrightarrow{xy} Q'_1}{P_1|Q_1 \xrightarrow{\sigma} P'_1|Q'_1}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} P'_1 \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \bar{x}y \cdot \sigma$$

For inductive hypothesis and lemma 3.2.2  $Q_1 \xrightarrow{xy} Q'_1$ . We can have two different cases now depending on where the first  $\bar{x}y$  is:

$\gamma_1 = \bar{x}y$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q'_1 \xrightarrow{\gamma_2} L_2|Q'_1 \cdots \xrightarrow{\gamma_k} L_k|Q'_1 \xrightarrow{\gamma_{k+1}} P'_1|Q'_1$$

we derive the first transition with rule *Com3L*, whether for the other transitions we use the rule *Par1L*. Since  $\gamma_1 \cdot \dots \cdot \gamma_{k+1} = \bar{x}y \cdot \sigma$  and  $\gamma_1 = \bar{x}y$  then  $\epsilon \cdot \gamma_2 \cdot \dots \cdot \gamma_{k+1} = \sigma$

$\gamma_i = \bar{x}y$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1 \xrightarrow{\gamma_{i+1}} L_{i+1}|Q'_1 \cdots \xrightarrow{\gamma_k} L_k|Q'_1 \xrightarrow{\gamma_{k+1}} P'_1|Q'_1$$

we derive the transition  $L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1$  with rule *Com2L*, whether for the other transitions of the premises we use the rule *Par1L*.

$\gamma_{k+1} = \bar{x}y$  : cannot happen because  $\sigma$  is not empty.

□

We would like to prove the converse of the previous proposition, namely: if there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

then  $P \xrightarrow{\sigma} Q$ . But this is false as shown by those examples:

**Example** Interleaving

$$\begin{array}{c}
 \text{SOutLow} \frac{}{\overline{xy}. \underline{ab}. \overline{xy}. 0 \xrightarrow{\overline{xy}} * \underline{ab}. \overline{xy}. 0} \quad \text{EInp} \frac{}{x(y). 0 \xrightarrow{xy} 0} \\
 \text{Com3L} \frac{}{\overline{xy}. \underline{ab}. \overline{xy}. 0 | x(y). 0 \xrightarrow{\epsilon} * \underline{ab}. \overline{xy}. 0 | 0} \\
 \text{Par1L} \frac{}{\overline{xy}. \underline{ab}. \overline{xy}. 0 | x(y). 0 | x(y). 0 \xrightarrow{\epsilon} * \underline{ab}. \overline{xy}. 0 | 0 | x(y). 0} \\
 \\
 \text{SOutLow} \frac{}{\underline{ab}. \overline{xy}. 0 \xrightarrow{\overline{ab}} * \overline{xy}. 0} \\
 \text{StarOut} \frac{}{* \underline{ab}. \overline{xy}. 0 \xrightarrow{\overline{ab}} * \overline{xy}. 0} \\
 \text{Par1L} \frac{}{* \underline{ab}. \overline{xy}. 0 | 0 \xrightarrow{\overline{ab}} * \overline{xy}. 0 | 0} \\
 \text{Par1L} \frac{}{* \underline{ab}. \overline{xy}. 0 | 0 | x(y). 0 \xrightarrow{\overline{ab}} * \overline{xy}. 0 | 0 | x(y). 0} \\
 \\
 \text{Out} \frac{}{\overline{xy}. 0 \xrightarrow{\overline{xy}} 0} \\
 \text{StarOut} \frac{}{* \overline{xy}. 0 \xrightarrow{\overline{xy}} 0} \\
 \text{Par1L} \frac{}{* \overline{xy}. 0 | 0 \xrightarrow{\overline{xy}} 0 | 0} \quad \text{EInp} \frac{}{x(y). 0 \xrightarrow{xy} 0} \\
 \text{Com4L} \frac{}{* \overline{xy}. 0 | 0 | x(y). 0 \xrightarrow{\tau} 0 | 0 | 0}
 \end{array}$$

this prove:

$$\overline{xy}. \underline{ab}. \overline{xy}. 0 | x(y). 0 | x(y). 0 \xrightarrow{\epsilon} * \underline{ab}. \overline{xy}. 0 | 0 | x(y). 0 \xrightarrow{\overline{ab}} * \overline{xy}. 0 | 0 | x(y). 0 \xrightarrow{\tau} 0 | 0 | 0$$

but there is no way to prove

$$\overline{xy}. \underline{ab}. \overline{xy}. 0 | x(y). 0 | x(y). 0 \xrightarrow{\overline{ab}} 0 | 0 | 0$$

**Example** Transactional synchronization

$$\begin{array}{c}
 \text{SOutLow} \frac{}{\overline{xy}. \overline{xy}. 0 \xrightarrow{\overline{xy}} * \overline{xy}. 0} \quad \text{EInp} \frac{}{x(y). x(y). 0 \xrightarrow{xy} x(y). 0} \\
 \text{Com3L} \frac{}{\overline{xy}. \overline{xy}. 0 | x(y). x(y). 0 \xrightarrow{\epsilon} * \overline{xy}. 0 | x(y). 0} \\
 \\
 \text{Out} \frac{}{\overline{xy}. 0 \xrightarrow{\overline{xy}} 0} \\
 \text{StarOut} \frac{}{* \overline{xy}. 0 \xrightarrow{\overline{xy}} 0} \quad \text{EInp} \frac{}{x(y). 0 \xrightarrow{xy} 0} \\
 \text{Com4L} \frac{}{* \overline{xy}. 0 | x(y). 0 \xrightarrow{\tau} 0 | 0}
 \end{array}$$

this prove:

$$\overline{xy}. \overline{xy}. 0 | x(y). x(y). 0 \xrightarrow{\epsilon} * \overline{xy}. 0 | x(y). 0 \xrightarrow{\tau} 0 | 0$$

but we cannot derive

$$\overline{xy}. \overline{xy}. 0 | x(y). x(y). 0 \xrightarrow{\tau} 0 | 0$$

also we do not want to derive this transaction because the second process does not start with a strong prefix.

There is a much weaker propositions we can prove:

**Proposition 3.2.4.** Let  $\rightarrow$  be the relation defined in table 3.1. Let  $\alpha$  be an action. If  $P \vdash^\alpha Q$  then  $P \xrightarrow{\alpha} Q$ .

*Proof.* The proof is by induction the depth of the derivation of  $P \vdash^\alpha Q$ :

**base case** in this case the derivation of this transition has depth one. The last(and only) rule used can be: *Out*, *EInp* or *Tau*; these rules are also in table 4.1 so we can derive  $P \xrightarrow{\alpha} Q$ .

**inductive case** in this case the last rule in the derivation can be: *Sum*, *Com1*, *Res*, *Par1L*, *Par1R*, *Cong*:

*Com1* :

$$\mathbf{Com1} \frac{P_1 \xrightarrow{\bar{x}y} Q_1 \quad P_2 \xrightarrow{xy} Q_2}{P_1|P_2 \xrightarrow{\tau} Q_1|Q_2}$$

for inductive hypothesis  $P_1 \xrightarrow{\bar{x}y} Q_1$  and  $P_2 \xrightarrow{xy} Q_2$  so for rule *Com*  $P_1|P_2 \xrightarrow{\tau} Q_1|Q_2$

*Sum* :

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\alpha} Q}{P_1 + P_2 \xrightarrow{\alpha} Q}$$

for inductive hypothesis  $P_1 \xrightarrow{\alpha} Q$  and for rule *Sum*  $P_1 + P_2 \xrightarrow{\alpha} Q$ .

*Res* the first transition is:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\alpha} Q_1 \quad z \notin n(\gamma_1)}{(\nu z)P_1 \xrightarrow{\alpha} (\nu z)Q_1}$$

for inductive hypothesis  $P_1 \xrightarrow{\alpha} Q_1$  and for rule *Res*  $(\nu z)P_1 \xrightarrow{\alpha} (\nu z)Q_1$ .

*others* : other cases are similar.

□

Since it's important to give a low level semantic which is equivalent to the high level one, we can propose a change to the low level semantic that gets closer to our purpose. We replace the rule *Com3L*, *Com3R*, *Com2L* and *Com2R* with:

$$\begin{array}{ll} \mathbf{Com2LStop} \frac{L_1 \xrightarrow{\bar{x}y} L_2 \quad P \xrightarrow{xy} Q}{L_1|P \xrightarrow{\epsilon} L_2|stop(Q)} & \mathbf{Com2RStop} \frac{P \xrightarrow{xy} Q \quad L_1 \xrightarrow{\bar{x}y} L_2}{P|L_1 \xrightarrow{\epsilon} stop(Q)|L_2} \\ \mathbf{Com3LStop} \frac{P \xrightarrow{\bar{x}y} L \quad Q \xrightarrow{xy} Q'}{P|Q \xrightarrow{\epsilon} L|stop(Q')} & \mathbf{Com3RStop} \frac{P \xrightarrow{xy} P' \quad Q \xrightarrow{\bar{x}y} L}{P|Q \xrightarrow{\epsilon} stop(P')|L} \end{array}$$

where  $stop(P)$  is a multi  $\pi$  process which cannot make any transition.

**Definition 3.2.4.** The *erase function*  $er$  is a function that eliminates the *stop* mark on processes. Its definition is straightforward.

**Proposition 3.2.5.** Let  $\rightarrow$  be the relation defined in table 3.1.

- If  $P \xrightarrow{\sigma} Q$  then there exist  $L_1, \dots, L_k$  with  $k \geq 0$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q' \quad \text{and} \quad \gamma_1 \dots \gamma_{k+1} = \sigma \quad \text{and} \quad er(Q') = Q$$

- If there exist  $L_1, \dots, L_k$  with  $k \geq 1$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q$$

where at most one  $\gamma$  is an output whether all the other  $\gamma$ s are  $\epsilon$  or  $\tau$  then  $P \xrightarrow{\tau} er(Q)$  or if there is an output  $\bar{x}y$  in the  $\gamma$ s then  $P \xrightarrow{\bar{x}y} er(Q)$ .

*Proof.* The proof of the first part of this proposition is almost exactly as the proof of proposition 3.2.3. The proof of the second part is by induction on the depth of the derivation of the first transition:

**base case** The last rule in the derivation of  $P \xrightarrow{\gamma_1} L_1$  can be only *SOutLow*:

$$\mathbf{SOutLow} \frac{}{\underbrace{\bar{x}y.P_1}_P \xrightarrow{\bar{x}y} \underbrace{*P_1}_{L_1}}$$

since  $*P_1$  has a mark at the top level, the last rule used to derive  $*P_1 \xrightarrow{\gamma_2}$  has to be *StarEps* so we have  $P_1 \xrightarrow{\gamma_2} L_2$  or  $P_1 \xrightarrow{\gamma_2} Q$  depending on  $k$ . We can build the following chain of transition:

$$P_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q$$

since  $\gamma_1$  is an output, the other  $\gamma$ s are  $\epsilon$  or  $\tau$ , then we can apply the inductive hypothesis to get  $P_1 \xrightarrow{\tau} er(Q)$ . Now a proof of the conclusion is

$$\mathbf{SOutTau} \frac{P_1 \xrightarrow{\tau} er(Q)}{\bar{x}y.P_1 \xrightarrow{\bar{x}y} er(Q)}$$

**inductive case** The last rule in the derivation of  $P \xrightarrow{\gamma_1} L_1$  can be:

*Sum* the first transition is:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\gamma_1} L_1}{P_1 + P_2 \xrightarrow{\gamma_1} L_1}$$

so we can build the following chain of transition:

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q$$

apply the inductive hypothesis to get  $P_1 \xrightarrow{\alpha} er(Q)$  where  $\alpha$  is  $\tau$  or an output. Now a proof of the conclusion is

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\alpha} er(Q)}{P_1 + P_2 \xrightarrow{\alpha} er(Q)}$$

*Res* the first transition is:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\gamma_1} L'_1 \quad z \notin n(\gamma_1)}{(\nu z)P_1 \xrightarrow{\gamma_1} (\nu z)L'_1}$$

given that  $L_1$  has a restriction at the top level, all the other intermediate processes  $L_2, \dots, L_k$  and  $Q$  have the same restriction at the top level. This is because the only rule whose conclusion is a transition that start from a possibly marked process with a restriction at its top level is *Res*. So the last rule used to prove all transition is *Res*.

$$\mathbf{Res} \frac{L'_k \xrightarrow{\gamma_{k+1}} Q' \quad z \notin n(\tau)}{(\nu z)L'_k \xrightarrow{\gamma_{k+1}} (\nu z)Q'} \quad \mathbf{Res} \frac{L'_i \xrightarrow{\gamma_i} L'_{i+1} \quad z \notin n(\epsilon)}{(\nu z)L'_i \xrightarrow{\gamma_i} (\nu z)L'_{i+1}}$$

we can build the following chain of transitions:

$$P_1 \xrightarrow{\gamma_1} L'_1 \xrightarrow{\gamma_2} L'_2 \cdots L'_{k-1} \xrightarrow{\gamma_k} L'_k \xrightarrow{\gamma_{k+1}} Q'$$

then apply the inductive hypothesis to get  $P_1 \xrightarrow{\alpha} er(Q')$ . A proof of the conclusion can be

$$\mathbf{Res} \frac{P_1 \xrightarrow{\alpha} er(Q') \quad z \notin n(\tau)}{(\nu z)P_1 \xrightarrow{\alpha} (\nu z)er(Q') = er((\nu z)Q')}$$

*Cong* the last rule of the derivation of the first transition is:

$$\mathbf{Cong} \frac{P' \equiv P \quad \vdash^{\gamma_1}_{\rightarrow} L_1}{P \vdash^{\gamma_1}_{\rightarrow} L_1}$$

We derive the following chain of transition:

$$P' \vdash^{\gamma_1}_{\rightarrow} L_1 \vdash^{\gamma_2}_{\rightarrow} L_2 \cdots L_{k-1} \vdash^{\gamma_k}_{\rightarrow} L_k \vdash^{\gamma_{k+1}}_{\rightarrow} Q$$

for inductive hypothesis  $P' \xrightarrow{\alpha} er(Q)$ . A proof of the conclusion is

$$\mathbf{Cong} \frac{P' \equiv P \quad P' \xrightarrow{\alpha} er(Q)}{P \xrightarrow{\alpha} er(Q)}$$

*Com3LStop* : the last part of the derivation of the first transition is:

$$\mathbf{Com3LStop} \frac{P_1 \xrightarrow{\bar{x}y} L'_1 \quad P_2 \xrightarrow{xy} Q_2}{P_1|P_2 \xrightarrow{\epsilon} L'_1|stop(Q_2)}$$

the derivations of all other transitions can end only with an instance of *Par1L* so we have:

$$\mathbf{Par1L} \frac{L'_i \vdash^{\gamma_i}_{\rightarrow} L'_{i+1}}{L'_i|stop(Q_2) \vdash^{\gamma_i}_{\rightarrow} L'_{i+1}|stop(Q_2)} \quad \mathbf{Par1L} \frac{L'_k \vdash^{\gamma_{k+1}}_{\rightarrow} Q_1}{L'_i|stop(Q_2) \vdash^{\gamma_{k+1}}_{\rightarrow} Q_1|stop(Q_2)}$$

We derive the following chain of transition:

$$P_1 \xrightarrow{\bar{x}y} L'_1 \vdash^{\gamma_2}_{\rightarrow} L'_2 \cdots L'_{k-1} \vdash^{\gamma_k}_{\rightarrow} L'_k \vdash^{\gamma_{k+1}}_{\rightarrow} Q_1$$

for inductive hypothesis  $P_1 \xrightarrow{\bar{x}y} er(Q_1)$ . A proof of the conclusion is

$$\mathbf{EComSeq} \frac{P_1 \xrightarrow{\bar{x}y} er(Q_1) \quad P_2 \xrightarrow{xy} Q_2}{P_1|P_2 \xrightarrow{\tau} er(Q_1)|Q_2}$$

*Par1L* : the last part of the derivation of the first transition is:

$$\mathbf{Par1L} \frac{P_1 \vdash^{\gamma_1}_{\rightarrow} L'_1}{P_1|P_2 \vdash^{\gamma_1}_{\rightarrow} L'_1|P_2}$$

there can be three cases:

- the derivations of all the other transitions end with an instance of *Par1L*. We derive the following chain of transition:

$$P_1 \vdash^{\gamma_1}_{\rightarrow} L'_1 \vdash^{\gamma_2}_{\rightarrow} L'_2 \cdots L'_{k-1} \vdash^{\gamma_k}_{\rightarrow} L'_k \vdash^{\gamma_k}_{\rightarrow} Q_1$$

for inductive hypothesis  $P_1 \xrightarrow{\alpha} er(Q_1)$ . A proof of the conclusion is

$$\mathbf{Par} \frac{P_1 \xrightarrow{\alpha} er(Q_1)}{P_1|P_2 \xrightarrow{\alpha} er(Q_1)|P'_2}$$

- there is one derivation that ends with an instance of *Com2LStop* and the derivations of all the other transitions end with an instance of *Par1L*. We present here the case when the second transition ends with a *Com2LStop*, the other cases are similar. So

$$\mathbf{Com2LStop} \frac{L'_2 \xrightarrow{\bar{x}y} L'_2 \quad P_2 \xrightarrow{xy} P'_2}{L'_2|P_2 \xrightarrow{\epsilon} L'_2|stop(P'_2)}$$

We derive the following chain of transition:



---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(z).P \xrightarrow{xw} P\{w/z\}}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$
<b>SOut</b> $\frac{P \xrightarrow{\gamma} P'}{\bar{x}y.P \xrightarrow{\bar{x}y.\gamma} P'}$	$\gamma$ is a non empty sequence of outputs	
<b>EComSeq</b> $\frac{P \xrightarrow{\bar{x}y.\sigma} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\sigma} P' Q'}$	<b>ECom</b> $\frac{P \xrightarrow{\bar{x}y} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\tau} P' Q'}$	
<b>ParL</b> $\frac{P \xrightarrow{\sigma} P' \quad bn(\sigma) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\sigma} P' Q}$	<b>ParR</b> $\frac{Q \xrightarrow{\sigma} Q' \quad bn(\sigma) \cap fn(P) = \emptyset}{P Q \xrightarrow{\sigma} P Q'}$	
<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\sigma)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$	<b>Ide</b> $\frac{A(\tilde{x}) \stackrel{def}{=} P \quad P\{\tilde{y}/\tilde{x}\} \xrightarrow{\sigma} Q}{A \xrightarrow{\sigma} Q}$	
<b>SumL</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>SumR</b> $\frac{Q \xrightarrow{\sigma} Q'}{P + Q \xrightarrow{\sigma} Q'}$	

---

Table 3.3: Multi $\pi$  early semantic without structural congruence

$P_1 \xrightarrow{\epsilon} L'_1 \xrightarrow{\bar{x}y} L'_2 \xrightarrow{\epsilon} \dots \xrightarrow{\epsilon} L'_{k-1} \xrightarrow{\epsilon} L'_k \xrightarrow{\tau} Q_1$   
 for inductive hypothesis  $P_1 \xrightarrow{\bar{x}y} er(Q_1)$ . A proof of the conclusion is

$$\mathbf{EComSeq} \frac{P_1 \xrightarrow{\bar{x}y} er(Q_1) \quad P_2 \xrightarrow{xy} P'_2}{P_1|P_2 \xrightarrow{\tau} er(Q_1)|P'_2}$$

- the derivation of the last transition ends with an instance of *Com4L* and the derivations of all the other transitions end with an instance of *Par1L*. We derive the following chain of transition:

$P_1 \xrightarrow{\epsilon} L'_1 \xrightarrow{\epsilon} L'_2 \dots L'_{k-1} \xrightarrow{\epsilon} L'_k \xrightarrow{\bar{x}y} Q_1$   
 for inductive hypothesis  $P_1 \xrightarrow{\bar{x}y} er(Q_1)$ . A proof of the conclusion is

$$\mathbf{EComSeq} \frac{P_1 \xrightarrow{\bar{x}y} er(Q_1) \quad P_2 \xrightarrow{xy} P'_2}{P_1|P_2 \xrightarrow{\tau} er(Q_1)|P'_2}$$

□

### 3.2.3 Early operational semantic without structural congruence

**Definition 3.2.5.** The *late transition relation without structural congruence* is the smallest relation induced by the rules in table 3.3.

**Example** Scope extrusion with strong prefixing(1).  $x \notin fn(y(z).Q|a(b).R|y(z).S)$ . The following is the desired transition:

$$(\nu x)(\bar{y}x.\bar{a}b.\bar{y}x.\bar{a}b.P)|y(z).Q|a(b).R|y(z).S|a(b).T \xrightarrow{\tau} (\nu x)(P|Q\{x/z\}|S\{x/z\})|R|T$$

It is possible to infer this transition in the semantic with structural congruence. But without structural congruence and the following scope extrusion rules

$$\mathbf{Opn} \frac{P \xrightarrow{\sigma} P' \quad y \in obj(\sigma) \quad y \notin sbj(\sigma)}{(\nu y)P \xrightarrow{opn(\sigma,y)} P'}$$

$$\text{Cls} \frac{P \xrightarrow{\bar{x}(z)} P' \quad Q \xrightarrow{xz} Q'}{P|Q \xrightarrow{\tau} (\nu z)(P'|Q')} \quad \text{ClsSeq} \frac{P \xrightarrow{\bar{x}(z) \cdot \sigma} P' \quad Q \xrightarrow{xz} Q'}{P|Q \xrightarrow{\sigma} (\nu z)(P'|Q')}$$

we can only infer

$$(\nu x)(\bar{y}x.\bar{a}b.\bar{y}x.\bar{a}b.P)|y(z).Q|a(b).R|y(z).S|a(b).T \xrightarrow{\tau} (\nu x)((\nu x)(P|Q\{x/z\})|R|S\{x/z\})|T$$

This transition is not what we want because now the scope of the inner  $\nu x$  hides in  $P$  the scope of the outer  $\nu x$ , so  $P$  and  $S$  cannot use  $x$  to communicate. But with these rules:

$$\begin{aligned} \text{Opn} & \frac{P \xrightarrow{\sigma} P' \quad y \in \text{obj}(\sigma) \quad y \notin \text{sbj}(\sigma)}{(\nu y)P \xrightarrow{\text{opn}(\sigma, y)} P'} \\ \text{Cls} & \frac{P \xrightarrow{\bar{x}(y) \cdot (\nu y)} P' \quad Q \xrightarrow{xy} Q'}{P|Q \xrightarrow{\tau} (\nu y)(P'|Q')} \quad \text{ClsSeq1} \frac{P \xrightarrow{\bar{x}(y) \cdot (\nu y) \cdot \sigma} P' \quad Q \xrightarrow{xy} Q'}{P|Q \xrightarrow{\sigma} (\nu y)(P'|Q')} \\ \text{ClsSeq2} & \frac{P \xrightarrow{\bar{x}(y) \cdot \sigma} P' \quad Q \xrightarrow{xy} Q'}{P|Q \xrightarrow{\sigma} P'|Q'} \quad \sigma \text{ does not start with } \nu \end{aligned}$$

$$\begin{aligned} \text{opn}(\bar{x}y, y) &= \bar{x}(y) \cdot (\nu y) & \text{opn}(\bar{x}y \cdot \sigma, y) &= \begin{cases} \bar{x}(y) \cdot \text{opn}(\sigma, y) & \text{if } y \in \text{obj}(\sigma) \\ \bar{x}(y) \cdot (\nu y) \cdot \text{opn}(\sigma, y) & \text{if } y \notin \text{obj}(\sigma) \end{cases} \\ \text{opn}(\bar{x}z, y) &= \bar{x}z & \text{opn}(\bar{x}z \cdot \sigma, y) &= \bar{x}z \cdot \text{opn}(\sigma, y) \\ \text{opn}((\nu z), y) &= (\nu z) & \text{opn}((\nu z) \cdot \sigma, y) &= (\nu z) \cdot \text{opn}(\sigma, y) \end{aligned}$$

$$\begin{aligned} \text{sbj}(\tau) &= \emptyset & \text{sbj}(\bar{x}y) &= \{x\} & \text{sbj}(x(y)) &= \{x\} & \text{sbj}((\nu y)) &= \emptyset & \text{sbj}(\alpha \cdot \sigma) &= \text{sbj}(\alpha) \cup \text{sbj}(\sigma) \\ \text{obj}(\tau) &= \emptyset & \text{obj}(\bar{x}y) &= \{y\} & \text{obj}(x(y)) &= \{y\} & \text{obj}((\nu y)) &= \emptyset & \text{obj}(\alpha \cdot \sigma) &= \text{obj}(\alpha) \cup \text{obj}(\sigma) \end{aligned}$$

The following transition can be inferred:

$$(\nu x)(\bar{y}x.\bar{a}b.\bar{y}x.\bar{a}b.P)|y(z).Q|a(b).R|y(z).S|a(b).T \xrightarrow{\tau} (\nu x)(P|Q\{x/z\})|R|S\{x/z\})|T$$

**Example** Scope intrusion without strong prefixing.

$$\bar{y}x.P|(\nu x)(y(z).Q) \xrightarrow{\tau} P|(\nu w)(Q\{w/x\}\{x/z\})$$

This transition cannot be derived without *alpha* conversion.

**Example** Scope extrusion without strong prefixing.

$$\begin{aligned} \text{Out} & \frac{}{\bar{y}x.P \xrightarrow{\bar{y}x} P} \\ \text{Opn} & \frac{}{(\nu x)(\bar{y}x.P) \xrightarrow{\bar{y}(x)(\nu x)} P} \quad \text{EInp} \frac{}{y(z).Q \xrightarrow{yx} Q\{x/z\}} \\ \text{Cls} & \frac{}{(\nu x)(\bar{y}x.P)|y(z).Q \xrightarrow{\tau} (\nu x)(P|Q\{x/z\})} \end{aligned}$$

**Example** Scope extrusion with strong prefixing(2).  $x \in \text{fn}(y(z).Q|a(b).R|y(z).S)$  and  $x' \notin \text{fn}(y(z).Q|a(b).R|y(z).S)$

$$(\nu x)(\bar{y}x.\bar{a}b.\bar{y}x.\bar{a}b.P)|y(z).Q|a(b).R|y(z).S|a(b).T \xrightarrow{\tau} (\nu x')(P\{x'/x\}|Q\{x'/z\})|R|S\{x'/z\})|T$$

This transition cannot be derived without  $\alpha$  conversion.

**Example** Scope intrusion with strong prefixing.

$$\bar{y}x.\bar{a}b.P|(\nu x)(y(z).Q)|(\nu b)(a(c).R) \xrightarrow{\tau} P|(\nu w)(Q\{w/x\}\{x/z\})|(\nu d)(R\{d/b\}\{b/c\})$$

This transition cannot be derived without  $\alpha$  conversion.

### 3.3 Strong bisimilarity and equivalence

#### 3.3.1 Strong bisimilarity

In the following section,  $\rightarrow$  is the transition relation defined in table 3.3.

**Definition 3.3.1.** A *strong early bisimulation* is a symmetric binary relation  $\mathbf{S}$  on multi  $\pi$  processes such that for all  $P$  and  $Q$ :  $PSQ$ ,  $P \xrightarrow{\gamma} P'$  and  $bn(\gamma)$  is fresh imply that

$$\exists Q' : Q \xrightarrow{\gamma} Q' \text{ and } P' \mathbf{S} Q'$$

The *strong early bisimilarity*, written  $\sim_E$ , is the union of all strong early bisimulation. Two processes  $P, Q$  are *strong early bisimilar*, written  $P \sim_E Q$ , if they are related by the strong early bisimilarity.

**Proposition 3.3.1.** The strong early bisimilarity is a strong early bisimulation.

**Proposition 3.3.2.**  $\sim_E$  is an equivalence relation.

*Proof.* :

**Reflexivity** The identity relation on processes is a strong early bisimulation.

**Simmetry** It is in the definition.

**Transitivity** The composition  $\sim_E \sim_E$  is a strong early bisimulation.

□

**Definition 3.3.2.** A *strong early bisimulation up to  $\sim_E$*  is a symmetric binary relation  $\mathbf{S}$  on multi  $\pi$  processes such that for all  $P$  and  $Q$ :  $PSQ$ ,  $P \xrightarrow{\gamma} P'$  and  $bn(\gamma)$  is fresh imply that

$$\exists P'', Q', Q'' : Q \xrightarrow{\gamma} Q' \text{ and } P' \sim_E P'' \mathbf{S} Q'' \sim_E Q'$$

Two processes  $P, Q$  are *strong early bisimilar up to  $\sim_E$* , written  $P \sim_E^{up} Q$ , if they are related by a strong early bisimulation up to  $\sim_E$ .

**Proposition 3.3.3.**  $P \sim_E^{up} Q$  imply  $P \sim_E Q$ .

*Proof.* Let  $\mathbf{S}$  be a bisimulation up to  $\sim_E$  such that  $PSQ$ . It can be proved that  $\sim_E \mathbf{S} \sim_E$  is a bisimulation: let  $A \sim_E B \mathbf{S} C \sim_E D$

$$\begin{aligned} A \xrightarrow{\gamma} A' \wedge A \sim_E B \wedge \text{definition 3.3.1} &\Rightarrow \exists B' : B \xrightarrow{\gamma} B' \wedge A' \sim_E B' \\ B \mathbf{S} C \wedge \text{definition 3.3.2} &\Rightarrow \exists C' C'' B'' : C \xrightarrow{\gamma} C' \wedge B' \sim_E B'' \mathbf{S} C'' \sim_E C' \\ C \xrightarrow{\gamma} C' \wedge C \sim_E D \wedge \text{definition 3.3.1} &\Rightarrow \exists D' : D \xrightarrow{\gamma} D' \wedge C' \mathbf{S} D' \\ A' \sim_E B' \sim_E B'' \mathbf{S} C'' \sim_E C' \sim_E D' \wedge \text{transitivity of } \sim_E &\Rightarrow A' \sim_E B'' \mathbf{S} C'' \sim_E D' \end{aligned}$$

It is easy to see that the symmetric also holds.

□

**Proposition 3.3.4.**  $\equiv_\alpha$  is a strong bisimulation.

*Proof.* We prove that if  $P \equiv_\alpha Q$  and  $P \xrightarrow{\gamma} P'$  then  $Q \xrightarrow{\gamma} Q'$  and  $P' \equiv_\alpha Q'$ . The symmetric holds because  $\alpha$  equivalence is symmetric. The proof proceed by induction on the derivation of  $P \xrightarrow{\gamma} P'$ . The last rule used can be:

**Out** :  $P$  is  $\bar{x}y.P_1$  and  $\gamma$  is  $\bar{x}y$  for some names  $x, y$  and process  $P_1$ .  $P \equiv_\alpha Q$  and the inversion lemma for  $\alpha$  equivalence imply  $Q$  is  $\bar{x}y.Q_1$  and  $P_1 \equiv_\alpha Q_1$  for a process  $Q_1$ . Rule *Out* proves  $\bar{x}y.Q_1 \xrightarrow{\bar{x}y} Q_1$

**EInp** :  $P$  is  $x(y).P_1$  and  $\gamma$  is  $xz$  for some names  $x, y, z$  and process  $P_1$ .  $P \equiv_\alpha Q$  and the inversion lemma for  $\alpha$  equivalence imply  $Q$  is  $x(w).Q_1$  and  $P_1 \equiv_\alpha Q_1\{y/w\}$  for a process  $Q_1$  such that  $y \notin fn(Q_1)$  and a name  $w$  which is not necessarily different from  $y$ . Rule *EInp* proves  $x(w).Q_1 \xrightarrow{xz} Q_1$

**Res** : similar to the previous case.

**Tau** :  $P$  is  $\tau.P_1$  and  $\gamma$  is  $\tau$  for some process  $P_1$ .  $P \equiv_\alpha Q$  and the inversion lemma for  $\alpha$  equivalence imply  $Q$  is  $\tau.Q_1$  and  $P_1 \equiv_\alpha Q_1$  for a process  $Q_1$ . Rule *Tau* proves  $\tau.Q_1 \xrightarrow{\tau} Q_1$

**SOut** :  $P$  is  $\overline{xy}.P_1$  and  $\gamma$  is  $\overline{xy} \cdot \gamma'$  for some names  $x, y$ , process  $P_1$  and a non empty sequence of outputs  $\gamma'$ .  $P \equiv_\alpha Q$  and the inversion lemma for  $\alpha$  equivalence imply  $Q$  is  $\overline{xy}.Q_1$  and  $P_1 \equiv_\alpha Q_1$  for a process  $Q_1$ . The premise is  $P_1 \xrightarrow{\gamma'} P'_1$ , for inductive hypothesis  $Q_1 \xrightarrow{\gamma'} Q'_1$  and  $P'_1 \equiv_\alpha Q'_1$ . Rule *SOut* proves  $\overline{xy}.Q_1 \xrightarrow{\overline{xy} \cdot \gamma'} Q_1$

**EComSeq, ECom, ParL, ParR, SumL, SumR, Ide** : similar to the previous case.

**Opn** :

**Cls** :

**ClsSeq1** :

**ClsSeq2** :

□

**Proposition 3.3.5.**  $\sim_E$  is preserved by all operators except input prefixing.

*Proof.* The proof goes by cases on operators:

**Output prefixing** The relation  $\{(\overline{xy}.P, \overline{xy}.Q) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation.

**Strong output prefixing** The relation  $\{(\overline{xy}.P, \overline{xy}.Q) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation: there are two cases to consider:

- If there exists a transition  $P \xrightarrow{\gamma} P'$  where  $\gamma$  is a non empty sequence of outputs then we can apply the rule *SOut*:

$$\frac{P \xrightarrow{\gamma} P'}{\overline{xy}.P \xrightarrow{\overline{xy} \cdot \gamma} P'}$$

$P \xrightarrow{\gamma} P'$  and  $P \sim_E Q$  imply  $Q \xrightarrow{\gamma} Q'$  and  $P' \sim_E Q'$ . For rule *SOut*:  $\overline{xy}.Q \xrightarrow{\overline{xy} \cdot \gamma} Q'$  so the conclusion holds.

- Otherwise there is no transition starting from  $\overline{xy}.P$  or from  $\overline{xy}.Q$  so these processes are strongly bisimilar.

**Tau prefixing** The relation  $\{(\tau.P, \tau.Q) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation.

**Summation** The relation  $\{(P + R, Q + R) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation.

**Restriction** The relation  $\{((\nu x)P, (\nu x)Q) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation.

**Parallel composition** The relation  $\{(P|R, Q|R) : P \sim_E Q\} \cup \sim_E$  is a strong early bisimulation. The last rule applicable to  $P|R$  can be:

**ECom** :

$$\frac{P \xrightarrow{\overline{xy}} P' \quad R \xrightarrow{xy} R'}{P|R \xrightarrow{\tau} P'|R'}$$

$P \xrightarrow{\overline{xy}} P'$  and  $P \sim_E Q$  imply that there exists a process  $Q'$  such that  $Q \xrightarrow{\overline{xy}} Q'$  and  $P' \sim_E Q'$ . So for rule *ECom*:  $Q|R \xrightarrow{\tau} Q'|R'$  and  $P'|R' \sim_E Q'|R'$

**Cls** :

$$\frac{P \xrightarrow{\bar{x}(y) \cdot (\nu y)} P' \quad R \xrightarrow{xy} R'}{P|R \xrightarrow{\tau} (\nu y)(P'|R')}$$

$P \xrightarrow{\bar{x}(y) \cdot (\nu y)} P'$  and  $P \sim_E Q$  imply that there exists a process  $Q'$  such that  $Q \xrightarrow{\bar{x}y \cdot (\nu y)} Q'$  and  $P' \sim_E Q'$ . So for rule *Cls*:  $Q|R \xrightarrow{\tau} (\nu y)(Q'|R')$  and  $(\nu y)(P'|R') \sim_E (\nu y)(Q'|R')$

**ClsSeq1, ClsSeq2, ParL, ParR** similar.

□

**Example**  $\sim_E$  is not in general preserved by input prefixing because:

$$a(x).0|\bar{b}y.0 \sim_E a(x).\bar{b}y.0 + \bar{b}y.a(x).0$$

but

$$c(a).(a(x).0|\bar{b}y.0) \not\sim_E c(a).(a(x).\bar{b}y.0 + \bar{b}y.a(x).0)$$

because

$$\begin{aligned} c(a).(a(x).0|\bar{b}y.0) &\xrightarrow{cb} b(x).0|\bar{b}y.0 \xrightarrow{\tau} 0|0 \\ c(a).(a(x).\bar{b}y.0 + \bar{b}y.a(x).0) &\xrightarrow{cb} b(x).\bar{b}y.0 + \bar{b}y.b(x).0 \not\xrightarrow{\tau} \end{aligned}$$

### 3.3.2 Open bisimilarity

**Definition 3.3.3.** A *distinction* is a finite symmetric and irreflexive binary relation on names. A substitution  $\sigma$  *respects* a distinction  $D$  if

$$\forall a, b. aDb \Rightarrow a\sigma \neq b\sigma$$

We write  $D\sigma$  for the composition of the two relation.

**Definition 3.3.4.**  $P$  and  $Q$  are *strongly  $D$  equivalent*, written  $P \sim^D Q$ , if for all substitution  $\sigma$  respecting  $D$ :  $P\sigma \sim_E Q\sigma$ .

**Theorem 3.3.6.**  $\sim^D$  is a congruence.

*Proof.*  $\sim^D$  is an equivalence relation because  $\sim_E$  is an equivalence relation.

*Reflexivity* Since  $\sim_E$  is reflexive, for all substitution  $\sigma$  respecting  $D$ :  $P\sigma \sim_E Q\sigma$  so  $P \sim^D P$

*Symmetry* Let  $P \sim^D Q$  then for all substitution  $\sigma$  respecting  $D$ :  $P\sigma \sim_E Q\sigma$ . Since  $\sim_E$  is symmetric  $Q\sigma \sim_E P\sigma$  so  $Q \sim^D P$

*Transitivity* Let  $P \sim^D Q$  and  $Q \sim^D R$  then for all substitution  $\sigma$  respecting  $D$ :  $P\sigma \sim_E Q\sigma$  and  $Q\sigma \sim_E R\sigma$ . Since  $\sim_E$  is transitive  $P\sigma \sim_E R\sigma$  so  $P \sim^D R$ .

$\sim^D$  is preserved by every operator. Let  $P \sim^D Q$  and let  $\sigma$  be a substitution respecting  $D$  so  $P\sigma \sim_E Q\sigma$ :

**Output prefixing** The only transitions starting from  $(\bar{x}y.P)\sigma$  and  $(\bar{x}y.Q)\sigma$  are

$$(\bar{x}y.P)\sigma \xrightarrow{(\bar{x}y)\sigma} P\sigma \sim_E Q\sigma \xleftarrow{(\bar{x}y)\sigma} (\bar{x}y.Q)\sigma$$

so  $(\bar{x}y.P)\sigma \sim_E (\bar{x}y.Q)\sigma$  and  $\bar{x}y.P \sim^D \bar{x}y.Q$

**Strong output prefixing** There are two cases to consider:

- If there exists a transition  $P\sigma \xrightarrow{\gamma} P'$  where  $\gamma$  is a non empty sequence of outputs then

$$\mathbf{SOut} \frac{P\sigma \xrightarrow{\gamma} P'}{(\bar{x}y.P)\sigma \xrightarrow{(\bar{x}y)\sigma \cdot \gamma} P'}$$

$P\sigma \xrightarrow{\gamma} P'$  and  $P\sigma \sim_E Q\sigma$  imply  $Q\sigma \xrightarrow{\gamma} Q'$  and  $P' \sim_E Q'$ . For rule *SOut*

$$(\underline{\bar{x}y}.Q)\sigma \xrightarrow{(\bar{x}y)\sigma \cdot \gamma} Q'$$

so  $(\underline{\bar{x}y}.P)\sigma \sim_E (\underline{\bar{x}y}.Q)\sigma$  and  $\underline{\bar{x}y}.P \dot{\sim}^D \underline{\bar{x}y}.Q$ .

- Otherwise there is no transition starting from  $(\underline{\bar{x}y}.P)\sigma$  or from  $(\underline{\bar{x}y}.Q)\sigma$  so  $(\underline{\bar{x}y}.P)\sigma \dot{\sim}_E (\underline{\bar{x}y}.Q)\sigma$  and  $\underline{\bar{x}y}.P \dot{\sim}^D \underline{\bar{x}y}.Q$ .

**Input prefixing** Let  $z \notin \text{bn}(P) \cup \text{bn}(Q)$  and let  $y \notin \text{img}(\sigma)$ . The transitions starting from  $(x(y).P)\sigma$  and  $(x(y).Q)\sigma$  are

$$x\sigma(y).P\sigma \xrightarrow{x\sigma z} P\sigma\{z/y\} \quad x\sigma(y).Q\sigma \xrightarrow{x\sigma z} Q\sigma\{z/y\}$$

?

**Tau prefixing**

**Summation**

**Restriction**

**Parallel composition**

□

## Chapter 4

# Multi $\pi$ calculus with strong input

### 4.1 Syntax

As we did with  $\pi$  calculus, we suppose that we have a countable set of names  $\mathbb{N}$ , ranged over by lower case letters  $a, b, \dots, z$ . These names are used for communication channels and values. Furthermore we have a set of identifiers, ranged over by  $A$ . We represent the agents or processes by upper case letters  $P, Q, \dots$ . A multi  $\pi$  process, in addition to the same actions of a  $\pi$  process, can perform also a strong prefix input:

$$\pi ::= \bar{x}y \mid x(z) \mid \underline{x(y)} \mid \tau$$

The processes are defined, just as original  $\pi$  calculus, by the following grammar:

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A(y_1, \dots, y_n)$$

and they have the same intuitive meaning as for the  $\pi$  calculus. The strong prefix input allows a process to make an atomic sequence of actions, so that more than one process can synchronize on this sequence. For the moment we allow the strong prefix to be on input names only. Also one can use the strong prefix only as an action prefixing for processes that can make at least a further action.

Multi  $\pi$  calculus is a conservative extension of the  $\pi$  calculus in the sense that: any  $\pi$  calculus process  $p$  is also a multi  $\pi$  calculus process and the semantic of  $p$  according to the SOS rules of  $\pi$  calculus is the same as the semantic of  $p$  according to the SOS rules of multi  $\pi$  calculus. We have to extend the following definition to deal with the strong prefix:

$$B(\underline{x(y)}.Q, I) = \{y, \bar{y}\} \cup B(Q, I) \quad F(\underline{x(y)}.Q, I) = \{x, \bar{x}\} \cup (F(Q, I) - \{y, \bar{y}\})$$

In this setting two processes cannot synchronize on a sequence of actions with length greater than one so we cannot have transactional synchronization but we can have multi-party synchronization.

### 4.2 Operational semantic

#### 4.2.1 Early operational semantic with structural congruence

The semantic of a multi  $\pi$  process is a labeled transition system such that

- the nodes are multi  $\pi$  calculus processes. The set of nodes is  $\mathbb{P}_m$
- the actions are multi  $\pi$  calculus actions. The set of actions is  $\mathbb{A}_m$ , we use  $\alpha, \alpha_1, \alpha_2, \dots$  to range over the set of actions, we use  $\sigma, \sigma_1, \sigma_2, \dots$  to range over the set  $\mathbb{A}_m^+ \cup \{\tau\}$ .
- the transition relations is  $\rightarrow \subseteq \mathbb{P}_m \times (\mathbb{A}_m^+ \cup \{\tau\}) \times \mathbb{P}_m$

In this case, a label can be a sequence of prefixes, whether in the original  $\pi$  calculus a label can be only a prefix. We use the symbol  $\cdot$  to denote the concatenation operator.

---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>EInp</b> $\frac{}{x(y).P \xrightarrow{xz} P\{z/y\}}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$
<b>SInpTau</b> $\frac{P\{y/z\} \xrightarrow{\tau} P'}{x(z).P \xrightarrow{xy} P'}$	<b>SInp</b> $\frac{P\{y/z\} \xrightarrow{ab} P'}{x(z).P \xrightarrow{xy \cdot ab} P'}$	<b>SInpSeq</b> $\frac{P\{y/z\} \xrightarrow{\sigma} P' \quad  \sigma  > 1}{x(z).P \xrightarrow{xy \cdot \sigma} P'}$
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\alpha} Q}{P \xrightarrow{\alpha} Q}$	<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\sigma)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$
<b>Par</b> $\frac{P \xrightarrow{\sigma} P'}{P Q \xrightarrow{\sigma} P' Q}$	<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$	
<b>ECom</b> $\frac{P \xrightarrow{xy} P' \quad Q \xrightarrow{\bar{x}y} Q'}{P Q \xrightarrow{\tau} P' Q'}$	<b>EComSeq</b> $\frac{P \xrightarrow{xy \cdot \sigma} P' \quad Q \xrightarrow{\bar{x}y} Q'}{P Q \xrightarrow{\sigma} P' Q'}$	

---

Table 4.1: Multi  $\pi$  early semantic with structural congruence

**Definition 4.2.1.** The *early transition relation with structural congruence* is the smallest relation induced by the rules in table 4.1 where *inpSeq* is a non empty sequence of input actions and  $\sigma$  is a sequence of any action.

**Example** Multi-party synchronization We show an example of a derivation of three processes that synchronize.

$$\begin{array}{c}
\text{EInp} \frac{}{(x(b).P)\{y/a\} \xrightarrow{xz} P\{y/a\}\{z/b\}} \\
\text{SInp} \frac{}{x(a).(x(b).P) \xrightarrow{xy \cdot xz} P\{y/a\}\{z/b\}} \\
\text{EComSeq} \frac{}{x(a).x(b).P|\bar{x}y.Q \xrightarrow{xz} P\{y/a\}\{z/b\}|Q} \\
\text{Out} \frac{}{\bar{x}y.Q \xrightarrow{\bar{x}y} Q} \\
\text{EComSng} \frac{x(a).x(b).P|\bar{x}y.Q \xrightarrow{xz} P\{y/a\}\{z/b\}|Q \quad \text{Out} \frac{}{\bar{x}z.R \xrightarrow{\bar{x}z} R}}{(x(a).x(b).P|\bar{x}y.Q)|\bar{x}z.R \xrightarrow{\tau} (P\{y/a\}\{z/b\}|Q)|R}
\end{array}$$

**Lemma 4.2.1.** If  $P \xrightarrow{\sigma} Q$  then only one of the following cases hold:

- $|\sigma| = 1$
- $|\sigma| > 1$ , the actions in  $\sigma$  are input.

#### 4.2.2 Late operational semantic with structural congruence

**Definition 4.2.2.** The *late transition relation with structural congruence* is the smallest relation induced by the rules in table 4.2.

**Example** Multi-party synchronization We show an example of a derivation of three processes that synchronize with the late semantic. The three processes are  $x(a).x(b).P$ ,  $\bar{x}y.Q$  and  $\bar{x}z.R$ . We assume modulo  $\alpha$  conversion that:

$$a \notin fn(x(b)) \cup fn(x(a).x(b).P)$$

and

$$c \notin fn(\bar{x}y.Q)$$



---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>LInp</b> $\frac{}{x(y).P \xrightarrow{x(y)} P}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$
<b>SInpTau</b> $\frac{P \xrightarrow{\tau} P'}{x(z).P \xrightarrow{x(z)} P'}$	<b>SInp</b> $\frac{P \xrightarrow{a(b)} P'}{x(z).P \xrightarrow{x(z).a(b)} P'}$	<b>SInpSeq</b> $\frac{P \xrightarrow{\sigma} P' \quad  \sigma  > 1}{x(z).P \xrightarrow{x(z).\sigma} P'}$
<b>LComSeq</b> $\frac{P \xrightarrow{x(y).\sigma} P' \quad Q \xrightarrow{\bar{x}z} Q'}{P Q \xrightarrow{\sigma\{z/y\}} P'\{z/y\} Q'}$	<b>LCom</b> $\frac{P \xrightarrow{x(y)} P' \quad Q \xrightarrow{\bar{x}z} Q'}{P Q \xrightarrow{\tau} P'\{z/y\} Q'}$	
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\sigma} Q}{P \xrightarrow{\sigma} Q}$	
<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$	<b>Par</b> $\frac{P \xrightarrow{\sigma} P' \quad bn(\sigma) \cup fn(Q) = \emptyset}{P Q \xrightarrow{\sigma} P' Q}$	
<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x}{(\nu z)P \xrightarrow{\bar{x}(z)} P'}$		

---

Table 4.2: Multi  $\pi$  late semantic with structural congruence

$$\begin{array}{c}
\text{LInp} \frac{}{x(b).P \xrightarrow{x(b)} P} \\
\text{SInp} \frac{}{x(a).x(b).P \xrightarrow{x(a).x(b)} P} \quad \text{Out} \frac{}{\bar{x}y.Q \xrightarrow{\bar{x}y} Q} \\
\text{LComSeq} \frac{}{x(a).x(b).P|\bar{x}y.Q \xrightarrow{x(b)} P\{y/a\}|Q} \\
\text{LCom} \frac{x(a).x(b).P|\bar{x}y.Q \xrightarrow{x(b)} P\{y/a\}|Q \quad \text{Out} \frac{}{\bar{x}z.R \xrightarrow{\bar{x}z} R}}{(x(a).x(b).P|\bar{x}y.Q)|\bar{x}z.R \xrightarrow{\tau} (P\{y/a\}|Q)\{z/b\}|R = (P\{y/a\}\{z/b\}|Q)|R}
\end{array}$$

### 4.2.3 Low level semantic

This section contains the definition of an alternative semantic for multi  $\pi$ . First we define a low level version of the multi  $\pi$  calculus (here with strong prefixing on input only), we call this language low multi  $\pi$ . The low multi  $\pi$  is the multi  $\pi$  enriched with a marked or intermediate process  $*P$ :

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A \mid *P$$

$$\pi ::= \bar{x}y \mid x(y) \mid \underline{x}(y) \mid \tau$$

**Definition 4.2.3.** The low level transition relation is the smallest relation induced by the rules in table 4.3 in which  $P$  stands for a process without mark,  $L$  stands for a process with mark and  $S$  can stand for both.

**Lemma 4.2.2.** For all unmarked processes  $P, Q$  and marked processes  $L_1, L_2$ .

- if  $P \xrightarrow{\alpha} L_1$  or  $L_1 \xrightarrow{\alpha} L_2$  then  $\alpha$  can only be an input or an  $\epsilon$
- if  $L_1 \xrightarrow{\alpha} P$  then  $\alpha$  is an input or a  $\tau$
- if  $P \xrightarrow{\alpha} Q$  then  $\alpha$  is not an  $\epsilon$

**Definition 4.2.4.** Let  $P, Q$  be unmarked processes and  $L_1, \dots, L_{k-1}$  marked processes. We define the derivation relation  $\rightarrow_s$  in the following way:

---

<b>Out</b> $\frac{}{\bar{x}y.P \mapsto P}$	<b>EInp</b> $\frac{}{x(y).P \mapsto P\{z/y\}}$	<b>Tau</b> $\frac{}{\tau.P \mapsto P}$
<b>StarInp</b> $\frac{P \mapsto S'}{*P \mapsto S'}$	<b>SInpLow</b> $\frac{}{\underline{x(z)}.P \mapsto *P\{y/z\}}$	<b>StarEps</b> $\frac{P \mapsto^\epsilon S'}{*P \mapsto^\epsilon S'}$
<b>Com1</b> $\frac{P \mapsto P' \quad Q \mapsto Q'}{P Q \mapsto P' Q'}$		
<b>Com2L</b> $\frac{L_1 \mapsto L_2 \quad P \mapsto Q}{L_1 P \mapsto L_2 Q}$	<b>Com2R</b> $\frac{P \mapsto Q \quad L_1 \mapsto L_2}{P L_1 \mapsto Q L_2}$	
<b>Com3L</b> $\frac{P \mapsto L \quad Q \mapsto Q'}{P Q \mapsto L Q'}$	<b>Com3R</b> $\frac{Q \mapsto Q' \quad P \mapsto L}{Q P \mapsto Q' L}$	
<b>Com4L</b> $\frac{L \mapsto P \quad Q \mapsto Q'}{L Q \mapsto P Q'}$	<b>Com4R</b> $\frac{Q \mapsto Q' \quad L \mapsto P}{L Q \mapsto P Q'}$	
<b>Res</b> $\frac{S \mapsto S' \quad y \notin n(\gamma)}{(\nu y)S \mapsto (\nu y)S'}$	<b>Opn</b> $\frac{P \mapsto Q \quad y \neq x}{(\nu y)P \mapsto Q}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \mapsto S}{P \mapsto S}$
<b>Par1L</b> $\frac{S \mapsto S'}{S Q \mapsto S' Q}$	<b>Par1R</b> $\frac{S \mapsto S'}{Q S \mapsto Q S'}$	<b>Sum</b> $\frac{P \mapsto S}{P + Q \mapsto S}$

---

Table 4.3: Low multi  $\pi$  early semantic with structural congruence

$$\mathbf{Low} \frac{P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} Q \quad k \geq 1}{P \xrightarrow{\gamma_1 \cdots \gamma_k} Q}$$

We need to be precise about the concatenation operator  $\cdot$  since we have introduced the new label  $\epsilon$ . Let  $a$  be an action such that  $a \neq \tau$  and  $a \neq \epsilon$  then the following rules hold:

$$\begin{aligned} \epsilon \cdot a &= a \cdot \epsilon = a & \epsilon \cdot \epsilon &= \epsilon & \tau \cdot \epsilon &= \epsilon \cdot \tau = \tau \\ \tau \cdot a &= a \cdot \tau = a & \tau \cdot \tau &= \tau \end{aligned}$$

**Example** Multi-party synchronization We show an example of a derivation of three processes that synchronize.

$$\begin{array}{c} \mathbf{SInpLow} \frac{}{\underline{x(a)}.x(b).P \xrightarrow{xy} *(x(b).P\{y/a\})} \quad \mathbf{Out} \frac{}{\overline{xy}.Q \xrightarrow{\overline{xy}} Q} \\ \mathbf{Com3L} \frac{}{\underline{x(a)}.x(b).P|\overline{xy}.Q \xrightarrow{\epsilon} *(x(b).P\{y/a\})|Q} \\ \mathbf{Par1L} \frac{}{(\underline{x(a)}.x(b).P|\overline{xy}.Q)|\overline{xz}.R \xrightarrow{\epsilon} (*(x(b).P\{y/a\})|Q)|\overline{xz}.R} \\ \\ \mathbf{EInp} \frac{}{x(b).P\{y/a\} \xrightarrow{xz} P\{y/a\}\{z/b\}} \\ \mathbf{Star} \frac{}{*(x(b).P\{y/a\}) \xrightarrow{xz} P\{y/a\}\{z/b\}} \\ \mathbf{Par1L} \frac{}{*(x(b).P\{y/a\})|Q \xrightarrow{xz} P\{y/a\}\{z/b\}|Q} \\ \\ \mathbf{Com4L} \frac{}{*(x(b).P\{y/a\})|Q \xrightarrow{xz} P\{y/a\}\{z/b\}|Q} \quad \mathbf{Out} \frac{}{\overline{xz}.R \xrightarrow{\overline{xz}} R} \\ \frac{}{(\underline{x(a)}.x(b).P|\overline{xy}.Q)|\overline{xz}.R \xrightarrow{\tau} (P\{y/a\}\{z/b\}|Q)|R} \end{array}$$

**Proposition 4.2.3.** Let  $\rightarrow$  be the relation defined in table 4.1. If  $P \xrightarrow{\sigma} Q$  then there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

*Proof.* The proof is by induction on the depth of the derivation tree of  $P \xrightarrow{\sigma} Q$ :

**base case**

If the depth is one then the rule used have to be one of: *EInp*, *Out*, *Tau*. These rules are also in table 4.3 so we can derive  $P \xrightarrow{\sigma} Q$ .

**inductive case**

If the depth is greater than one then the last rule used in the derivation can be:

*SInpSeq* : the last part of the derivation tree looks like this:

$$\mathbf{SInpSeq} \frac{P_1\{y/z\} \xrightarrow{\sigma} Q \quad |\sigma| > 1}{\underline{x(z)}.P_1 \xrightarrow{xy \cdot \sigma} Q}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1\{y/z\} \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

then a proof of the conclusion follows from:

$$\mathbf{SInpLow} \frac{}{\underline{x(z)}.P_1 \xrightarrow{xy} *P_1\{y/z\}} \quad \mathbf{Star} \frac{P_1\{y/z\} \xrightarrow{\gamma_1} L_1}{*P_1\{y/z\} \xrightarrow{\gamma_1} L_1}$$

where *Star* means *StarInp* or *StarEps*, note that  $\gamma_1$  is an input or an *epsilon* because of 4.2.1.

*SInp* : this case is similar to the previous.

*SInpTau* : this case is similar to the previous observing that  $xy \cdot \tau = xy$ .

*Sum* : the last part of the derivation tree looks like this:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\sigma} Q}{P_1 + P_2 \xrightarrow{\sigma} Q}$$

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \sigma$$

A proof of the conclusion is:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\gamma_1} L_1}{P_1 + P_2 \xrightarrow{\gamma_1} L_1}$$

*Cong* : this case is similar to the previous.

*ECom* : the last part of the derivation tree looks like this:

$$\mathbf{ECom} \frac{P_1 \xrightarrow{xy} P'_1 \quad Q_1 \xrightarrow{\bar{xy}} Q'_1}{P_1|Q_1 \xrightarrow{\tau} P'_1|Q'_1}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} P'_1 \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = xy$$

and there exist  $R_1, \dots, R_h$  and  $\delta_1, \dots, \delta_{h+1}$  with  $h \geq 0$  such that

$$Q_1 \xrightarrow{\delta_1} R_1 \xrightarrow{\delta_2} R_2 \cdots R_{h-1} \xrightarrow{\delta_h} R_h \xrightarrow{\delta_{h+1}} Q'_1 \quad \text{and} \quad \delta_1 \cdot \dots \cdot \delta_{h+1} = \bar{xy}$$

For lemma 4.2.2 there cannot be an output action in a transition involving marked processes so  $h$  must be 0 and  $Q_1 \xrightarrow{\delta_1} Q'_1$  with  $\delta_1 = \bar{xy}$ . We can have three different cases now:

$\gamma_1 = xy$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q'_1 \xrightarrow{\epsilon} L_2|Q'_1 \cdots \xrightarrow{\epsilon} L_k|Q'_1 \xrightarrow{\tau} P'_1|Q'_1$$

we derive the first transition with rule *Com3L*, whether for the other transition we use the rule *Par1L*.

$\gamma_i = xy$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1 \xrightarrow{\epsilon} L_{i+1}|Q'_1 \cdots \xrightarrow{\epsilon} L_k|Q'_1 \xrightarrow{\tau} P'_1|Q'_1$$

we derive the transaction  $L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1$  with rule *Com2L*, whether for the other transactions we use the rule *Par1L*.

$\gamma_{k+1} = xy$  similar.

*Res* : the last part of the derivation tree looks like this:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\sigma} Q_1 \quad z \notin n(\sigma)}{(\nu z)P_1 \xrightarrow{\sigma} (\nu z)Q_1}$$

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q_1 \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \sigma$$

We can apply the rule *Res* to each of the previous transitions because

$$z \notin n(\sigma) \text{ implies } z \notin n(\gamma_i) \text{ for each } i$$

and then get a proof of the conclusion:

$$(\nu z)P_1 \xrightarrow{\gamma_1} (\nu z)L_1 \xrightarrow{\gamma_2} (\nu z)L_2 \cdots (\nu z)L_{k-1} \xrightarrow{\gamma_k} (\nu z)L_k \xrightarrow{\gamma_{k+1}} (\nu z)Q_1$$

*Par* : this case is similar to the previous.

*EComSeq* : the last part of the derivation tree looks like this:

$$\mathbf{EComSeq} \frac{P_1 \xrightarrow{xy \cdot \sigma} P'_1 \quad Q_1 \xrightarrow{\bar{x}y} Q'_1}{P_1|Q_1 \xrightarrow{\sigma} P'_1|Q'_1}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} P'_1 \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = xy \cdot \sigma$$

For inductive hypothesis and lemma 4.2.2  $Q_1 \xrightarrow{\bar{x}y} Q'_1$ . We can have two different cases now depending on where the first  $xy$  is:

$\gamma_1 = xy$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q'_1 \xrightarrow{\gamma_2} L_2|Q'_1 \cdots \xrightarrow{\gamma_k} L_k|Q'_1 \xrightarrow{\gamma_{k+1}} P'_1|Q'_1$$

we derive the first transition with rule *Com3L*, whether for the other transactions we use the rule *Par1L*. Since  $\gamma_1 \cdot \dots \cdot \gamma_{k+1} = xy \cdot \sigma$  and  $\gamma_1 = xy$  then  $\epsilon \cdot \gamma_2 \cdot \dots \cdot \gamma_{k+1} = \sigma$

$\gamma_i = xy$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1 \xrightarrow{\gamma_{i+1}} L_{i+1}|Q'_1 \cdots \xrightarrow{\gamma_k} L_k|Q'_1 \xrightarrow{\gamma_{k+1}} P'_1|Q'_1$$

we derive the transition  $L_{i-1}|Q_1 \xrightarrow{\epsilon} L_i|Q'_1$  with rule *Com2L*, whether for the other transactions of the premises we use the rule *Par1L*.

$\gamma_{k+1} = xy$  : cannot happen because  $\sigma$  is not empty.

□

**Proposition 4.2.4.** Let  $\rightarrow$  be the relation defined in table 4.1. Let  $\alpha$  be an action. If  $P \xrightarrow{\alpha} Q$  then  $P \xrightarrow{\alpha} Q$ .

*Proof.* The proof is by induction the depth of the derivation of  $P \xrightarrow{\alpha} Q$ :

**base case** in this case the derivation of this transition has depth one. The last(and only) rule used can be: *Out*, *EInp* or *Tau*; these rules are also in table 4.1 so we can derive  $P \xrightarrow{\alpha} Q$ .

**inductive case** in this case the last rule in the derivation can be: *Sum*, *Com1*, *Res*, *Par1L*, *Par1R*, *Cong*, *Opn*:

*Com1* :

$$\mathbf{Com1} \frac{P_1 \xrightarrow{xy} Q_1 \quad P_2 \xrightarrow{\bar{x}y} Q_2}{P_1|P_2 \xrightarrow{\tau} Q_1|Q_2}$$

for inductive hypothesis  $P_1 \xrightarrow{xy} Q_1$  and  $P_2 \xrightarrow{\bar{x}y} Q_2$  so for rule *Com*  $P_1|P_2 \xrightarrow{\tau} Q_1|Q_2$

*Sum* :

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\alpha} Q}{P_1 + P_2 \xrightarrow{\alpha} Q}$$

---

<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$	<b>LInp</b> $\frac{w \notin (fn(P) - \{z\})}{x(z).P \xrightarrow{x(w)} P\{w/z\}}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$
<b>SInp</b> $\frac{P\{w/z\} \xrightarrow{\sigma} P' \quad w \notin (fn(P) - \{z\})}{x(y).P \xrightarrow{x(w) \cdot \sigma} P'}$	$\sigma$ is a non empty sequence of inputs	
<b>LComSeq</b> $\frac{P \xrightarrow{x(y) \cdot \sigma} P' \quad Q \xrightarrow{\bar{x}z} Q' \quad bn(\sigma) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\sigma\{z/y\}} P'\{z/y\} Q'}$	<b>LCom</b> $\frac{P \xrightarrow{x(y)} P' \quad Q \xrightarrow{\bar{x}z} Q'}{P Q \xrightarrow{\tau} P'\{z/y\} Q'}$	
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Opn</b> $\frac{P \xrightarrow{\bar{x}z} P' \quad z \neq x \quad w \notin fn(P' - \{y\})}{(\nu z)P \xrightarrow{\bar{x}(z)} P'\{z/w\}}$	
<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$	<b>Par</b> $\frac{P \xrightarrow{\sigma} P' \quad bn(\sigma) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\sigma} P' Q}$	
<b>Close</b> $\frac{P \xrightarrow{x(w)} P' \quad Q \xrightarrow{\bar{x}(w)} Q'}{P Q \xrightarrow{\tau} (\nu w)(P' Q')}$	<b>CloseSeq</b> $\frac{P \xrightarrow{x(w) \cdot \sigma} P' \quad Q \xrightarrow{\bar{x}(w)} Q'}{P Q \xrightarrow{\sigma} (\nu w)(P' Q')}$	

---

Table 4.4: Multi  $\pi$  late semantic without structural congruence

for inductive hypothesis  $P_1 \xrightarrow{\alpha} Q$  and for rule *Sum*  $P_1 + P_2 \xrightarrow{\alpha} Q$ .

*Res* the first transition is:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\alpha} Q_1 \quad z \notin n(\gamma_1)}{(\nu z)P_1 \xrightarrow{\alpha} (\nu z)Q_1}$$

for inductive hypothesis  $P_1 \xrightarrow{\alpha} Q_1$  and for rule *Res*  $(\nu z)P_1 \xrightarrow{\alpha} (\nu z)Q_1$ .

*others* : other cases are similar.

□

#### 4.2.4 Late operational semantic without structural congruence

**Definition 4.2.5.** The *late transition relation without structural congruence* is the smallest relation induced by the rules in table 4.4.

**Example** Multi-party synchronization We show an example of a derivation of three processes that synchronize with the late semantic. The three processes are  $\underline{x(a)}.x(b).P$ ,  $\bar{x}y.Q$  and  $\bar{x}z.R$ . We assume that:

$$\begin{array}{c}
 b \notin fn(\underline{x(a)}.x(b).P|\bar{x}y.Q) \\
 \\
 \mathbf{LCom} \frac{\underline{x(a)}.x(b).P|\bar{x}y.Q \xrightarrow{x(b)} P\{y/a\}|Q \quad \mathbf{Out} \frac{}{\bar{x}z.R \xrightarrow{\bar{x}z} R}}{(\underline{x(a)}.x(b).P|\bar{x}y.Q)|\bar{x}z.R \xrightarrow{\tau} (P\{y/a\}|Q)\{z/b\}|R} \\
 \\
 \mathbf{LInp} \frac{}{x(b).P \xrightarrow{x(b)} P} \\
 \mathbf{SInp} \frac{}{x(a).x(b).P \xrightarrow{x(a) \cdot x(b)} P} \\
 \mathbf{LComSeq} \frac{x(a).x(b).P \xrightarrow{x(a) \cdot x(b)} P \quad \mathbf{Out} \frac{}{\bar{x}y.Q \xrightarrow{\bar{x}y} Q}}{\underline{x(a)}.x(b).P|\bar{x}y.Q \xrightarrow{x(b)} P\{y/a\}|Q}
 \end{array}$$

In the following example we assume that all bound variables have different names.

**Example** Scope intrusion without strong prefixing.

$$\bar{y}x.P|(\nu x)(y(z).Q) \xrightarrow{\tau} P|(\nu w)(Q\{w/x\}\{x/z\})$$

This cannot be proved without explicit  $\alpha$  conversion.

**Example** Scope intrusion with strong prefixing.

$$(\nu x)(\nu b)\underline{y(z)}.a(c).P|y(x).Q|a(b).R \xrightarrow{\tau} ((\nu x')(\nu b')P\{x'/x\}\{x/z\}|Q)\{b'/b\}\{b/c\}|R$$

This cannot be proved without explicit  $\alpha$  conversion.

**Example** Scope extrusion without strong prefixing.  $x \notin fn(Q)$

$$\text{Cls} \frac{\text{Opn} \frac{\text{Out} \frac{\bar{y}x.P \xrightarrow{\bar{y}x} P}{(\nu x)(\bar{y}x.P) \xrightarrow{\bar{y}(x)} P}}{(\nu x)(\bar{y}x.P)|y(z).Q \xrightarrow{\tau} (\nu x)(P|Q\{x/z\})} \quad \text{LInp} \frac{y(z).Q \xrightarrow{y(x)} Q\{x/z\}}{(\nu x)(\bar{y}x.P)|y(z).Q \xrightarrow{\tau} (\nu x)(P|Q\{x/z\})}}$$

**Example** Scope extrusion with strong prefixing.

$$\begin{aligned} & \underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R|(\nu z_2)\bar{y}z_2.S \xrightarrow{\tau} (\nu z_2)((\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R|S) \\ & \text{ClsSeq} \frac{\text{SInp} \frac{\text{SInp} \frac{\text{LInp} \frac{y(x_2).P\{x_1/z_1\} \xrightarrow{y(z_2)} P\{x_1/z_1\}\{x_2/z_2\}}{(a(b).y(x_2).P)\{x_1/z_1\} \xrightarrow{a(b).y(z_2)} P\{x_1/z_1\}\{x_2/z_2\}}}{y(x_1).a(b).y(x_2).P \xrightarrow{y(z_1).a(b).y(z_2)} P\{x_1/z_1\}\{x_2/z_2\}}}{(\nu z_1)\bar{y}z_1.Q \xrightarrow{y(z_1)} Q}}{y(x_1).a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q \xrightarrow{a(b).y(x_2)} (\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)} \\ & \text{ClsSeq} \frac{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q \xrightarrow{a(b).y(z_2)} (\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)}{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R \xrightarrow{y(z_2)} (\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R}} \quad \text{Out} \frac{\bar{a}b.R \xrightarrow{\bar{a}b} R}{\bar{a}b.R \xrightarrow{\bar{a}b} R} \\ & \text{ClsSeq} \frac{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R \xrightarrow{y(z_2)} (\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R}}{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R \xrightarrow{y(z_2)} (\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R}} \quad \text{Out} \frac{\bar{y}z_2.S \xrightarrow{\bar{y}z_2} S}{\bar{y}z_2.S \xrightarrow{\bar{y}z_2} S} \\ & \text{Opn} \frac{(\nu z_2)\bar{y}z_2.S \xrightarrow{\bar{y}(z_2)} S}{(\nu z_2)\bar{y}z_2.S \xrightarrow{\bar{y}(z_2)} S} \\ & \text{Cls} \frac{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R|(\nu z_2)\bar{y}z_2.S \xrightarrow{\tau} (\nu z_2)((\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R|S)}}{\underline{y(x_1)}.a(b).y(x_2).P|(\nu z_1)\bar{y}z_1.Q|\bar{a}b.R|(\nu z_2)\bar{y}z_2.S \xrightarrow{\tau} (\nu z_2)((\nu z_1)(P\{z_1/x_1\}\{x_2/z_2\}|Q)|R|S)}} \end{aligned}$$





## Chapter 5

# Multi $\pi$ calculus with strong input and output

### 5.1 Syntax

As we did with multi  $\pi$  calculus, we suppose that we have a countable set of names  $\mathbb{N}$ , ranged over by lower case letters  $a, b, \dots, z$ . These names are used for communication channels and values. Furthermore we have a set of identifiers, ranged over by  $A$ . We represent the agents or processes by upper case letters  $P, Q, \dots$ . A multi  $\pi$  process, in addition to the same actions of a  $\pi$  process, can perform also a strong prefix:

$$\pi ::= \bar{x}y \mid x(z) \mid \underline{x}(y) \mid \bar{x}y \mid \tau$$

The process are defined, just as original  $\pi$  calculus, by the following grammar:

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A(y_1, \dots, y_n)$$

and they have the same intuitive meaning as for the  $\pi$  calculus. The strong prefix input allows a process to make an atomic sequence of actions, so that more than one process can synchronize on this sequence.

We have to extend the following definition to deal with the strong prefix:

$$\begin{aligned} B(x(y).Q, I) &= \{y, \bar{y}\} \cup B(Q, I) & F(x(y).Q, I) &= \{x, \bar{x}\} \cup (F(Q, I) - \{y, \bar{y}\}) \\ B(\bar{x}y.Q, I) &= B(Q, I) & F(\bar{x}y.Q, I) &= \{x, \bar{x}, y, \bar{y}\} \cup F(Q, I) \end{aligned}$$

### 5.2 Operational semantic

#### 5.2.1 Early operational semantic with structural congruence

**Definition 5.2.1.** The *early transition relation with structural congruence* is the smallest relation induced by the rules in table 5.1:

The names  $\sigma, \sigma_1, \sigma_2, \sigma_3$  are non empty sequences of actions and are also not  $\tau$ . The relation  $ESync$  is defined by the axioms in table 5.2

**Example Transactional synchronization.** This is an example of two processes that synchronize over a sequence of actions of length two:

$$\bar{a}x.\bar{a}y.P|a(w).a(z).Q \xrightarrow{\tau} P|Q\{x/w\}\{y/z\}$$

We start first noticing that

$$\text{S4R} \frac{\text{S1R} \frac{}{Sync(\bar{a}y, ay, \tau)}}{Sync(\bar{a}x \cdot \bar{a}y, ax \cdot ay, \tau)}$$

and that

---

<b>Inp</b> $\frac{}{x(y).P \xrightarrow{xz} P\{z/x\}}$	<b>Tau</b> $\frac{}{\tau.P \xrightarrow{\tau} P}$	<b>Out</b> $\frac{}{\bar{x}y.P \xrightarrow{\bar{x}y} P}$
<b>SInp</b> $\frac{\{z/y\}P \xrightarrow{\sigma} P' \quad \sigma \neq \tau}{x(y).P \xrightarrow{xz \cdot \sigma} P'}$	<b>SOut</b> $\frac{P \xrightarrow{\sigma} P' \quad \sigma \neq \tau}{\bar{x}y.P \xrightarrow{\bar{x}y \cdot \sigma} P'}$	
<b>ECom</b> $\frac{P \xrightarrow{\sigma_1} P' \quad Q \xrightarrow{\sigma_2} Q' \quad ESync(\sigma_1, \sigma_2, \sigma_3)}{P Q \xrightarrow{\sigma_3} P' Q'}$		
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\alpha} Q}{P \xrightarrow{\alpha} Q}$	
<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$	<b>Par</b> $\frac{P \xrightarrow{\sigma} P'}{P Q \xrightarrow{\sigma} P' Q}$	

---

Table 5.1: Multi  $\pi$  early semantic with structural congruence

---

<b>S1L</b> $\frac{}{ESync(xy, \bar{x}y, \tau)}$	<b>S1R</b> $\frac{}{ESync(\bar{x}y, xy, \tau)}$
<b>S2L</b> $\frac{}{ESync(xy, \bar{x}y \cdot \sigma, \sigma)}$	<b>S2R</b> $\frac{}{ESync(\bar{x}y \cdot \sigma, xy, \sigma)}$
<b>S3L</b> $\frac{}{ESync(xy \cdot \sigma, \bar{x}y, \sigma)}$	<b>S3R</b> $\frac{}{ESync(\bar{x}y, xy \cdot \sigma, \sigma)}$
<b>S4L</b> $\frac{ESync(\sigma_1, \sigma_2, \sigma_3)}{ESync(xy \cdot \sigma_1, \bar{x}y \cdot \sigma_2, \sigma_3)}$	<b>S4R</b> $\frac{ESync(\sigma_1, \sigma_2, \sigma_3)}{ESync(\bar{x}y \cdot \sigma_1, xy \cdot \sigma_2, \sigma_3)}$

---

Table 5.2: Synchronization relation

$$\text{SOUT} \frac{\text{OUT} \frac{}{\bar{a}y.P \xrightarrow{\bar{a}y} P}}{\bar{a}x.\bar{a}y.P \xrightarrow{\bar{a}x.\bar{a}y} P} \quad \text{SINP} \frac{\text{INP} \frac{}{(a(z).Q)\{x/w\} \xrightarrow{ay} Q\{x/w\}\{y/z\}}}{a(w).a(z).Q \xrightarrow{ax.ay} Q}$$

and in the end we just need to apply the rule **LCom**

**Example Multi-party synchronization.** In this example we have three processes that want to synchronize:

$$\begin{array}{c} \text{ECom} \frac{\bar{a}f.\bar{b}g.P|a(w).Q \xrightarrow{\bar{b}g} P|Q\{f/w\} \quad \text{Inp} \frac{}{b(y).R \xrightarrow{bg} R\{g/y\}} \quad \text{S1R} \frac{}{\text{Sync}(\bar{b}g, bg, \tau)}}{(\bar{a}f.\bar{b}g.P|a(w).Q)|b(y).R \xrightarrow{\tau} (P|Q\{f/w\})|R\{g/y\}} \\ \\ \text{LCom} \frac{\bar{a}f.\bar{b}g.P \xrightarrow{\bar{a}f.\bar{b}g} P \quad \text{Inp} \frac{}{a(w).Q \xrightarrow{af} Q\{f/w\}} \quad \text{S2R} \frac{}{\text{Sync}(\bar{a}f \cdot \bar{b}g, af, \bar{b}g)}}{\bar{a}f.\bar{b}g.P|a(w).Q \xrightarrow{\bar{b}g} P|Q\{f/w\}} \\ \\ \text{SOut} \frac{\text{Out} \frac{}{\bar{b}g.P \xrightarrow{\bar{b}g} P}}{\bar{a}f.\bar{b}g.P \xrightarrow{\bar{a}f.\bar{b}g} P} \end{array}$$

### 5.2.2 Late operational semantic with structural congruence

The semantic of a multi  $\pi$  process is labeled transition system such that

- the nodes are multi  $\pi$  calculus process. The set of node is  $\mathbb{P}_m$
- The set of actions is  $\mathbb{A}_m$  and can contain
  - bound output  $\bar{x}(y)$
  - unbound output  $\bar{x}y$
  - bound input  $x(z)$

We use  $\alpha, \alpha_1, \alpha_2, \dots$  to range over the set of actions, we use  $\sigma, \sigma_1, \sigma_2, \dots$  to range over the set  $\mathbb{A}_m^+ \cup \{\tau\}$ .

- the transition relations is  $\rightarrow \subseteq \mathbb{P}_m \times (\mathbb{A}_m^+ \cup \{\tau\}) \times \mathbb{P}_m$

In this case, a label can be a sequence of prefixes, whether in the original  $\pi$  calculus a label can be only a prefix. We use the symbol  $\cdot$  to denote the concatenation operator.

**Definition 5.2.2.** The *late transition relation with structural congruence* is the smallest relation induced by the rules in table 5.3:

In what follows, the names  $\delta, \delta_1, \delta_2$  represents substitutions, they can also be empty; the names  $\sigma, \sigma_1, \sigma_2, \sigma_3$  are non empty sequences of actions. The relation *Sync* is defined by the axioms in table 5.4

**Example Transactional synchronization.** This is an example of two processes that synchronize over a sequence of actions of length two:

$$\bar{a}x.\bar{a}y.P|a(w).a(z).Q \xrightarrow{\tau} P|Q\{x/w\}\{y/z\}$$

We start first noticing that

$$\text{S4R} \frac{\text{S1R} \frac{}{\text{Sync}(\bar{a}y, a(z)\{x/w\}, \tau, \{y/z\})}}{\text{Sync}(\bar{a}x \cdot \bar{a}y, a(w) \cdot a(z), \tau, \{x/w\}\{y/z\})}$$

and that

---

<b>Pref</b> $\frac{\alpha \text{ not a strong prefix}}{\alpha.P \xrightarrow{\alpha} P}$	<b>Par</b> $\frac{P \xrightarrow{\sigma} P' \quad bn(\sigma) \cap fn(Q) = \emptyset}{P Q \xrightarrow{\sigma} P' Q}$
<b>SOut</b> $\frac{P \xrightarrow{\sigma} P' \quad \sigma \neq \tau}{\bar{x}y.P \xrightarrow{\bar{x}y \cdot \sigma} P'}$	<b>LCom</b> $\frac{P \xrightarrow{\sigma_1} P' \quad Q \xrightarrow{\sigma_2} Q' \quad Sync(\sigma_1, \sigma_2, \sigma_3, \delta_1, \delta_2)}{P Q \xrightarrow{\sigma_3} P'\delta_1 Q'\delta_2}$
<b>Sum</b> $\frac{P \xrightarrow{\sigma} P'}{P + Q \xrightarrow{\sigma} P'}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\alpha} Q}{P \xrightarrow{\alpha} Q}$
<b>Res</b> $\frac{P \xrightarrow{\sigma} P' \quad z \notin n(\alpha)}{(\nu z)P \xrightarrow{\sigma} (\nu z)P'}$	<b>SInp</b> $\frac{P \xrightarrow{\sigma} P' \quad \sigma \neq \tau}{\underline{x}(y).P \xrightarrow{x(y) \cdot \sigma} P'}$

---

Table 5.3: Multi  $\pi$  late semantic with structural congruence

---

S1L $\frac{}{Sync(x(y), \bar{x}z, \tau, \{z/y\}, \{\})}$	S1R $\frac{}{Sync(\bar{x}z, x(y), \tau, \{\}, \{z/y\})}$
S2L $\frac{}{Sync(x(y), \bar{x}z \cdot \sigma, \sigma, \{z/y\}, \{\})}$	S2R $\frac{}{Sync(\bar{x}z \cdot \sigma, x(y), \sigma, \{\}, \{z/y\})}$
S3L $\frac{}{Sync(x(y) \cdot \sigma, \bar{x}z, \sigma\{z/y\}, \{z/y\}, \{\})}$	S3R $\frac{}{Sync(\bar{x}z, x(y) \cdot \sigma, \sigma\{z/y\}, \{\}, \{z/y\})}$
S4L $\frac{Sync(\sigma_1, \sigma_2\{z/y\}, \sigma_3, \delta_1, \delta_2)}{Sync(x(y) \cdot \sigma_1, \bar{x}z \cdot \sigma_2, \sigma_3, \{z/y\}\delta_1, \delta_2)}$	S4R $\frac{Sync(\sigma_1, \sigma_2\{z/y\}, \sigma_3, \delta_1, \delta_2)}{Sync(\bar{x}z \cdot \sigma_1, x(y) \cdot \sigma_2, \sigma_3, \delta_1, \{z/y\}\delta_2)}$

---

Table 5.4: Synchronization relation

$$\text{SOUT} \frac{\text{PREF} \frac{\overline{a}y.P \xrightarrow{\overline{a}y} P}}{\overline{a}x.\overline{a}y.P \xrightarrow{\overline{a}x.\overline{a}y} P} \quad \text{SINP} \frac{\text{PREF} \frac{a(z).Q \xrightarrow{a(z)} Q}}{a(w).a(z).Q \xrightarrow{a(w).a(z)} Q}$$

and in the end we just need to apply the rule **LCom**


**Example Multi-party synchronization.** In this example we have three processes that want to synchronize:

$$\begin{array}{c} \text{LCom} \frac{\overline{a}f.\overline{b}g.P|a(w).Q \xrightarrow{\overline{b}g} P|Q\{f/w\} \quad \text{Pref} \frac{}{b(y).R \xrightarrow{b(y)} R} \quad \text{S1R} \frac{}{\text{Sync}(\overline{b}g, b(y), \tau, \emptyset, \{g/y\})}}{(\overline{a}f.\overline{b}g.P|a(w).Q)|b(y).R \xrightarrow{\tau} (P|Q\{f/w\})|R\{g/y\}} \\ \\ \text{LCom} \frac{\overline{a}f.\overline{b}g.P \xrightarrow{\overline{a}f.\overline{b}g} P \quad \text{Pref} \frac{}{a(w).Q \xrightarrow{a(w)} Q} \quad \text{S2R} \frac{}{\text{Sync}(\overline{a}f.\overline{b}g, a(w), \tau, \emptyset, \{f/w\})}}{\overline{a}f.\overline{b}g.P|a(w).Q \xrightarrow{\overline{b}g} P|Q\{f/w\}} \\ \\ \text{SOut} \frac{\text{Out} \frac{}{\overline{b}g.P \xrightarrow{\overline{b}g} P}}{\overline{a}f.\overline{b}g.P \xrightarrow{\overline{a}f.\overline{b}g} P} \end{array}$$

**Example Cigarette smokers' problem.** In this problem there are four processes: an agent and three smokers. Each smoker continuously makes a cigarette and smokes it. To make a cigarette each smoker needs three ingredients: tobacco, paper and matches. One of the smokers has paper, another tobacco and the third matches. The agent has an infinite supply of the ingredients. The agent places two of the ingredients on the table. The smoker who has the remaining ingredient take the others from the table, make a cigarette and smokes. Then the cycle repeats. A solution to the problem is the following:

$$\begin{aligned} \text{Agent} &\stackrel{\text{def}}{=} \overline{\text{tob}}.\overline{\text{mat}}.\text{end}().\text{Agent} + \overline{\text{mat}}.\overline{\text{pap}}.\text{end}().\text{Agent} + \overline{\text{pap}}.\overline{\text{tob}}.\text{end}().\text{Agent} \\ S_{\text{pap}} &\stackrel{\text{def}}{=} \overline{\text{tob}}().\text{mat}().\overline{\text{smoke}}.\text{end}.S_{\text{pap}} \\ S_{\text{tab}} &\stackrel{\text{def}}{=} \overline{\text{mat}}().\text{pap}().\overline{\text{smoke}}.\text{end}.S_{\text{tab}} \\ S_{\text{mat}} &\stackrel{\text{def}}{=} \overline{\text{pap}}().\text{tob}().\overline{\text{smoke}}.\text{end}.S_{\text{mat}} \\ \text{CSP} &\stackrel{\text{def}}{=} (\nu \text{tob}, \text{pap}, \text{mat}, \text{end})(\text{Agent}|S_{\text{tob}}|S_{\text{mat}}|S_{\text{pap}}) \end{aligned}$$

The semantic of *CSP* is the following graph:



where

$$\begin{aligned}
PAP &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|S_{tob}|S_{mat}|\overline{smoke}.end.S_{pap}) \\
TOB &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|\overline{smoke}.end.S_{tob}|S_{mat}|S_{pap}) \\
MAT &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|S_{tob}|\overline{smoke}.end.S_{mat}|S_{pap}) \\
PAP_1 &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|S_{tob}|S_{mat}|\overline{end}.S_{pap}) \\
TOB_1 &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|\overline{end}.S_{tob}|S_{mat}|S_{pap}) \\
MAT_1 &\stackrel{def}{=} (\nu tob, pap, mat, end)(end().Agent|S_{tob}|\overline{end}.S_{mat}|S_{pap})
\end{aligned}$$

### 5.2.3 Low level semantic

This section contains the definition of an alternative semantic for multi  $\pi$ . First we define a low level version of the multi  $\pi$  calculus, we call this language low multi  $\pi$ . The low multi  $\pi$  is the multi  $\pi$  enriched with a marked or intermediate process  $*P$ :

$$P, Q ::= 0 \mid \pi.P \mid P|Q \mid P + Q \mid (\nu x)P \mid A(x_1, \dots, x_n) \mid *P$$

$$\pi ::= \bar{x}y \mid x(y) \mid \bar{x}y \mid x(y) \mid \tau$$

**Definition 5.2.3.** The low level transition relation is the smallest relation induced by the rules in table 5.5 in which  $P$  stands for a process without mark,  $L$  stands for a process with mark and  $S$  can stand for both.

**Proposition 5.2.1.** Let  $\rightarrow$  be the relation defined in table 5.1. If  $P \xrightarrow{\sigma} Q$  then there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \sigma$$

*Proof.* The proof is by induction on the depth of the derivation tree of  $P \xrightarrow{\sigma} Q$ :

**base case**

If the depth is one then the rule used have to be one of: *EInp*, *Out*, *Tau*. These rules are also in table 5.5 so we can derive  $P \xrightarrow{\sigma} Q$ .

**inductive case**

If the depth is greater than one then the last rule used in the derivation can be:

*SOut* : the last part of the derivation tree looks like this:

$$\mathbf{SOut} \frac{P_1 \xrightarrow{\sigma} Q \quad \sigma \neq \tau}{\bar{x}y.P_1 \xrightarrow{\bar{x}y.\sigma} Q}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdot \dots \cdot \gamma_{k+1} = \sigma$$

then a proof of the conclusion follows from:

$$\mathbf{SOutLow} \frac{}{\bar{x}y.P_1 \xrightarrow{\bar{x}y} *P_1} \quad \mathbf{Star} \frac{P_1 \xrightarrow{\gamma_1} L_1}{*P_1 \xrightarrow{\gamma_1} L_1}$$

*SInp* : this case is similar to the previous.

*Sum* : the last part of the derivation tree looks like this:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\sigma} Q}{P_1 + P_2 \xrightarrow{\sigma} Q}$$

---

<b>Out</b> $\frac{}{\bar{x}y.P \mapsto P}$	<b>EInp</b> $\frac{}{x(y).P \mapsto P\{z/y\}}$	<b>Tau</b> $\frac{}{\tau.P \mapsto P}$
<b>SOutLow</b> $\frac{}{\bar{x}y.P \mapsto *P}$	<b>SInpLow</b> $\frac{}{x(y).P \mapsto *P\{z/y\}}$	
<b>StarEps</b> $\frac{S \xrightarrow{\epsilon} S'}{*S \xrightarrow{\epsilon} S'}$	<b>StarInp</b> $\frac{S \xrightarrow{xy} S'}{*S \xrightarrow{xy} S'}$	<b>StarOut</b> $\frac{S \xrightarrow{\bar{xy}} S'}{*S \xrightarrow{\bar{xy}} S'}$
<b>Par1R</b> $\frac{S \xrightarrow{\gamma} S'}{Q S \xrightarrow{\gamma} Q S'}$	<b>Par1L</b> $\frac{S \xrightarrow{\gamma} S'}{S Q \xrightarrow{\gamma} S' Q}$	
<b>Sum</b> $\frac{P \xrightarrow{\gamma} S}{P+Q \xrightarrow{\gamma} S}$	<b>Cong</b> $\frac{P \equiv P' \quad P' \xrightarrow{\gamma} S}{P \xrightarrow{\gamma} S}$	<b>Res</b> $\frac{S \xrightarrow{\gamma} S' \quad y \notin n(\gamma)}{(\nu y)S \xrightarrow{\gamma} (\nu y)S'}$
<b>Com1</b> $\frac{P \xrightarrow{\bar{xy}} P' \quad Q \xrightarrow{xy} Q'}{P Q \xrightarrow{\tau} P' Q'}$		
<b>Com2LOut</b> $\frac{L_1 \xrightarrow{\bar{xy}} L'_1 \quad L_2 \xrightarrow{xy} S}{L_1 L_2 \xrightarrow{\epsilon} L'_1 S}$	<b>Com2ROut</b> $\frac{L_1 \xrightarrow{xy} S \quad L_2 \xrightarrow{\bar{xy}} L'_2}{L_1 L_2 \xrightarrow{\epsilon} S L'_2}$	
<b>Com2LInp</b> $\frac{L_1 \xrightarrow{\bar{xy}} S \quad L_2 \xrightarrow{xy} L'_2}{L_1 L_2 \xrightarrow{\epsilon} S L'_2}$	<b>Com2RInp</b> $\frac{L_1 \xrightarrow{xy} L'_1 \quad L_2 \xrightarrow{\bar{xy}} S}{L_1 L_2 \xrightarrow{\epsilon} L'_1 S}$	
<b>Com3LOut</b> $\frac{Q \xrightarrow{\bar{xy}} S \quad P \xrightarrow{xy} L}{Q P \xrightarrow{\epsilon} S L}$	<b>Com3ROut</b> $\frac{P \xrightarrow{xy} L \quad Q \xrightarrow{\bar{xy}} S}{P Q \xrightarrow{\epsilon} L S}$	
<b>Com3LInp</b> $\frac{Q \xrightarrow{xy} S \quad P \xrightarrow{\bar{xy}} L}{Q P \xrightarrow{\epsilon} S L}$	<b>Com3RInp</b> $\frac{P \xrightarrow{\bar{xy}} L \quad Q \xrightarrow{xy} S}{P Q \xrightarrow{\epsilon} L S}$	
<b>Com4L</b> $\frac{L_1 \xrightarrow{\bar{xy}} P \quad L_2 \xrightarrow{xy} Q}{L_1 L_2 \xrightarrow{\tau} P Q}$	<b>Com4R</b> $\frac{L_1 \xrightarrow{xy} P \quad L_2 \xrightarrow{\bar{xy}} Q}{L_1 L_2 \xrightarrow{\tau} P Q}$	

---

Table 5.5: Low multi  $\pi$  early semantic with structural congruence

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

A proof of the conclusion is:

$$\mathbf{Sum} \frac{P_1 \xrightarrow{\gamma_1} L_1}{P_1 + P_2 \xrightarrow{\gamma_1} L_1}$$

*Cong* : this case is similar to the previous.

*Res* : the last part of the derivation tree looks like this:

$$\mathbf{Res} \frac{P_1 \xrightarrow{\sigma} Q_1 \quad z \notin n(\sigma)}{(\nu z)P_1 \xrightarrow{\sigma} (\nu z)Q_1}$$

for the inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} Q_1 \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma$$

We can apply the rule *Res* to each of the previous transitions because

$$z \notin n(\sigma) \text{ implies } z \notin n(\gamma_i) \text{ for each } i$$

and then get a proof of the conclusion:

$$(\nu z)P_1 \xrightarrow{\gamma_1} (\nu z)L_1 \xrightarrow{\gamma_2} (\nu z)L_2 \cdots (\nu z)L_{k-1} \xrightarrow{\gamma_k} (\nu z)L_k \xrightarrow{\gamma_{k+1}} (\nu z)Q_1$$

*Par* : this case is similar to the previous.

*ECom* : the last part of the derivation tree looks like this:

$$\mathbf{ECom} \frac{P_1 \xrightarrow{\sigma_1} P'_1 \quad Q_1 \xrightarrow{\sigma_2} Q'_1 \quad ESync(\sigma_1, \sigma_2, \sigma_3)}{P_1|Q_1 \xrightarrow{\sigma_3} P'_1|Q'_1}$$

for inductive hypothesis there exist  $L_1, \dots, L_k$  and  $\gamma_1, \dots, \gamma_{k+1}$  with  $k \geq 0$  such that

$$P_1 \xrightarrow{\gamma_1} L_1 \xrightarrow{\gamma_2} L_2 \cdots L_{k-1} \xrightarrow{\gamma_k} L_k \xrightarrow{\gamma_{k+1}} P'_1 \quad \text{and} \quad \gamma_1 \cdots \gamma_{k+1} = \sigma_1$$

and there exist  $R_1, \dots, R_h$  and  $\delta_1, \dots, \delta_{h+1}$  with  $h \geq 0$  such that

$$Q_1 \xrightarrow{\delta_1} R_1 \xrightarrow{\delta_2} R_2 \cdots R_{h-1} \xrightarrow{\delta_h} R_h \xrightarrow{\delta_{h+1}} Q'_1 \quad \text{and} \quad \delta_1 \cdots \delta_{h+1} = \sigma_2$$

We proceed by cases on the derivation of  $ESync(\sigma_1, \sigma_2, \sigma_3)$ . We show just some cases because the others are similar.

*S1L* Suppose that  $\delta_1$  is  $\bar{x}y$  (the other cases are similar), so the other  $\delta$ s are  $\epsilon$  or  $\tau$ . We can have three different cases now each :

$\gamma_1 = xy$  : The other  $\gamma$ s are  $\epsilon$  or  $\tau$ . A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|R_1 \xrightarrow{\epsilon} L_2|R_1 \cdots \xrightarrow{\epsilon} P'_1|R_1 \xrightarrow{\epsilon} P'_1|R_2 \cdots \xrightarrow{\epsilon} P'_1|Q'_1$$

we derive the first transition with rule *Com3ROut*, whether for the other transition we use the rules *Par1L*, *Par1R*, *Par3L* or *Par3R*.

$\gamma_i = xy$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\tau} L_i|R_1 \xrightarrow{\epsilon} L_{i+1}|R_1 \cdots \xrightarrow{\epsilon} P'_1|R_1 \xrightarrow{\epsilon} P'_1|R_2 \cdots \xrightarrow{\epsilon} P'_1|Q'_1$$



we derive the transaction  $L_{i-1}|Q_1 \xrightarrow{\tau} L_i|R_1$  with rule *Com5L*, whether for the other transactions we use some rule for parallel.

$\gamma_{k+1} = xy$  similar.

*S2R* : We suppose that  $\delta_1 = xy$  and so other  $\delta$ s are  $\epsilon$  or  $\tau$ , the other cases are similar. We can have two different cases now depending on where the first  $\bar{xy}$  is:

$\gamma_1 = \bar{xy}$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\tau} L_1|R_1 \xrightarrow{\gamma_2} L_2|R_1 \cdots \xrightarrow{\gamma_{k+1}} P'_1|R_1 \xrightarrow{\delta_2} P'_1|R_2 \cdots \xrightarrow{\delta_{h+1}} P'_1|Q'_1$$

we derive the first transition with rule *Com3L*, whether for the other transactions we use some rule for parallel. Since  $\gamma_1 \cdots \gamma_{k+1} = \bar{xy} \cdot \sigma$  and  $\gamma_1 = \bar{xy}$  then  $\tau \cdot \gamma_2 \cdots \gamma_{k+1} \cdot \epsilon \cdots \epsilon \cdot \tau = \sigma$

$\gamma_i = \bar{xy}$  : A proof of the conclusion is:

$$P_1|Q_1 \xrightarrow{\epsilon} L_1|Q_1 \cdots \xrightarrow{\epsilon} L_{i-1}|Q_1 \xrightarrow{\tau} L_i|R_1 \xrightarrow{\gamma_{i+1}} L_{i+1}|R_1 \cdots \xrightarrow{\gamma_k} P'_1|R_1 \xrightarrow{\delta_2} P'_1|R_2 \cdots \xrightarrow{\delta_{h+1}} P'_1|Q'_1$$

we derive the transition  $L_{i-1}|Q_1 \xrightarrow{\tau} L_i|Q'_1$  with rule *Com2L*, whether for the other transactions of the premises we use the rule *Par1L*.

$\gamma_{k+1} = \bar{xy}$  : cannot happen because  $\sigma$  is not empty.

*S4R* We have three cases:  $|\sigma_1| = |\sigma_2|$ ,  $|\sigma_1| > |\sigma_2|$  or  $|\sigma_2| > |\sigma_1|$ . In the first case  $|\sigma_3|$  must be  $\tau$  and we can build a chain of transition as in the previous cases. In the second case there is a prefix of  $\sigma_1$  which synchronize with  $\sigma_2$  and  $\sigma_3$  is the rest of  $\sigma_1$ , in this case we can also build a chain of transition as in the previous cases. The third case is symmetric to the second.

□

The converse of lemma 5.2.1 does not hold because the low semantic allow to express interleaving behaviour. But there is the following weaker result:

**Proposition 5.2.2.** Let  $\rightarrow$  be the relation defined in table 5.1, let  $\alpha$  be an action and  $P, Q$  be processes. If  $P \xrightarrow{\alpha} Q$  then  $P \xrightarrow{\alpha} Q$ .

*Proof.* The proof is an easy induction on the proof tree of  $P \xrightarrow{\alpha} Q$ .

□



# Bibliography

- [1] Roberto Gorrieri, Cristian Versari, *Multi  $\pi$ : a calculus for mobile multi-party and transactional communication*.
- [2] Robin Milner, Joachim Parrow, David Walker, *A calculus of mobile processes, part II*, 1990.
- [3] Roberto Gorrieri, *A fully-abstract semantics for atomicity*, Dipartimento di scienze dell'informazione, Università di Bologna.
- [4] Joachim Parrow, *An introduction to the  $\pi$  calculus*, Department Teleinformatics, Rotal Institute of Technology, Stockholm.
- [5] Davide Sangiorgi, David Walker, *The  $\pi$ -calculus*, Cambridge University Press.
- [6] Davide Sangiorgi, *A theory of bisimulation for the  $\pi$ -calculus*, Acta informatica, 33(1):69-97, 1996.
- [7] Milner, Robin, *Communicating and mobile systems: the  $\pi$ -calculus*, Cambridge University Press.
- [8] MohammedReza Mousavi, Michel A Reniers, *Congruence for structural congruences*, Department of Computer Science, Eindhoven University of Technology.