

Copernicus Atmosphere Monitoring Service
Technical User Guide

Final Report

Nikolaos Kalogeras
National Observatory of Athens
space15009@uop.gr

Ioannis Karavasilis
National Observatory of Athens
space15011@uop.gr

Vyron Vasileiadis
National Observatory of Athens
space15003@uop.gr

Abstract

All Human, and not only, activities, are affected by the chemical composition of the air close to the surface of the Earth, generally referred as Air Quality (AQ). Low quality Air Exposure, affects a large percentage of human and animal population in the form of respiratory diseases and allergic responses. European commission is very sensitive with issues concerning public health and environment, and supports numerous efforts that describe and predict parameters and procedures that affect air quality. One of the most recent programme is the Copernicus Atmosphere Monitoring Service (CAMS).

The purpose of this guide is to provide individuals, not necessarily scientists, who are looking for the kind of data, that will allow them to create, defend or even debunk Legislation & Policies concerning factors that affect AQ, with the help of CAMS. In order to evaluate CAMS functionality three different factors affecting AQ are examined: dust aerosols, pollen, ozone and mononitrogen oxides.

Contents

1. Introduction	1
2. Study Area, Data & Materials	1
2.1. Pollutants	1
2.1.1. Aerosols: The case of Dust	1
2.1.2. NO _x : Nitric Oxide (NO) + Nitrogen Dioxide (NO ₂)	1
2.1.3. Ozone (O ₃)	2
2.1.4. Pollen	2
2.2. Data generation	3
2.3. Product Services	4
2.3.1. Global	7
2.3.2. European-scale	8
2.4. Data access	6
3. Models	7
3.1. Global	7
3.1.1. C-IFS	7
3.1.2. IFS-LMD	8
3.2. European-scale	8
3.2.1. CHIMERE	10
3.2.2. EMEP	10
3.2.3. EURAD-IM	11
3.2.4. LOTOS-EUROS	11
3.2.5. MATCH	12
3.2.6. MOCAGE	13
3.2.7. SILAM	14
3.2.8. ENSEMBLE	14
3.2.9. Availability statistics	15
4. Results & Discussions	15
5. Conclusions	21
References	25

Figures

1. Available options and details of CAMS Catalogue product <i>Global analyses of chemical species - nitrogen dioxide</i>	17
2. Global analysis of NO ₂ concentration for December 4th, 2016	18
3. User Interface of ECMWF website for downloading data of global analyses and forecasts.	19
4. Additional filtering of global analyses' and forecasts' data before downloading.	19
5. Available options in <i>Ensemble Analysis and Forecast</i> tab	20
6. Outline of analyses and forecast maps of regional products.	21
7. Analysis of NO ₂ concentration on 00:00 UTC December 4th, 2016 issued from the ENSEMBLE model.	22
8. Maximum concentrations of NO ₂ on December 4th, 2016 as calculated from all models and their ensemble.	22
9. 4-day forecast of O ₃ , NO ₂ , SO ₂ and PM ₁₀ at the city of Athens for the period between December 4th - December 8th, 2016.	23
10. Forecast and analysis maps of NO ₂ concentration for De- cember 4th, 2016 issued by the CHIMERE model.	24
(a) Forecast issued on December 3rd, 2016	??
(b) Analysis issued on December 5th, 2016	??

Tables

1. General characteristics of the regional models.	9
2. Characteristics of the daily assimilation chains of the regional models	9
3. Products of CHIMERE model	10
4. Products of EMEP model	11
5. Products of EURAD-IM model	12
6. Products of LOTOS-EUROS model	12
7. Products of MATCH model	13
8. Products of MOCAGE model	14
9. Products of SILAM model	14
10. Products of SILAM model	15
11. Availability statistics of models' outputs for usage in ENSEMBLE	16

1. Introduction

The chemical composition of the air close to Earth's surface, generally referred as *air quality* (AQ), directly affects human and animal health and also the vegetation. Recently, the World Health Organization reported that in 2012 around 3.7 million deaths were attributable to ambient air pollution ¹

Since the Helsinki Protocol in 1985, many regions and countries, including the European Union countries, have progressively put in place tools to regulate and to control the emissions of the main air pollutants. This has led to an important effort to monitor the air composition near the surface but also to develop air quality forecasting systems in experimental or operational modes [@ebel2007advanced; @menut2010atmospheric]. These tools can be used in cases of high pollution episodes to inform people and to take emergency measures to prevent harming effects. They can also be used for policy makers for the regulations on air pollutant emissions and for monitoring the effect of these regulations on air quality.

The Copernicus Atmosphere Monitoring Service (CAMS) has been developed to meet these needs, aiming at supporting scientists, policymakers, business and citizens with enhanced atmospheric environmental information.

In the following sections, we will study the functionality of CAMS from a user's perspective, the global and regional models it uses, and the data produced for three fundamental air pollutants: ozone, nitric oxide and nitrogen dioxide.

2. Study Area, Data & Materials

2.1. Pollutants

2.1.1. Aerosols: The case of Dust

Aerosols are tiny solid and liquid particles, that are suspended in the atmosphere. Examples of aerosols are windblown dust, sea salts, volcanic ash, smoke from wildfires, and pollution from factories. Depending upon their size, type, and location, aerosols can either cool the surface, or warm it. They can help clouds to form, or they can inhibit cloud formation. And if inhaled, some aerosols can be harmful to people's health.

Aerosols are either created naturally (nearly 90% of the calculated mass), or generated from human activities (the remaining 10%). Dust is one of the most abundant aerosols, since sandstorms whip small pieces of mineral dust from deserts into the atmosphere. Dust tends to consists of larger particles, than their human-made counterparts.

2.1.2. NO_x: Nitric Oxide (NO) + Nitrogen Dioxide (NO₂)

NO_x is a generic term for the mono-nitrogen oxides [@mollenhauer2010handbook; @Omidvarborna2015113], nitric oxide (NO) and nitrogen dioxide (NO₂). They

¹http://www.who.int/phe/health_topics/outdoorair/databases/en/

are produced from the reaction among nitrogen, oxygen and even hydrocarbons (during combustion), especially at high temperatures [@Omidvarborna2015113; @annamalai2007combustion].

Because NO_x are transparent to most wavelengths of light, they allow the vast majority of photons to pass through and, therefore, have a lifetime of at least several days. Because NO₂ is recycled from NO by the photo reaction of VOC (volatile organic compounds) to make more ozone, NO₂ seems to have an even longer lifetime and is capable of traveling considerable distances before creating ozone. Weather systems usually travel over the earth's surface and allow the atmospheric effects to move downwind for several hundred miles. This was noted in EPA reports more than twenty years ago. [@epabulletin] These reports found that each major city on the East coast has a plume of ozone that extends more than a hundred miles out to sea before concentrations drop to 100 parts per billion (ppb).

NO_x mainly impacts on respiratory conditions causing inflammation of the airways at high levels. Long term exposure can decrease lung function, increase the risk of respiratory conditions and increases the response to allergens. NO_x also contributes to the formation of fine particles².

2.1.3. Ozone (O₃)

Ozone (O₃) in the troposphere (the lowermost part of the atmosphere, from the surface to 6-15 km height depending on the latitude) is highly relevant for the Earth's climate, ecosystems, and human health. Tropospheric ozone is the third largest contributor to greenhouse radiative forcing after carbon dioxide and methane [@forster2007changes]. It is part of the Earth's shield against ultraviolet radiation, particularly when there is stratospheric ozone depletion [@sabziparvar1998changes].

Ozone plays a crucial role in tropospheric chemistry as the main precursor for the OH radical which determines the oxidation capacity of the troposphere [@seinfeld2006atmospheric]. It is a toxic air pollutant affecting human health [@bell2006exposure] and agriculture [@fowler2008ground]. Furthermore, through plant damage, it impedes the uptake of carbon into the biosphere [@sitch2007indirect].

Accurate long-term measurements of ozone in the troposphere, including near the earth surface in unpolluted and polluted environments, are needed in order to assess the impacts of tropospheric ozone on the earth system, human health and ecosystems, and to detect changes in the atmospheric composition which could aggravate or reduce these impacts because of changing ozone precursor emissions or climate change.

2.1.4. Pollen

Pollen is a very fine powder produced by trees, flowers, grasses, and weeds in order to fertilize other plants of the same species. Then it is transported through

²<http://www.icopal-noxite.co.uk/nox-problem/nox-pollution.aspx#sthash.esEBqdj6.dpuf>

air, especially through Spring months.

These loose particles are one of the most common causes of allergies throughout the World, such as the Oral Allergy Syndrome (OAS)[^{pollen1}]. At the United States, only, pollen allergies affect up to 30 percent of adults and 40 percent of children. Other reports increase those numbers up to 75 percent of adult population, whereas asthma triggered by pollen affects around 300 million people worldwide[^{pollen2}].

Pollen is considered a chemically non-reactive aerosol, whose atmospheric dispersion only depends on the basic properties of the grains, such as their aerodynamic diameter and density.

2.2. Data generation

CAMS is based on the processing of environmental data collected from Earth observation satellites and in situ sensors. Space segment is divided into two groups of missions: 1. The Sentinels, which are currently being developed especially for Copernicus programme. Sentinel-1,-2, -3, -5P and -6 are dedicated satellites, while Sentinel-4 and 5 are instruments onboard EUMETSAT's weather satellites. European Space Agency (ESA) is responsible for Sentinel-1,-2 and 5P in the future, while EUMETSAT is responsible for operating the Sentinel-3,-6 and the instruments at -4 and -5. 2. The Contributing Missions, which are operated by national, European or international organizations and already provide a wealth of data for Copernicus services. ESA and EUMETSAT will coordinate the delivery of data of these missions.

Sentinel 4, 5 and 5P are dedicated satellites for Atmosphere Monitoring Services.

The Sentinel-4 mission aims to support operational services covering air-quality near-real time applications, air-quality protocol monitoring and climate protocol monitoring. Also, wants to achieve this with a high revisit time over Europe.¹ It uses Ultraviolet-Visible-Near-Infrared (UVN) instrument, a high-resolution spectrometer covering the ultraviolet, visible and near-infrared bands. The spectral resolution is 0.5 nm in the ultraviolet and visible bands, with the goal of 0.12 nm in near-infrared². Moreover, it will use data from Eumetsat's thermal InfraRed Sounder (IRS), embarked on the MTG-Sounder (MTG-S) satellite. After the MTG-S satellite is in orbit, the Sentinel-4 mission also includes data from Eumetsat's Flexible Combined Imager (FCI) embarked on the MTG-Imager (MTG-I) satellite. Sentinel-4 will be a payload embarked on EUMETSAT's Meteosat Third Generation (MTG), which is scheduled to be launched around 2019.

Sentinel-5 will be part of EUMETSAT's Metop Second Generation (Metop-SG), which is to be launched in 2020. It uses an Ultraviolet Visible Near-infrared Shortwave (UVNS) spectrometer and data from Eumetsat's IRS, the Visible Infrared Imager (VII) and the Multi-Viewing Multi-Channel Multi-Polarization Imager (3MI)¹. The Sentinel-5/UVNS instrument is a high-resolution spectrometer operating in the ultraviolet to shortwave infrared range with 7 different spectral bands. Its spatial resolution is below 8km for wavelengths above 300nm

and below 50km for wavelength below 300nm². The orbit is high inclination (approximately 98.7°) LEO, with orbital cycle 29 days (14 orbits per day, 412 orbits per cycle)³. SENTINEL-5 mission will continue spectrometer measurements of previous missions, such as the Global Ozone Monitoring Experiment (GOME) (1995-2011), GOME-2 (2006 till now), with a second GOME-2 on MetOp-B satellite (2012 till now), SCanning Imaging Absorption spectroMeter for Atmospheric CartographHY (SCIAMACHY) on ESA's ENVISAT mission (2002 – 2012) and Ozone Monitoring Instrument (OMI) on NASA's AURA spacecraft (2004 till now)¹, but with a far better spatial resolution. Products from Sentinel-5 are associated with three levels of processing, with Level-0 not distributed to end-users, Level 1B delivered less than 135 minutes from sensing and Level-2 delivered within 165 minutes.

Sentinel-5 Precursor mission will fill the five-year gap until the launch of Sentinel-5. It will use the TROPOMI UV-VIS-NIR-SWIR instrument which is an advanced spectrometer. It will use 8 spectral bands. It follows a high inclination (98.74°), Sun-synchronous, Low-Earth (824 km) polar orbit., with a repeat cycle of 17 days. For Near Real Time processing the following time constraints must be met: For Level-1B should be delivered by 2 h 15 min after sensing and Level-2 up to 3 h after sensing.

It is obvious that in order for CAMS to take full advantage of Sentinels' potential it has to wait a few years. Until then it uses data from the contributing missions, a large number of stored data from previous observations and data gathered daily from ground stations.

2.3. Product Services

CAMS provides products on two scales: global and European-scale.

2.3.1. Global

The global products are provided by the ECMWF Integrated Forecast System (IFS). Products concerning chemical species (O_3 , NO, NO_2) use the Composition-Integrated Forecast System (C-IFS). Products concerning aerosols use a simple bin scheme (IFS-LMD), but there are plans for a more elaborate modal scheme for the future (IFS-GLOMAP). Currently there are no global products for pollen.

The services consist of daily near real-time analyses and forecasts up to five days, and a re-analysis, currently for the period 2003 - 2012. The geographical area covered is the whole Earth, from 180°W to 180° and 90°S to 90°N. Vertically global services cover both the troposphere and stratosphere.

Global analyses of aerosol concentrations - dust

This service provides daily analyses of dust concentrations. The Service became operational on July 1st 2015 and provides data every 6 hours.

Global analyses of chemical species

This service provides daily analyses of chemical species, including O₃, NO and NO₂. The service became operational on July 1st 2015 and provides data every 6 hours.

Global forecasts of aerosol concentrations - dust

This service provides daily forecasts up to 5 days of dust concentrations. The service became operational on July 1st 2015.

Global forecasts of chemical species

This service provides daily forecasts up to 5 days of chemical species, including O₃, NO and NO₂. The service became operational on July 1st 2015.

MACC global reanalysis of aerosol concentrations - dust

This service provides global reanalysis of dust concentrations. The service time coverage spans from January 2003 to 31 December 2012, and has a time resolution of 3 hours.

MACC global reanalysis of assimilated chemical species

This service provides global reanalysis of chemical species that are directly constrained by observations. Only O₃ and NO₂ reanalysis data are available. This service time coverage spans from January 2003 to 31 December 2012, and has a time resolution of 3 hours. Horizontal resolution is approximately 60 km.

MACC global reanalysis of non-assimilated chemical species

This service provides global reanalysis of chemical species that are not directly constrained by observations. Only NO and NO₂ reanalysis data are available. This service time coverage spans from January 2003 to 31 December 2012, and has a time resolution of 3 hours. Horizontal resolution is 1.125° x 1.125°.

2.3.2. European-scale

The European-scale regional products are provided for the European domain (25°W to 45°E and from 30°N to 70°N). The service provides daily 4-day forecasts of the main air quality species, including O₃, NO, NO₂, and birch pollen and analyses of the day before, from 7 state-of-the-art atmospheric chemistry models and from the median ensemble calculated from the 7 model forecasts. The 7 atmospheric chemistry models are: CHIMERE, EMEP, EURAD-IM, LOTOS-EUROS, MATCH, MOCAGE, SILAM. The regional service also provides daily posteriori interim re-analyses based on fast-track in-situ observations, and annual re-analyses based on fully validated in-situ observations. There are no regional products concerning dust concentrations.

European-scale air quality analysis

This service provides European-scale air quality analyses for every hour of the previous day for every of the 7 models and the ensemble median. The service became operational on October 1st 2015.

European-scale air quality forecast

This service provides European-scale air quality forecasts for every hour up to 4 days in advance for every of the 7 models and the ensemble median. The service became operational on October 1st 2015.

2.4. Data access

Products delivered by CAMS are provided free of charge, since October 1st 2015. There is a universal search engine available on the CAMS web portal. The default procedure that should be used by common user is the following:

1. Go to the CAMS website <http://atmosphere.copernicus.eu>
2. Click CATALOGUE. The CAMS catalogue lists the various CAMS products.
3. Filter the catalogue for the service or product of interest. The filters change depending on the selection.
4. Once a suitable product is identified, click on its name. A popup window opens, showing details and a preview.
5. At the bottom of the window follow the links to Plots (maps and graphs), Data Access, Documentation, etc. These links take you to pages specific to the selected product.
6. From here the procedure is different for every product, depending on the provider.

In particular:

Global products are served by European Centre for Medium-Range Weather Forecasts (ECMWF). A registration is required to access the data. CAMS provides its global data sets in two different ways.

One option is the interactive web interface of ECMWF³ data server which provides batch data access to real-time analyses and forecasts, reanalysis results in GRIB and NetCDF format. It also provides a programmatic way of accessing data using python language. Detailed information can be found in ECMWF wiki⁴.

The second option is through the ECMWF FTP server⁵, which provides archived real-time forecasts of the latest three days in GRIB and NetCDF format. CAMS data files can be accessed through directory `/DATA/CAMS_NREALTIME/${YYYYMMDDhh}`

³<http://apps.ecmwf.int/datasets/data/cams-nrealtime/levtype=sfc/>

⁴<https://software.ecmwf.int/wiki/display/WEBAPI/Access+ECMWF+Public+Datasets>

⁵<ftp://dissemination.ecmwf.int/>

and follow a naming convention proposed by World Meteorological Organization (WMO). Detailed information can be found in CAMS website⁶.

European-scale products are accessed directly through CAMS web portal⁷. Files are available in Grib2 format and Netcdf format for model ENSEMBLE for both Forecast (surface as well as other levels) & Analysis (surface). Files are available only in Netcdf format for models: CHIMERE, EMEP, EURAD, LOTOSEUROS, MATCH, MOCAGE, SILAM for both Forecast (surface as well as other levels) & Analysis (surface). There are six available choices:

- **Download online data** Files produced less than 30 days ago are available. Available formats are GRIB2, NetCDF or TXT, depending on the type of product.
- **Download archived data** Archived data are older than 30 days. Archiving starts in October 2015. Available formats are GRIB2, NetCDF or TXT, depending on the type of product.
- **Download reanalysis data** Download numerical data about atmospheric composition in Europe (reanalysis). Archiving starts in 2015 and the available data format is NetCDF.
- **Download on map viewing** Visualize data over a map of Europe. Very useful for an interesting air quality phenomenon spotted on a map view.
- **Data server - subscribe mode** Data server subscribe mode allows users to subscribe to a product, to choose the delivery type (ftp,...), as well as the delivery time (upon arrival, on a specific chosen time).
- **Data server - catalogue** Consult the catalogue of metadata, in order to discover all the CAMS Regional Air Quality products.

3. Models

3.1. Global

The operational CAMS global assimilation and forecasting system uses fully integrated chemistry in the ECMWF Integrated Forecasting System (IFS).

3.1.1. C-IFS

For chemical species, the version of the IFS is named Composition-IFS (C-IFS). C-IFS makes it possible

- to use the detailed meteorological simulation of the IFS for the simulation of the fate of constituents,
- to simulate atmospheric chemistry globally at high resolution because of the computational parallel efficiency of the IFS,

⁶<http://atmosphere.copernicus.eu/ftp-access-global-data>

⁷http://www.regional.atmosphere.copernicus.eu/index.php?category=data_access&subensemble=dataserver_services

- to use the IFS data assimilation system to assimilate observation of atmospheric composition and
- to simulate feedback between atmospheric composition and weather. Different chemical schemes can be used with the current operational version being based on the CB05 scheme.

C-IFS replaces the coupled system IFS-MOZART for forecast and assimilation of reactive gases within the pre-operational Copernicus Atmosphere Monitoring Service.

3.1.2. IFS-LMD

The IFS-LMD aerosol scheme is the scheme that was introduced to add aerosol modelling to the ECMWF IFS forecasting system. It is currently used for the daily analysis and 5-day forecast and was also used for the MACC reanalysis.

The initial package of ECMWF physical parameterizations dedicated to aerosol processes mainly follows the aerosol treatment in the LOA/LMD-Z model (Boucher et al. 2002; Reddy et al. 2005). Five types of tropospheric aerosols are considered: sea salt, dust, organic and black carbon, and sulphate aerosols. Prognostic aerosols of natural origin, such as mineral dust and sea salt are described using three size bins. For dust, radius bin limits are at 0.03, 0.55, 0.9, and 20 microns while for sea-salt radius bin limits are at 0.03, 0.5, 5 and 20 microns.

A detailed description of the ECMWF forecast and analysis model including aerosol processes is given in Morcrette et al. (2009) and Benedetti et al. (2009).

3.2. European-scale

Each of the seven models is run at its own horizontal and vertical resolutions, with the horizontal resolutions varying between 10 and 20 km. This range of resolutions is not designed to reproduce local aspects of air pollution but to provide concentrations of pollutants at the regional scale that can then be used in particular as boundary conditions for air quality forecasts at finer resolution.

The range of the forecasts is 96h from 00:00 UTC on Day0 with hourly outputs on eight vertical levels (surface, 50, 250, 500, 1000, 2000, 3000 and 5000m). Day0 is defined as the day when the forecast is run. The forecast initial time/date is Day0 at 00:00 UTC and final time/date is Day3 at 24:00UTC. For each time step (1h), the individual model fields are interpolated on these vertical levels and on the same regular 0.1° latitude by 0.1° longitude grid over the European domain. It is from these re-gridded fields that the ensemble median and verification products are calculated.

The analysis at the surface for Day0–1 (the day before Day0) is run daily a posteriori on Day0 using the assimilation of the hourly data from the AQ monitoring stations available in Europe between 00:00 and 23:00UTC on Day0–1. Like for the forecasts, Day0 is defined as the day when the analysis is run. The analysis initial time/date is Day0–1 at 00:00 UTC and final time/date is

Day0–1 at 23:00 UTC. Similarly to the forecasts, the hourly individual model fields are interpolated on the same 0.1° latitude by 0.1° longitude grid. The analyses are only produced at the surface level.

An overview of the characteristics of the seven models can be seen in tables 1 and 2. Detailed descriptions of each model are presented in the following sections.

Model	Operated By	Horizontal Resolution	Vertical Resolution
CHIMERE	INERIS, France	$0.1^\circ \times 0.1^\circ$	8 levels
			Top at 500 hPa
EMEP	MET Norway, Norway	$0.25^\circ \times 1.25^\circ$	20 levels
			Top at 100 hPa
EURAD-IM	RIU UK, Germany	15km on a Lambert conformal projection	23 levels
			Top at 100 hPa
LOTOS-EURO	KNMI, Netherlands	$0.25^\circ \times 1.25^\circ$	4 levels
			Top at 3.5 km
MATCH	SMHI, Sweden	$0.2^\circ \times 0.2^\circ$	52 levels
			Top at 300 hPa
MOCAGE	Météo-France, France	$0.2^\circ \times 0.2^\circ$	47 levels
			Top at 5 hPa
SILAM	FMI, Finland	$0.15^\circ \times 0.15^\circ$	8 levels
			Top at 6.7 km

Table 1. General characteristics of the regional models.

Model	Assimilation Method	Observation assimilated	Pollutants analysed
CHIMERE	Optimal interpolation	O ₃ from surface stations	O ₃ , NO, NO ₂
EMEP	3DVar	NO ₂ columns from OMI	O ₃ , NO, NO ₂
		NO ₂ from surface stations	O ₃ , NO, NO ₂
EURAD-IM	3DVar	O ₃ , NO, NO ₂ from surface stations	O ₃ , NO, NO ₂
		NO ₂ columns from OMI and GOME-2	O ₃ , NO, NO ₂
LOTOS-EURO	Ensemble Kalman filter	O ₃ from surface stations	O ₃ , NO, NO ₂
MATCH	3DVar	O ₃ , NO ₂ from surface stations	O ₃ , NO, NO ₂
MOCAGE	3DVar	O ₃ from surface stations	O ₃ , NO, NO ₂
SILAM	4DVar	O ₃ , NO ₂ from surface stations	O ₃ , NO, NO ₂

Table 2. Characteristics of the daily assimilation chains of the regional models

3.2.1. CHIMERE

CHIMERE is an Eulerian chemistry-transport model able to simulate concentration fields of gaseous and aerosols species at a regional scale [@menut2013chimere]. The model is developed under the General Public License licence⁸. CHIMERE is used for analysis of pollution events, process studies [@beekmann2010modelling], experimental and operational forecasts [@rouil2009prev], regional climate studies and trends [@colette2011air], among others.

CHIMERE runs over a range of spatial scale from the regional scale (several thousand kilometres) to the urban scale (100-200 Km) with resolutions from 1-2 Km to 100 Km. The model runs over the GEMS-MACC domain with a 0.1° resolution and 8 vertical levels extending from the surface up to 500 hPa, covering the whole troposphere.

CHIMERE reproduces nicely the day to day O₃ variation similarly at urban and rural sites with an overestimation which is higher during the winter at urban sites. CHIMERE reproduces the daily NO_x variability along the year but underestimates significantly the concentration especially during the cold season.

Availability information of CHIMERE products as of May 2016 are presented in Table 3.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	6:00 UTC	09:45 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 3. Products of CHIMERE model

3.2.2. EMEP

The EMEP/MSC-W model has been developed at the EMEP Meteorological Synthesizing Centre-West at the Norwegian Meteorological Institute. The model has been publicly available as open-source code since 2008, and a detailed description is given in Simpson et al. (2012).

One strength of the EMEP model is that its domain extends throughout the whole troposphere, thus taking accurate account of long-range transport of pollutants in the free troposphere. As the model is designed mainly for background concentrations, urban increments have not been implemented as in some other models with equally coarse resolution, leading to somewhat lower performance in urban and sub-urban areas. However, being one of the main

⁸Official website: <http://www.lmd.polytechnique.fr/chimere/>

research tools under the UN LRTAP (Long-range Transboundary Air Pollution) convention, the EMEP model is evaluated continuously against measurements of a large range of chemical parameters (including air concentrations, depositions, and trends) ensuring modelling capability with very good overall performance [@jonson2006first; @fagerli2008trends; @genberg2013light].

Availability information of EMEP products as of May 2016 are presented in Table 4.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	2:00 UTC	08:45 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 4. Products of EMEP model

3.2.3. EURAD-IM

EURAD-IM is an Eulerian meso-scale chemistry transport model involving advection, diffusion, chemical transformation, wet and dry deposition and sedimentation of tropospheric trace gases and aerosols (Hass et al., 1995, Memmesheimer et al., 2004). It includes 3DVar and 4DVar chemical data assimilation (Elbern et al., 2007) and is able to run in nesting mode. EURAD-IM has been applied on several recent air pollution studies [@monteiro2013bias; @paschali2015inverse].

Availability information of EURAD-IM products as of May 2016 are presented in Table 5.

3.2.4. LOTOS-EUROS

The 3-D chemistry-transport model LOTOS-EUROS [@schaap2008lotos] is developed by the Dutch institutes TNO, RIVM and, more recently, KNMI. It is used for regional-scale air quality forecasts in Europe and the Netherlands [@de2011six].

The model extends up to 3.5km above sea level, with three dynamic layers and a fixed 25m thick surface layer. The lowest dynamic layer is the mixing layer, followed by two reservoir layers. The height of the mixing layer is obtained from the ECMWF meteorological input data used to drive the model.

The LOTOS-EUROS model has been designed as a model of intermediate complexity, to favour short computation times. For this, the vertical top of the operational model version is limited and covers only the boundary layer

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	6:00 UTC	09:45 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Until April 20th, 2016 NO observations from the German Umweltbundesamt (UBA) were assimilated. Thereafter surface in situ observations from UBA have been removed from the EURAD-IM assimilation chain, to be compliant with CAMS requirements.

Table 5. Products of EURAD-IM model

and reservoir layers (up to 3.5km); effectively, the model therefore employs only four dynamic layers. Concentrations from the free troposphere are taken from the global boundary conditions, and therefore fully incorporate the knowledge, assimilations, and validation efforts present in the global model.

In spite of the limited complexity, the model performs well in simulation of O₃ [@curier2012improving].

Apart from the relative short run-through time, the strength of the model is in the detailed description of anthropogenic emissions, given the close cooperation with the developers of the TNO-MACC emission inventory; this is for example shown by excellent simulation of boundary layer NO₂ [@vlemmix2015max].

Availability information of LOTOS-EUROS products as of May 2016 are presented in Table 6.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	4:00 UTC	10:00 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 6. Products of LOTOS-EUROS model

3.2.5. MATCH

The MATCH model has been developed at SMHI over the past 20 years and is applied for emergency purposes as well as for regional-scale chemistry modelling [@langner1998validation].

The strength of the MATCH model is that it spans vertically the troposphere and makes use of the same vertical layers as provided from the IFS model up to 300hPa. This means about 50 layers in the vertical and the lowest one just 20m thick and about 15 in the boundary layer. Using the same vertical resolution as the IFS model is an advantage because no vertical interpolation is required. Nevertheless, since the MATCH model has been developed mainly using HIRLAM (HIgh-Resolution Limited Area Model) data with a coarser vertical resolution, the use of the high-resolution vertical levels from IFS may lead to less accurate chemistry forecasts compared to the HIRLAM version

Availability information of MATCH products as of May 2016 are presented in Table 7.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	3:00 UTC	10:00 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 7. Products of MATCH model

3.2.6. MOCAGE

The MOCAGE model [@josse2004radon] has been developed at Météo-France since 2000. Its assimilation system has been developed jointly with CERFACS. This model and its assimilation system have been successfully used for tropospheric and stratospheric research [@bousserez2007evaluation; @lacressoniere2014european] and also for operational purposes [@rouil2009prev].

The strength of MOCAGE is that it simulates the air composition of the whole troposphere and lower stratosphere. Thus, it provides a full representation of transport processes, in particular boundary layer–troposphere and troposphere–stratosphere exchanges, and the time evolution of stratospheric conditions for accurate photolysis rate calculations at the surface.

The MOCAGE assimilation system in its MACC configuration produces robust analyses for both O₃ and NO₂ as illustrated in the annual re-analysis reports ⁹

Availability information of MOCAGE products as of May 2016 are presented in Table 8.

⁹<http://www.gmes-atmosphere.eu/documents/maccii/deliverables/eva/>

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	0:30 UTC	10:30 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 8. Products of MOCAGE model

3.2.7. SILAM

SILAM is a meso-to-global-scale dispersion model [sofiev2008construction] that is used for atmospheric composition, emergencies, composition–climate interactions, and air quality modeling purposes. The model has been applied with resolutions ranging from 1km up to 3°, incorporates eight chemical and physical transformation modules and covers the troposphere and the stratosphere. SILAM is publicly available since 2007 and is used as an operational and research tool.

Availability information of SILAM products as of May 2016 are presented in Table 9.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	3:00 UTC	9:30 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO*, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

* Not assimilated

Table 9. Products of SILAM model

3.2.8. ENSEMBLE

To process the ensemble median, all seven individual models are first interpolated to a common 0.1°×0.1° horizontal grid. For each grid point, the ensemble model value is calculated as the median value of the individual model forecasts or analyses available. The median is defined as the value having 50% of individual models with higher values and 50% with lower values. This method is rather insensitive to outliers in the forecasts or analyses and is very efficient computationally. It is also little sensitive if a particular model forecast or analysis is

occasionally missing. These properties are very useful from an operational point of view.

For the forecasts, the ENSEMBLE is produced for all levels and all pollutants. For the analyses, the individual assimilation systems provide only analyses at the surface level.

Availability information of ENSEMBLE products as of May 2016 are presented in Table 10.

	Forecast	Analysis
Altitudes	Surface, 50m, 250m, 500m, 1000m, 2000m, 3000, 5000m	Surface
Available at	6:45 UTC for D0-D1 8:45 UTC for D2-D3	11:30 UTC for the day before
Pollutants	O ₃ , NO, NO ₂	O ₃ , NO, NO ₂
Timespan	0-96h, hourly	0-24h for the day before, hourly

Table 10. Products of SILAM model

3.2.9. Availability statistics

The statistics in Table 11 describe the ratio of days for which the models' outputs were available on time to be included in the ENSEMBLE fields (analyses and forecasts) that are computed at METEO-FRANCE. They are based on the following schedule:

- forecasts data before: 05:30 UTC for Day0-Day1 (up to 48h), 07:30 UTC for Day2-Day3 (from 49h to 96h)
- analyses data before: 11:00 UTC

Statistics are based on the quarter of March - May 2016, which is the latest with published data¹⁰.

4. Results & Discussions

In this section, we will reproduce a real-life case of air quality monitoring, to test the validity of CAMS forecast services and evaluate its easiness of use. Specifically, we will study the concentration of nitrogen dioxide on December 4th, 2016 in global and European scale.

First, we access the online catalogue available on CAMS website. We set the provided filters as following:

- Product family = { Global analyses, Global forecasts, Regional analyses, Regional forecasts }

¹⁰<https://atmosphere.copernicus.eu/documentation-regional-systems>

Model	Forecast Availability	Analysis Availability
CHIMERE	Day0: 85%	83%
	Day1: 85%	
	Day2: 85%	
	Day3: 75%	
EMEP	Day0: 97%	97%
	Day1: 96%	
	Day2: 96%	
	Day3: 96%	
EURAD-IM	Day0: 99%	98%
	Day1: 99%	
	Day2: 99%	
	Day3: 99%	
LOTOS-EURO	Day0: 97%	97%
	Day1: 97%	
	Day2: 97%	
	Day3: 97%	
MATCH	Day0: 99%	97%
	Day1: 99%	
	Day2: 99%	
	Day3: 97%	
MOCAGE	Day0: 100%	98%
	Day1: 100%	
	Day2: 99%	
	Day3: 99%	
SILAM	Day0: 97%	96%
	Day1: 97%	
	Day2: 97%	
	Day3: 97%	

Table 11. Availability statistics of models' outputs for usage in ENSEMBLE

- Parameter = { Birch pollen, Dust concentration, Nitrogen dioxide, Nitrogen monoxide, Ozone }

64 results are returned. Analytically:

- 8 regional forecasts for birch pollen (7 individual models plus ensemble)
- 8 regional analyses for birch pollen (7 individual models plus ensemble)
- 1 global analysis for dust concentration

- 1 global forecast for dust concentration
- 1 global analysis for nitrogen dioxide
- 1 global forecast for nitrogen dioxide
- 8 regional forecasts for nitrogen dioxide (7 individual models plus ensemble)
- 8 regional analyses for nitrogen dioxide (7 individual models plus ensemble)
- 1 global analysis for nitrogen monoxide
- 1 global forecast for nitrogen monoxide
- 8 regional forecasts for nitrogen monoxide (7 individual models plus ensemble)
- 8 regional analyses for nitrogen monoxide (7 individual models plus ensemble)
- 1 global analysis for ozone
- 1 global forecast for ozone
- 8 regional forecasts for ozone (7 individual models plus ensemble)
- 8 regional analyses for ozone (7 individual models plus ensemble)

At first glance, we can verify that CAMS provides both global and regional products for chemical species (O_3 , NO, NO_2), only global products for dust concentration and only regional products for birch pollen.

All products when clicked lead to a page with several links available. This can be seen in Figure 1 for the case of global analysis. We are primarily concerned about the provided plots and data. As it is mentioned before, the procedure onward is different for global and regional products.




Figure 1. Available options and details of CAMS Catalogue product *Global analyses of chemical species - nitrogen dioxide*

For global products, *Plots* links to a page within CAMS website where an interactive global map is provided. Figure 2 shows the generated map of global

analysis of NO₂ concentration for 00:00 UTC December 4th, 2016. Information about NO₂ concentration is color-layered on the map with a measurement unit legend provided. The user is able to adjust the date/time and the vertical area of interest.




Figure 2. Global analysis of NO₂ concentration for December 4th, 2016

It should be noted that the plots service seems confusing at first, since both global analyses and forecasts link to the same page. The distinction between analysis and forecast is based on the date/time selected in respect to current date/time. Also, the plot service is time-limited in the sense that provides analyses for the past 5 days and forecasts for the following 5 days only.

Global product *Data download* link to a page within CAMS website where the user can choose to download data through an interactive interface or through ftp access. Interactive data access link to ECMWF website. The user is presented a rich web form to select the exact product of interest. This is shown in Figure 3. A date-span or specific months can be selected, different vertical levels (surface or pressure dependent or model dependent), multiple time-steps and a huge parameter list, which of course includes NO₂. A drawback is that data for the past four days in respect to current date is not available.

After the user has set the filters he is able to download data in both GRID and NetCDF format by clicking the relative link. In both cases he is led to an *Additional filtering* form where he can define the area of interest and the grid resolution as shown in Figure 4. The data is then available for downloading.

For regional products, both *Plots* and *Data download* link to the regional subdomain of CAMS website. The subdomain provides a universal access to individual models and ensemble plots and data for both analyses and forecasts. This is done by an easy-to-use tab section as seen in Figure.

In *Ensemble Analysis and Forecast* tab, except from forecasts and analyses maps, additional plots of daily mean and maximum values as well as EPSgrams for 41 major European cities are available (Figure #ensemble-tab).

Hourly Ensemble Maps (as well as all maps) provide a friendly user interface (UI) with details of the product, a time navigation bar, filters for base date/time, model, vertical level and parameter options. The outline is shown in Figure 6.

CAMS Near-real-time

Select date

Select a date in the interval 2012-07-05 to 2016-11-30

Start date: 2012-07-05 End date: 2016-11-30

Select a list of months

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2012												2013											
2014												2015											
2016																							
Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

Select All or Clear

Select time

00:00:00 06:00:00 12:00:00 18:00:00

Select All or Clear

Select step

<input type="checkbox"/> 0	<input type="checkbox"/> 3	<input type="checkbox"/> 6	<input type="checkbox"/> 9	<input type="checkbox"/> 12	<input type="checkbox"/> 15	<input type="checkbox"/> 18	<input type="checkbox"/> 21	<input type="checkbox"/> 24	<input type="checkbox"/> 27	<input type="checkbox"/> 30	<input type="checkbox"/> 33	<input type="checkbox"/> 36	<input type="checkbox"/> 39
<input type="checkbox"/> 42	<input type="checkbox"/> 45	<input type="checkbox"/> 48	<input type="checkbox"/> 51	<input type="checkbox"/> 54	<input type="checkbox"/> 57	<input type="checkbox"/> 60	<input type="checkbox"/> 63	<input type="checkbox"/> 66	<input type="checkbox"/> 69	<input type="checkbox"/> 72	<input type="checkbox"/> 75	<input type="checkbox"/> 78	<input type="checkbox"/> 81
<input type="checkbox"/> 84	<input type="checkbox"/> 87	<input type="checkbox"/> 90	<input type="checkbox"/> 93	<input type="checkbox"/> 96	<input type="checkbox"/> 99	<input type="checkbox"/> 102	<input type="checkbox"/> 105	<input type="checkbox"/> 108	<input type="checkbox"/> 111	<input type="checkbox"/> 114	<input type="checkbox"/> 117	<input type="checkbox"/> 120	

Select All or Clear

Select parameter

<input type="checkbox"/> 2 metre dewpoint temperature	<input type="checkbox"/> 2 metre temperature
<input type="checkbox"/> 10 metre U wind component	<input type="checkbox"/> 10 metre V wind component
<input type="checkbox"/> UV biologically effective dose clear-sky	<input type="checkbox"/> UV biologically effective dose clear-sky
<input type="checkbox"/> Albedo	<input type="checkbox"/> Black Carbon Aerosol Optical Depth at 550nm
<input type="checkbox"/> Convective precipitation	<input type="checkbox"/> Dust Aerosol Optical Depth at 550nm

Figure 3. User Interface of ECMWF website for downloading data of global analyses and forecasts.

Additional filtering

Current request

Stream: Atmospheric model
Type: Analysis
Dataset: cams_nrealtime
Step: 0
Version: 1
Type of level: Surface
[See full request](#)

The request will be done using the following attributes:

Area: Default (as archived) ([change](#))

- Default (as archived)
- South Asia
- Inter-tropical band
- Northern Hemisphere
- Southern Hemisphere
- Tropical Pacific
- Europe
- North America
- Indonesia
- Custom: N W S E

Grid: Default (as archived) ([change](#))

- Default (as archived)
- 0.125x0.125
- 0.225x0.225
- 0.25x0.25
- 0.4x0.4
- 0.5x0.5
- 0.75x0.75

Figure 4. Additional filtering of global analyses' and forecasts' data before downloading.




Figure 5. Available options in *Ensemble Analysis and Forecast* tab

Hourly Ensemble Maps provide analysis and forecasts in the same UI. User first select a base date/time and then, depending on navigation's bar value, he selects forecast or analysis produced on that day. For example, if selected date/time is 00:00 UTC December 4th, 2016 and time navigation bar value is -10, then the analysis for 14:00 UTC December 3rd is depicted on the map. Similarly, given the same date/time and a value of +48 in time navigation bar, the forecast of 00:00 UTC December 6th is shown. A useful *Download PDF* button, which exports the map to PDF format, is also present.

In order to obtain the analysis of NO₂ concentration on 00:00 UTC December 4th, 2016 we set as date/time the December 5th and a time step of -24. Obtained map is shown in Figure 7.

Daily mean and maximum tab provides maps of the daily mean and maximum values of the concentration of pollutants as calculated by each model and ensemble. A useful feature is the ability to view the maps generated of all models side by side. This feature is shown in Figure 8 for NO₂ maximum concentration values on December 4th, 2016.

EPSgrams tab provide 4-days forecasts of O₃, NO₂, SO₂ and PM₁₀ at the location of 41 major European cities and their uncertainty. In Figure 9 forecasts for the city of Athens between December 4th and December 8th, 2016 are shown. *Individual Analyses* and *Individual Forecasts* tabs provide maps of pollutants' concentration based on individual models calculations. Analyses are available for the day before at an hourly step. Forecasts are available up to 4 days ahead at an hourly step. It is interesting to see the relevance of the forecast for a given date/time to the post-analysis. In Figure 10 we see the forecast of NO₂ concentration for December 4th, 2016 issued on December 3rd and the relevant analysis issued on December 5th. It is obvious that results are quite different. Data files are very easy to download. Under the filters section there is a *Direct Data Access* section with a checkbox for accepting the CAMS data licence. Once the checkbox is checked we are presented with the details of the product (model, pollutant, base date/time) and an optional selectbox with available options




Figure 6. Outline of analyses and forecast maps of regional products.

depending on the product. After an option is selected a direct download in NetCDF format is available by clicking the button.

5. Conclusions

Satellites are capable of measuring the most important air pollutants, greenhouse gases and particles such as those from dust and pollen. They provide timely, spatially explicit, large-scale information on distribution in a cost-effective manner. When these data are coupled with ground-based measurements from monitoring stations and atmospheric chemistry models, accurate information on air quality can be provided in near-real time to support European air quality policies, air pollution mitigation and public health protection.

CAMS provide services both global and regional, with each side having its advantages and drawbacks. For this reason, major updates are incorporated continually, new satellites are launched, and models used for prediction are also modified, in order to produce better assessments.

When studying the usability of CAMS one must have in mind that it is essentially composed of two different services: global and regional.

The global models C-IIFS and IFS-LMD are much more mature and tested. However, the website of ECMWF that provides the results is not very functional




Figure 7. Analysis of NO₂ concentration on 00:00 UTC December 4th, 2016 issued from the ENSEMBLE model.




Figure 8. Maximum concentrations of NO₂ on December 4th, 2016 as calculated from all models and their ensemble.

and can be frustrating for the common user.

On the other hand, the regional models are quite fresh and several were developed for a different purpose, however quarterly reports show that all models





Figure 9. 4-day forecast of O₃, NO₂, SO₂ and PM₁₀ at the city of Athens for the period between December 4th - December 8th, 2016.

are constantly improving over time¹¹. Moreover, the website of regional services is very functional and easy to use, even by a completely inexperienced user.

What anyone can conclude is that we are only at the beginning. CAMS

¹¹<https://atmosphere.copernicus.eu/documentation-regional-systems>



(a) Forecast issued on December 3rd, (b) Analysis issued on December 5th, 2016

Figure 10. Forecast and analysis maps of NO₂ concentration for December 4th, 2016 issued by the CHIMERE model.

is a very powerful and important service, which has enormous possibilities of continuous and large-scale monitoring with advanced satellite-based systems. Also, can provide timely and reliable long-term information for assessment and verification purposes. It is sure that with its predictions can contribute to EU efforts to monitor health-related environmental phenomena.

References