

Beckmann's approach to multi-item multi-bidder auctions

arXiv:2203.06837

Alexander V. Kolesnikov (HSE)

Fedor Sandomirskiy (Caltech, fsandomi@caltech.edu, <https://www.fedors.info/>)

Aleh Tsyvinski (Yale)

Alexander P. Zimin (MIT)

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$		
$n \geq 2:$		

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	Single-item monopolist Myerson (1981): posted price is optimal	
$n \geq 2:$		

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	Single-item monopolist Myerson (1981): posted price is optimal	
$n \geq 2:$	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport^a <hr/> ^a C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	<p>Single-item monopolist</p> <p>Myerson (1981): posted price is optimal</p>	<p>Multi-item monopolist</p> <ul style="list-style-type: none"> optimal mechanisms known in particular cases connection to optimal transport^a <hr/> <p>^aC.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica</p>
$n \geq 2$:	<p>Classic auctions</p> <p>Myerson (1981): 2nd-price auction with reserve is optimal</p>	<p>Multi-item auctions</p> <ul style="list-style-type: none"> almost nothing known about optimal mechanisms^a Our paper: connection to optimal transport <hr/> <p>^aCombined obstacles of multidimensional screening and multidimensional majorization ²</p>

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport^a <hr/> ^a C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	Multi-item auctions <ul style="list-style-type: none">• almost nothing known about optimal mechanisms^a• Our paper: connection to optimal transport <hr/> ^a Combined obstacles of multidimensional screening and multidimensional majorization ²

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM. Beckmann (1952) **A continuous model of transportation** *Econometrica*

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM. Beckmann (1952) **A continuous model of transportation** *Econometrica*

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

- **Econ applications of optimal transport**
 - Monge-Kantorovich: Daskalakis et al. (2017), Kleiner, Manelli (2019), Boerma et al.(2021), Chiapporiet et al. (2010), Galichon (2021), Steinerberger, Tsyvinski (2019), Gensbittel (2015), Arieli et al.,(2022), Guo, Shmaya (2021)
 - Beckmann: Fajgelbaum, Schaal (2020), Allenand, Arkolakis (2014), Santambrogio (2015)
- **Non-transport duality in auction design** Giannakopoulos, Koutsoupas (2018), Cai et al. (2019), Bergemann et al. (2016)
- **Simple mechanisms with good revenue guarantees** Hart, Reny (2019), Haghpasand, Hartline (2021), Babaioff et al. (2020,2021), Hart, Nisan (2017), Jehiel et al. (2007), Yao (2017), and many more...
- **Majorization in economics** Hart and Reny (2015), Kleiner et al. (2021), Arieli et al. (2019), Candogan, Strack (2021), Nikzad (2022)

- Known results: monopolist's problem and its dual
- The case of $n \geq 2$ bidders
 - Similarities and differences
 - Formal statement of duality theorem
- Applications and simulations

Warm-up: $n = m = 1$

- How to sell one good to one agent with the value $v \sim \rho(v) dv$?

Warm-up: $n = m = 1$

- How to sell one good to one agent with the value $v \sim \rho(v) dv$?
- **posted-price mechanism:** chose the best option from the menu
 - pay 0 and get nothing
 - pay t and get the good

Warm-up: $n = m = 1$

- How to sell one good to one agent with the value $v \sim \rho(v) dv$?
- **posted-price mechanism:** chose the best option from the menu
 - pay 0 and get nothing
 - pay t and get the good

best price $t^* = \arg \max_t t \cdot \int_t^\infty \rho(v) dv$

Warm-up: $n = m = 1$

- How to sell one good to one agent with the value $v \sim \rho(v) dv$?
- **posted-price mechanism:** chose the best option from the menu
 - pay 0 and get nothing
 - pay t and get the good

best price $t^* = \arg \max_t t \cdot \int_t^\infty \rho(v) dv$

Theorem (Myerson (1981))

Posted price mechanism with t^* maximizes revenue

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal?
- Is bundling all goods together always optimal?
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal?
- Is bundling all goods together always optimal?
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal?
- Is bundling all goods together always optimal?
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal?
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal?
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal? **No**¹
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism**: chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal? **No**¹
- Is $x \in \{0, 1\}^m$ enough?
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) Maximal revenue with multiple goods: Nonmonotonicity TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal? **No**¹
- Is $x \in \{0, 1\}^m$ enough? **No**¹
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal? **No**¹
- Is $x \in \{0, 1\}^m$ enough? **No**¹
- **menu mechanism:** chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' get x'
 -

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ goods, $n = 1$ agent: menu mechanisms

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- if agent gets a bundle $x = (x_1, \dots, x_m) \in [0, 1]^m$ and pays t , her utility is $\langle x, v \rangle - t$
- Is selling each good separately always optimal? **No**¹
- Is bundling all goods together always optimal? **No**¹
- Is $x \in \{0, 1\}^m$ enough? **No**¹
- **menu mechanism**: chose the best option from the menu
 - pay 0 get 0
 - pay t get x
 - pay t' pay x'
 -

Revelation principle

Any mechanism is equivalent to a menu mechanism

¹Hart, Reny (2015) **Maximal revenue with multiple goods: Nonmonotonicity** TE

$m \geq 2$ items, $n = 1$ agent: optimal menus

- a menu $M \subset \mathbb{R}_+ \times [0, 1]^m$
- utility obtained by an agent with values $v = (v_1, \dots, v_m)$:

$$u_M(v) = \max_{(t, x) \in M} \langle x, v \rangle - t,$$

- u_M is convex and

$$x(v) = \partial u_M(v), \quad t(v) = \langle x(v), v \rangle - u_M(v)$$

$m \geq 2$ items, $n = 1$ agent: optimal menus

- a menu $M \subset \mathbb{R}_+ \times [0, 1]^m$
- utility obtained by an agent with values $v = (v_1, \dots, v_m)$:

$$u_M(v) = \max_{(t, x) \in M} \langle x, v \rangle - t,$$

- u_M is convex and

$$x(v) = \partial u_M(v), \quad t(v) = \langle x(v), v \rangle - u_M(v)$$

$m \geq 2$ items, $n = 1$ agent: optimal menus

- a menu $M \subset \mathbb{R}_+ \times [0, 1]^m$
- utility obtained by an agent with values $v = (v_1, \dots, v_m)$:

$$u_M(v) = \max_{(t,x) \in M} \langle x, v \rangle - t,$$

- u_M is convex and

$$x(v) = \partial u_M(v), \quad t(v) = \langle x(v), v \rangle - u_M(v)$$

$m \geq 2$ items, $n = 1$ agent: optimal menus

- a menu $M \subset \mathbb{R}_+ \times [0, 1]^m$
- utility obtained by an agent with values $v = (v_1, \dots, v_m)$:

$$u_M(v) = \max_{(t,x) \in M} \langle x, v \rangle - t,$$

- u_M is convex and

$$x(v) = \partial u_M(v), \quad t(v) = \langle x(v), v \rangle - u_M(v)$$

Theorem (Rochet and Chone (1998))

$M \leftrightarrow u_M$ is a bijection between menus and convex u_M with $u_M(0) = 0$ and $\partial u_M \in [0, 1]^m$

$m \geq 2$ items, $n = 1$ agent: optimal menus

- a menu $M \subset \mathbb{R}_+ \times [0, 1]^m$
- utility obtained by an agent with values $v = (v_1, \dots, v_m)$:

$$u_M(v) = \max_{(t, x) \in M} \langle x, v \rangle - t,$$

- u_M is convex and

$$x(v) = \partial u_M(v), \quad t(v) = \langle x(v), v \rangle - u_M(v)$$

Theorem (Rochet and Chone (1998))

$M \leftrightarrow u_M$ is a bijection between menus and convex u_M with $u_M(0) = 0$ and $\partial u_M \in [0, 1]^m$

Revenue maximization becomes:

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) \, dv.$$

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) \, dv$$

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) \, dv =$$

[integrating by parts]

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) \, dv =$$

[integrating by parts]

$$= \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} u(v) \, d\psi,$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) \, dv$ (not necessary positive!)

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} u(v) \, d\psi$$

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} u(v) \, d\psi$$

What is the dual?

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} u(v) \, d\psi$$

What is the dual?

Definition: 2nd-order stochastic dominance aka majorization

$$\mu \succeq \nu \iff \int g \, d\mu \geq \int g \, d\nu \text{ for any convex monotone } g$$

$m \geq 2$ items, $n = 1$ agent: optimal menus and transportation

$$R_m(\rho) = \max_{\substack{\text{convex } u \\ u(0) = 0, \partial u \in [0, 1]^m}} \int_{\mathbb{R}_+^m} u(v) \, d\psi$$

What is the dual?

Definition: 2nd-order stochastic dominance aka majorization

$$\mu \succeq \nu \iff \int g \, d\mu \geq \int g \, d\nu \text{ for any convex monotone } g$$

Theorem (Daskalakis et al (2017))

$$R_m(\rho) = \min_{\substack{\text{positive measures } \gamma \\ \text{on } \mathbb{R}_+^m \times \mathbb{R}_+^m \\ \gamma_1 - \gamma_2 \succeq \psi}} \int_{\mathbb{R}_+^m \times \mathbb{R}_+^m} \|v - v'\|_1 \, d\gamma(v, v')$$

This is Monge-Kantorovich problem with majorization

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach?

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach? To some extent.

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach? To some extent.

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach? To some extent.

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) *Implementation of reduced form mechanisms* ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach? To some extent.

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

New feasibility constraint

u corresponds to a symmetric n -agent mechanism if and only if

$$\partial_{v_i} u(v) \preceq z^{n-1} \quad \forall i = 1, \dots, m,$$

where $v \sim \rho$ and $z \sim \text{Uniform}([0, 1])$.

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: reduction to a single agent

Goal: maximize revenue over BIC, IR, symmetric n -agent mechanisms

Can we use the same approach? To some extent.

- **Reduced-forms mechanism:** expected allocation and payment of an agent as a function of her vector of values
- As before, one-agent mechanism \leftrightarrow convex u

New feasibility constraint

u corresponds to a symmetric n -agent mechanism if and only if

$$\partial_{v_i} u(v) \preceq z^{n-1} \quad \forall i = 1, \dots, m,$$

where $v \sim \rho$ and $z \sim \text{Uniform}([0, 1])$.

- $m = 1$ proved by Hart and Reny²; equivalent to earlier result by K.Border. General case follows, e.g., from Kleiner et al. (2021)

²S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

$m \geq 2$ items, $n \geq 2$ agents: primal problem

multi-bidder version of Rochet-Chone theorem

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1} \forall i}} n \cdot \int_{\mathbb{R}_+^m} u(v) \, d\psi(v),$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) \, dv$.

- Looks similar to one-agent case
- **Major obstacle:** Local feasibility constraint $\partial u \in [0, 1]^m$ is replaced by a non-local non-linear majorization constraint on gradient's distribution. Cannot get rid of u 's derivatives.

$m \geq 2$ items, $n \geq 2$ agents: primal problem

multi-bidder version of Rochet-Chone theorem

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1} \forall i}} n \cdot \int_{\mathbb{R}_+^m} u(v) \, d\psi(v),$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) \, dv$.

- Looks similar to one-agent case
- **Major obstacle:** Local feasibility constraint $\partial u \in [0, 1]^m$ is replaced by a non-local non-linear majorization constraint on gradient's distribution. Cannot get rid of u 's derivatives.

$m \geq 2$ items, $n \geq 2$ agents: primal problem

multi-bidder version of Rochet-Chone theorem

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1} \forall i}} n \cdot \int_{\mathbb{R}_+^m} u(v) \, d\psi(v),$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) \, dv$.

- Looks similar to one-agent case
- **Major obstacle:** Local feasibility constraint $\partial u \in [0, 1]^m$ is replaced by a non-local non-linear majorization constraint on gradient's distribution. Cannot get rid of u 's derivatives.

$m \geq 2$ items, $n \geq 2$ agents: primal problem

multi-bidder version of Rochet-Chone theorem

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1} \forall i}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v),$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) dv$.

- Looks similar to one-agent case
- **Major obstacle:** Local feasibility constraint $\partial u \in [0, 1]^m$ is replaced by a non-local non-linear majorization constraint on gradient's distribution. Cannot get rid of u 's derivatives.

What is the dual?

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- **Beckmann's problem:**

$$\text{Beck}_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$$

- **The choice of costs:**

for convex monotone φ_i on \mathbb{R}_+ with $\varphi_i(0) = 0$ define

$$\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|) \quad \text{where } \varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$$

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- **Beckmann's problem:**

$$\text{Beck}_\rho(\pi, \Phi) = \min_{f: \operatorname{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$$

- **The choice of costs:**

for convex monotone φ_i on \mathbb{R}_+ with $\varphi_i(0) = 0$ define

$$\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|) \quad \text{where } \varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$$

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- **Beckmann's problem:**

$$\text{Beck}_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$$

- **The choice of costs:**

for convex monotone φ_i on \mathbb{R}_+ with $\varphi_i(0) = 0$ define

$$\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|) \quad \text{where } \varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$$

Strong duality

$$R_{n,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- $\text{Beck}_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$
- $\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|)$ where $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$

Strong duality

$$R_{n,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

- **weak duality (\leq) is easy:** see the blackboard
 - gives complementary slackness conditions [link](#)
- **ingredients for strong duality:** careful choice of topology, a priori estimates on (u, φ_i) , functional minimax theorem
- **the existence of optimal vector field f :** allow for generalized fields (vector measures) and use weak derivatives

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- $\text{Beck}_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$
- $\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|)$ where $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$

Strong duality

$$R_{n,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

- **weak duality (\leq) is easy:** see the blackboard
 - gives complementary slackness conditions [link](#)
- **ingredients for strong duality:** careful choice of topology, a priori estimates on (u, φ_i) , functional minimax theorem
- **the existence of optimal vector field f :** allow for generalized fields (vector measures) and use weak derivatives

$m \geq 2$ items, $n \geq 2$ agents: dual problem

- $\text{Beck}_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv.$
- $\Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|)$ where $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$

Strong duality

$$R_{n,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

- **weak duality (\leq) is easy:** see the blackboard
 - gives complementary slackness conditions [link](#)
- **ingredients for strong duality:** careful choice of topology, a priori estimates on (u, φ_i) , functional minimax theorem
- **the existence of optimal vector field f :** allow for generalized fields (vector measures) and use weak derivatives

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals:
Monge-Kantorovich and Beckmann?

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals:
Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right]$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(\mathbf{1}) \, dz \right]$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \varphi_i(1) \right]$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \varphi_i(1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \varphi_i(1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \varphi_i(1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[\text{Beck}_\rho(\pi, \Phi) + \sum_{i=1}^m \varphi_i(1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \preceq \psi} \left[\text{Beck}_\rho(\pi, \Phi) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \preceq \psi} \left[\text{Beck}_\rho(\pi, \Phi) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \preceq \psi} \left[\text{Beck}_{\rho}(\pi, \Phi) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\text{Beck}_\rho \left(\pi, \|\cdot\|_1 \right) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\text{Beck}_\rho \left(\pi, \|\cdot\|_1 \right) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\text{Beck}_\rho(\pi, \|\cdot\|_1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\text{Beck}_{\rho}(\pi, \|\cdot\|_1) \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_{\rho}(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v') \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\pi \succeq \psi} \left[\min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v') \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\text{positive measures } \gamma \\ \gamma_1 - \gamma_2 \succeq \psi}} \left[\int \|v - v'\|_1 d\gamma(v, v') \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

$$R_{n=1,m}(\rho) = \min_{\substack{\text{positive measures } \gamma \\ \gamma_1 - \gamma_2 \succeq \psi}} \left[\int \|v - v'\|_1 d\gamma(v, v') \right]$$

- if we increase $\varphi_i(x)$, then $\varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$ decreases
- thus $\varphi_i(x) = x \cdot \varphi_i(1)$ in $[0, 1]$ and $\varphi_i(x) = +\infty$ for $x \geq 1$
- optimization over $\varphi_i(1)$ gives $\varphi_i(1) = 0$
- $\varphi_i^*(y) = y$ and so $\Phi = \|\cdot\|_1$

Theorem (Santambrogio (2015))

$$\text{Beck}_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

Corollary: duality by Daskalakis et al. (2017)

Why strong duality is useful?

- **Upper bound on revenue**
- **Controlling how far a given mechanism is from the optimum:**
numerical methods with provable approximation guarantees
- **Complementary slackness conditions**
 - can be used to show that a mechanism is not optimal if the conditions are infeasible.
Example: For $p(v) = p_1(v_1) \cdot \dots \cdot p_m(v_m)$, selling separately is never optimal.³
 - help to guess/construct an explicit solution and to prove its optimality (dual solution is a certificate)
Example: For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.⁴

³Obtained by P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) Mixed bundling auctions JET

⁴Obtained by A.Manelli, D.Vincent (2007) Multidimensional Mechanism Design JET

Why strong duality is useful?

- **Upper bound on revenue**
- **Controlling how far a given mechanism is from the optimum:**
numerical methods with provable approximation guarantees
- **Complementary slackness conditions**
 - can be used to show that a mechanism is not optimal if the conditions are infeasible.
Example: For $p(v) = p_1(v_1) \cdot \dots \cdot p_m(v_m)$, selling separately is never optimal.³
 - help to guess/construct an explicit solution and to prove its optimality (dual solution is a certificate)
Example: For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.⁴

³Obtained by P. Jehiel, M. Meyer-Ter-Vehn, B. Moldovanu (2007) *Mixed bundling auctions* JET

⁴Obtained by A. Manelli, D. Vincent (2007) *Multidimensional Mechanism Design* JET

Why strong duality is useful?

- **Upper bound on revenue**
- **Controlling how far a given mechanism is from the optimum:**
numerical methods with provable approximation guarantees
- **Complementary slackness conditions**
 - can be used to show that a mechanism is not optimal if the conditions are infeasible.
Example: For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.³
 - help to guess/construct an explicit solution and to prove its optimality (dual solution is a certificate)
Example: For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.⁴

³Obtained by P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

⁴Obtained by A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Why strong duality is useful?

- Upper bound on revenue
- Controlling how far a given mechanism is from the optimum: numerical methods with provable approximation guarantees
- Complementary slackness conditions
 - can be used to show that a mechanism is not optimal if the conditions are infeasible.
Example: For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.³
 - help to guess/construct an explicit solution and to prove its optimality (dual solution is a certificate)
Example: For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.⁴

³Obtained by P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

⁴Obtained by A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Why strong duality is useful?

- Upper bound on revenue
- Controlling how far a given mechanism is from the optimum: numerical methods with provable approximation guarantees
- Complementary slackness conditions
 - can be used to show that a mechanism is not optimal if the conditions are infeasible.
Example: For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.³
 - help to guess/construct an explicit solution and to prove its optimality (dual solution is a certificate)
Example: For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.⁴

³Obtained by P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

⁴Obtained by A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Numerical simulations


Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items?

Numerical simulations

Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items? **Perhaps, not.**

Numerical simulations


Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items? **Perhaps, not.**



Probability to receive the first item as a function of bidder's values (v_1, v_2) in the optimal auction

Numerical simulations

Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items? **Perhaps, not.**



Probability to receive the first item as a function of bidder's values (v_1, v_2) in the optimal auction

Remark: computing the optimum numerically is a non-trivial task requiring extra optimal transportation insights

[about algorithm](#)

- Another justification that mechanism design \simeq optimal transport
- Surprising appearance of Beckmann's problem, its first non-transportation application
- **Remaining question to unleash full power:** Guess a good candidate for an explicit solution? (for some particular ρ)

- Another justification that mechanism design \simeq optimal transport
- Surprising appearance of Beckmann's problem, its first non-transportation application
- **Remaining question to unleash full power:** Guess a good candidate for an explicit solution? (for some particular ρ)

Thank you!

Optimal u^{opt} , functions φ_i^{opt} , measure π^{opt} , and vector field f^{opt} satisfy:

$$\int u^{\text{opt}}(v) \, d\psi(v) = \int u^{\text{opt}}(v) \, d\pi^{\text{opt}}(v)$$

$$f_i^{\text{opt}}(v) \in \partial \varphi_i^{\text{opt}} \left(\frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right)$$

$$\int \varphi_i^{\text{opt}} \left(\frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right) \rho(v) \, dv = \int_0^1 \varphi_i^{\text{opt}}(z^{n-1}) \, dz$$

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ revenue curve

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$



Revenue as a function of the number of bidders n for two items with i.i.d. values uniform on $[0, 1]$. Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for $n \rightarrow \infty$ (the dashed line).

Remark: For $n = 2$, selling optimally improves upon selling separately by 5%



Revenue as a function of the number of bidders n for two items with i.i.d. values uniform on $[0, 1]$. Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for $n \rightarrow \infty$ (the dashed line).

Remark: For $n = 2$, selling optimally improves upon selling separately by 5%