

Beckmann's approach to multi-item multi-bidder auctions

arXiv:2203.06837

Alexander V. Kolesnikov (HSE)

Fedor Sandomirskiy (Caltech, fsandomi@caltech.edu, <https://www.fedors.info/>)

Aleh Tsyvinski (Yale)

Alexander P. Zimin (MIT)

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$		
$n \geq 2:$		

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	Single-item monopolist Myerson (1981): posted price is optimal	
$n \geq 2:$		

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	Single-item monopolist Myerson (1981): posted price is optimal	
$n \geq 2:$	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport^a <hr/> ^a C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport^a <hr/> ^a C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	Multi-item auctions <ul style="list-style-type: none">• almost nothing known about optimal mechanisms^a• Our paper: connection to optimal transport <hr/> ^a Combined obstacles of multidimensional and multi-agent mechanism design

Most profitable way to sell m unrelated items to n agents

Bayesian setting: independent private values, seller knows distribution

What is known?

	$m = 1$:	$m \geq 2$:
$n = 1$:	Single-item monopolist Myerson (1981): posted price is optimal	Multi-item monopolist <ul style="list-style-type: none">• optimal mechanisms known in particular cases• connection to optimal transport^a <hr/> ^a C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) Strong duality for a multiple-good monopolist Econometrica
$n \geq 2$:	Classic auctions Myerson (1981): 2nd-price auction with reserve is optimal	Multi-item auctions <ul style="list-style-type: none">• almost nothing known about optimal mechanisms^a• Our paper: connection to optimal transport <hr/> ^a Combined obstacles of multidimensional and multi-agent mechanism design

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

What will we see?

Strong duality (informal)

For $n \geq 1$ bidders with additive utilities over $m \geq 1$ items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive \Rightarrow discuss the right-hand side

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM. Beckmann (1952) **A continuous model of transportation** *Econometrica*

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM. Beckmann (1952) **A continuous model of transportation** *Econometrica*

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

Transportation problem

For given geographical distribution of production $\pi_+ \in \Delta(X)$ and consumption $\pi_- \in \Delta(X)$, $X \subset \mathbb{R}^d$, find least costly way of transportation

Monge-Kantorovich problem

Given:

π_+, π_- and transport costs $c(x, y)$

Find:

transportation plan γ solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for $n = 1$ bidder
- transportation is immediate

Beckmann's problem^a

Given:

π_+, π_- , costs $\Phi(x)$

Find:

flow $f: X \rightarrow \mathbb{R}^n$ solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

^aM.Beckmann (1952) **A continuous model of transportation** Econometrica

- **Econ applications of optimal transport**
 - Monge-Kantorovich: Daskalakis et al. (2017), Kleiner, Manelli (2019), Boerma et al.(2021), Chiapporiet et al. (2010), Galichon (2021), Steinerberger, Tsyvinski (2019), Gensbittel (2015), Arieli et al.,(2022), Guo, Shmaya (2021)
 - Beckmann: Fajgelbaum, Schaal (2020), Allenand, Arkolakis (2014), Santambrogio (2015)
- **Non-transport duality in auction design** Giannakopoulos, Koutsoupas (2018), Cai et al. (2019), Bergemann et al. (2016)
- **Simple mechanisms with good revenue guarantees** Hart, Reny (2019), Haghpasand, Hartline (2021), Babaioff et al. (2020,2021), Hart, Nisan (2017), Jehiel et al. (2007), Yao (2017), and many more...
- **Majorization in economics** Hart and Reny (2015), Kleiner et al. (2021), Arieli et al. (2019), Candogan, Strack (2021), Nikzad (2022)

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\text{optimal revenue} = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv$$

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv = \\ & \left[\text{integrating by parts} \right] \end{aligned}$$

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\text{optimal revenue} = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left(\langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv =$$

[integrating by parts]

$$= \max_{\substack{\text{convex } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi,$$

where $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) dv$ (signed measure!)

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

What is the dual?

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

What is the dual?

Definition: 2nd-order stochastic dominance aka **majorization**

$$\mu \succeq \nu \iff \int g d\mu \geq \int g d\nu \text{ for any convex monotone } g$$

Known results: $m \geq 2$ goods, $n = 1$ agent

- agent's values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms \Leftrightarrow interim utility functions

Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

Definition: 2nd-order stochastic dominance aka **majorization**

$$\mu \succeq \nu \iff \int g d\mu \geq \int g d\nu \text{ for any convex monotone } g$$

Theorem (Daskalakis et al (2017))

$$\begin{aligned} \text{optimal revenue} = & \min_{\substack{\text{positive measures } \gamma \\ \text{on } \mathbb{R}_+^m \times \mathbb{R}_+^m \\ \gamma_1 - \gamma_2 \succeq \psi}} \int_{\mathbb{R}_+^m \times \mathbb{R}_+^m} \|v - v'\|_1 d\gamma(v, v') \end{aligned}$$

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} &= n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution
- **Ingredients:**
 - **reduction:** n -agent mechanism \rightarrow 1-agent reduced form
 - characterization of feasible reduced forms via majorization:

$m = 1$ proved by Hart and Reny¹, equivalent to Border's theorem

¹S.Hart, P.Reny (2015) *Implementation of reduced form mechanisms* ET Bulletin

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution
- **Ingredients:**
 - **reduction:** n -agent mechanism \rightarrow 1-agent reduced form
 - **characterization of feasible reduced forms via majorization:**
 $m = 1$ proved by Hart and Reny¹, equivalent to Border's theorem

¹S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- n i.i.d agents with values $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

What is the dual?

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) \, d\psi \end{aligned}$$

What is the dual?

$$\text{Beckmann: } B_\rho(\pi, \Phi) = \min_{f: \operatorname{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv$$

Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) \, d\psi \end{aligned}$$

$$\text{Beckmann: } B_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv$$

Theorem (strong duality)

$$\begin{aligned} \text{optimal revenue} = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} \left[B_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right], \end{aligned}$$

$$\text{where } \Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|) \quad \text{and} \quad \varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

Beckmann's dual simplifies:

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_{\rho} \left(\pi, \| \cdot \|_1 \right)$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

Beckmann's dual simplifies:

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_{\rho}(\pi, \|\cdot\|_1)$$

Theorem (Santambrogio (2015))

$$B_{\rho}(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

The case for $n = 1$ agent

Question: How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

Beckmann's dual simplifies:

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_{\rho}(\pi, \|\cdot\|_1)$$

Theorem (Santambrogio (2015))

$$B_{\rho}(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

Corollary: duality by Daskalakis et al. (2017)

Strong duality \Rightarrow complementary slackness conditions

- allow to **disprove** optimality
 - **Example:** For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.¹
- help to **guess** an explicit solution and to **prove optimality**
 - **Example:** For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.²

¹P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

²A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Strong duality \Rightarrow complementary slackness conditions

- allow to **disprove** optimality
 - **Example:** For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.¹
- help to **guess** an explicit solution and to **prove optimality**
 - **Example:** For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.²

¹P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

²A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Strong duality \Rightarrow complementary slackness conditions

- allow to **disprove** optimality
 - **Example:** For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.¹
- help to **guess** an explicit solution and to **prove optimality**
 - **Example:** For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.²

¹P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

²A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Strong duality \Rightarrow complementary slackness conditions

- allow to **disprove** optimality
 - **Example:** For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.¹
- help to **guess** an explicit solution and to **prove optimality**
 - **Example:** For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.²

Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items?

¹P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

²A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

Strong duality \Rightarrow complementary slackness conditions

- allow to **disprove** optimality
 - **Example:** For $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$, selling separately is never optimal.¹
- help to **guess** an explicit solution and to **prove optimality**
 - **Example:** For $n = 1$ and $m = 2$ i.i.d. uniform items, selling each for $\frac{2}{3}$ or both for $\frac{4-\sqrt{2}}{3}$ is optimal.²


Question: Any hope for an explicit solution with $n \geq 2$ and $m = 2$ i.i.d. uniform items? **Perhaps, not**

¹P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

²A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET


Pictures for dessert: 2 bidders, 2 i.i.d. uniform items

Probability to receive the first item as a function of bidder's values (v_1, v_2) in the optimal auction (about algorithm):



Pictures for dessert: 2 bidders, 2 i.i.d. uniform items

Probability to receive the first item as a function of bidder's values (v_1, v_2) in the optimal auction (about algorithm):



Thank you!

Optimal u^{opt} , functions φ_i^{opt} , measure π^{opt} , and vector field f^{opt} satisfy:

$$\int u^{\text{opt}}(v) \, d\psi(v) = \int u^{\text{opt}}(v) \, d\pi^{\text{opt}}(v)$$

$$f_i^{\text{opt}}(v) \in \partial \varphi_i^{\text{opt}} \left(\frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right)$$

$$\int \varphi_i^{\text{opt}} \left(\frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right) \rho(v) \, dv = \int_0^1 \varphi_i^{\text{opt}}(z^{n-1}) \, dz$$

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$ [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of n agents can have q different values for each of m items \Rightarrow the dimension $\sim (q^n)^m$
 - intractable for $(m = 2, q = 100, n = 2)$ or for $(m = 2, q = 10, n = 4)$
 - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on n is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
 - μ on $[0, 1]$ majorizes ν if and only if there is γ on $[0, 1]^2$ with marginals μ on y and ν on x and such that $\int y d\gamma(y | x) \geq x$ for γ -almost all x
 - solve for (u, γ)
- **Performance:** algorithm handles $(m = 2, q = 100, n = 10)$



Revenue as a function of the number of bidders n for two items with i.i.d. values uniform on $[0, 1]$. Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for $n \rightarrow \infty$ (the dashed line).

Remark: For $n = 2$, selling optimally improves upon selling separately by 5%



Revenue as a function of the number of bidders n for two items with i.i.d. values uniform on $[0, 1]$. Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for $n \rightarrow \infty$ (the dashed line).

Remark: For $n = 2$, selling optimally improves upon selling separately by 5%