

# Improvable Equilibria

---

**Kirill Rudov** - UC Berkeley

**Fedor Sandomirskiy** - Princeton

**Leeat Yariv** - Princeton

Harvard-MIT theory seminar, February 20, 2025

Correlated Equilibria (CE) (Aumann, 1974) generalize Nash equilibria to allow correlation

- capture pre-play communication, intermediation, and joint randomization  
Bárány (1992), Ben-Porath (1998), Gerardi (2004), Lehrer and Sorin (1997)
- result from natural learning dynamics  
Foster and Vohra (1997), Fudenberg and Levine (1999), Hart and Mas-Colell (2000)
- efficiently computable  
Papadimitriou and Roughgarden (2008)

**Broad question:** When is there potential value in correlation?

CE  $\simeq$  adding a recommendation system on top of the existing interaction

- $\implies$  What interactions can be improved by a recommendation system?

CE  $\simeq$  adding a recommendation system on top of the existing interaction

- $\implies$  What interactions can be improved by a recommendation system?

CE  $\simeq$  outcomes of arbitrary pre-play communication protocols

- $\implies$  What strategic interactions are susceptible to communication / collusion?

## Normal-form game

$$\Gamma = \left( N, (A_i)_{i \in N}, (u_i: A \rightarrow \mathbb{R})_{i \in N} \right)$$

- $N = \{1, \dots, n\}$  is finite set of players
- $A_i$  is a finite set of actions of player  $i$
- $A = \prod_{i \in N} A_i$  is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$  is utility of player  $i$

**Definition (Aumann, 1974, 1987)**

A distribution  $\mu \in \Delta(A)$  is a CE if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all  $i \in N$  and all  $a_i, a'_i \in A_i$

**Interpretation:**  $\mu$  generated by a mediator and players best respond by adhering

**Remark:** Nash Equilibria (NE) are CE of the form  $\mu = \mu_1 \times \dots \times \mu_n$

## Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

# Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

## Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE



# Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**


## Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

**Our Question:** When is a Nash equilibrium extreme?

# Linear objectives and extreme points

Maximization of a linear objective over a polytope:



- If the optimum is unique, it is an extreme point
- In knife-edge cases, the whole face can be optimal

## Definition

Objectives with unique optima are **non-degenerate**

- Tiny perturbations can make degenerate non-degenerate

# Improvability of non-extreme equilibria

## Observation

NE is non-extreme  $\iff$  any non-degenerate linear objective can be improved

## Observation

NE is non-extreme  $\iff$  any non-degenerate linear objective can be improved

- Linear in probabilities, not in actions  $\Rightarrow$  a broad class of objectives
  - e.g., utilitarian welfare, revenue, maximizing/minimizing the probability of a particular action profile

## Observation

NE is non-extreme  $\iff$  any non-degenerate linear objective can be improved

- Linear in probabilities, not in actions  $\Rightarrow$  a broad class of objectives
  - e.g., utilitarian welfare, revenue, maximizing/minimizing the probability of a particular action profile
- The conclusion extends to convex objectives (**Bauer's maximum principle**)
  - including quasi-convex (Ball, 2023)

## Improvability of non-extreme equilibria 2

- Non-extreme equilibria are improvable **no matter** the objective
- A conservative notion, agnostic to the designer's objective
- Usually, assess outcomes for a **given** objective

## Improvability of non-extreme equilibria 2

- Non-extreme equilibria are improvable **no matter** the objective
- A conservative notion, agnostic to the designer's objective
- Usually, assess outcomes for a **given** objective

### Main Insight

Despite restrictiveness of improvability notion, **many** equilibria are improvable, i.e., **non-extreme**

- **Value of correlation in 2-player games:** Cripps (1995), Evangelista and Raghavan (1996), Canovas, Hansen, and Jaumard (1999), Nau, Canovas, and Hansen (2004), Peeters and Potters (1999), Calvó-Armengol (2006), Ashlagi, Monderer, and Tennenholtz (2008)
- **Extreme-point approach in info & mech. design:** Manelli and Vincent (2007), Kleiner, Moldovanu, and Strack (2021), Arieli, Babichenko, Smorodinsky, and Yamashita (2023), Yang and Zentefis (2024), Nikzad (2022), Kleiner, Moldovanu, Strack, and Whitmeyer (2024), Lahr and Niemeyer (2024)



- **Conditions for extremality:**  
in the space of action distributions and payoff space
- **Particular classes of games:**  
symmetric, having unique CE
- **Extensions:**  
Bayesian CE and Coarse CE

## Conditions for Extremality

---

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme  
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme  
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness

## Theorem

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme  
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness  
 $\Rightarrow$  2-player games not representative



# Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...

## Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

# Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize

# Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)

# Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize


- In a generic game, any NE is regular (Harsanyi, 1973)
- Hence, Theorem 1'  $\Rightarrow$  Theorem 1

## **Example: 2 Players vs 3 Players**

---

## Example: 2-Player Games

A version of the Game of Chicken by **Aumann (1974)**:



	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9

## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players

Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$



## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players  
Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from  $(6,6)$

## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players

Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**


## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players  
Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**
- Indeed, it is the optimum for a non-degenerate objective


weight of (Risky, Risky) & (Safe, Safe)  $\rightarrow \max$

## Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

## Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

## Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

## Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

## Example: 3-Player Games




		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player




## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation in  $p \Rightarrow$  irrational weights (Nash, 1950)


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
Safe		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation in  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation in  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates
- The mixed NE is **not extreme**

## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
Safe				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation in  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates
- The mixed NE is **not extreme**

More than 2 players mixing makes a difference...

## General Proof Intuition

---

**High-level idea:** When many players randomize, there are too many ways to correlate their actions  $\implies$  one must be beneficial

Focus on a particular example to illustrate

- Game with  $n$  players, each with 2 actions

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$



- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq k + 1$

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq k + 1$
- $\Rightarrow$  support of an extreme CE  $\mu$  is bounded by  $2n + 1$

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing

## General Proof Intuition

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles

## General Proof Intuition

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- We can replace  $2n + 1$  with  $2k + 1$  by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- We can replace  $2n + 1$  with  $2k + 1$  by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

**Conclusion:** NE with  $k \geq 3$  mixing agents cannot be extreme



- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- We can replace  $2n + 1$  with  $2k + 1$  by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

**Conclusion:** NE with  $k \geq 3$  mixing agents cannot be extreme

- The same argument applies to equilibria where players mix over the same number of pure strategies

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- We can replace  $2n + 1$  with  $2k + 1$  by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

**Conclusion:** NE with  $k \geq 3$  mixing agents cannot be extreme

- The same argument applies to equilibria where players mix over the same number of pure strategies
- The main difficulty is handling very asymmetric equilibria [▶ details](#)

## Extreme Points in Payoff Space

---

- The set of CE  $\subset \Delta(A)$  subset of a space of dimension  $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in  $\mathbb{R}^n$

## Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

- The set of CE  $\subset \Delta(A)$  subset of a space of dimension  $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in  $\mathbb{R}^n$

## Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

**Question:** What can we say about payoff-extreme equilibria?

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

- Projection of an extreme point **need not** be an extreme point of a projection



## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

- Projection of an extreme point **need not** be an extreme point of a projection
- $\Rightarrow$  pure NE and NE with 2 mixers **need not** be payoff-extreme
  - e.g, the mixed NE in the Game of Chicken

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \beta_i \sum_{a \in A} u_i(a) \mu(a) \rightarrow \max$$

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \beta_i \sum_{a \in A} u_i(a) \mu(a) \rightarrow \max$$

- The case  $\beta_1 = \dots = \beta_n = 1$  corresponds to **utilitarian welfare**

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \beta_i \sum_{a \in A} u_i(a) \mu(a) \rightarrow \max$$

- The case  $\beta_1 = \dots = \beta_n = 1$  corresponds to **utilitarian welfare**
- Non-degeneracy means unique optimum

# Payoff-extreme Equilibria

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \beta_i \sum_{a \in A} u_i(a) \mu(a) \rightarrow \max$$

- The case  $\beta_1 = \dots = \beta_n = 1$  corresponds to **utilitarian welfare**
- Non-degeneracy means unique optimum

## Proposition

In a generic game, utilitarian welfare is non-degenerate

## **Applications to Particular Classes of Games**

---

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$



## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

**Other Applications:** games where players want to mismatch actions of others

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

**Other Applications:** games where players want to mismatch actions of others

- e.g., network games (with substitutes), congestion games, all-pay auctions, Boston matching mechanism

# Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

# Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

**Take-away:** caution when focusing on symmetric mixed equilibria in symmetric games



## What Extreme CE Look Like

---

► skip

## What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?

# What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?

- For general games, we only know that extreme CE have small support
- For symmetric games and symmetric CE, we can say more

**Observation:**

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable

**Observation:**

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known

## Observation:

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known

## Theorem (de Finetti)

Any infinite exchangeable sequence  $a_1, a_2, a_3 \dots$  is a mixture of i.i.d. distributions

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### **Proposition 2**

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions



## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

# Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

- A version of Proposition 2 holds: sampling without replacement instead of i.i.d.

# Games with Unique Correlated Equilibrium

---

► skip

# Games with Unique Correlated Equilibrium

- Unique CE  $\implies$  CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

# Games with Unique Correlated Equilibrium

- Unique CE  $\implies$  CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

## Corollary

If a game has a unique CE, then it is either:

- a pure Nash, or
- a Nash where exactly two players randomize

# Games with Unique Correlated Equilibrium

- Unique CE  $\implies$  CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

## Corollary

If a game has a unique CE, then it is either:

- a pure Nash, or
  - a Nash where exactly two players randomize
- 
- No genericity assumption since games with a unique CE form an open set (Viossat, 2010)



# Bayesian Correlated Equilibria

---

► skip

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states  $\Theta$
- Private information: finite sets of types  $T_i$
- Prior  $\pi \in \Delta(\Theta \times T)$

# Bayesian Correlated Equilibria

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states  $\Theta$
- Private information: finite sets of types  $T_i$
- Prior  $\pi \in \Delta(\Theta \times T)$

## Definition

A distribution  $\psi \in \Delta(A \times \Theta \times T)$  is a BCE if

- its marginal on  $\Theta \times T$  coincides with  $\pi$
- no agent can gain by deviating from a recommended action  $a_i$  to another action  $a'_i$ , given her private type  $t_i$

$$\sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a, \theta, t) \geq \sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a'_i, a_{-i}, \theta, t)$$

# Bayesian Correlated Equilibria

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states  $\Theta$
- Private information: finite sets of types  $T_i$
- Prior  $\pi \in \Delta(\Theta \times T)$

## Definition

A distribution  $\psi \in \Delta(A \times \Theta \times T)$  is a BCE if

- its marginal on  $\Theta \times T$  coincides with  $\pi$
- no agent can gain by deviating from a recommended action  $a_i$  to another action  $a'_i$ , given her private type  $t_i$

$$\sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a, \theta, t) \geq \sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a'_i, a_{-i}, \theta, t)$$

A Bayesian Nash equilibrium (BNE) is a BCE where  $\theta, a_1, \dots, a_n$  are independent, conditional on  $t_1, \dots, t_n$

## Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ( $|\Theta| \geq 2$ ), or
- non-trivial private information ( $|T_i| \geq 2$  for at least 3 agents),

a BNE is an extreme point of BCE  $\Leftrightarrow$  it is pure

## Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ( $|\Theta| \geq 2$ ), or
- non-trivial private information ( $|T_i| \geq 2$  for at least 3 agents),

a BNE is an extreme point of BCE  $\Leftrightarrow$  it is pure

- Even minimal uncertainty—e.g., a single binary state—can be sufficient to render mixed BNE non-extreme
  - Contrast with complete information games, where two agents can mix without losing extremality

## Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ( $|\Theta| \geq 2$ ), or
- non-trivial private information ( $|T_i| \geq 2$  for at least 3 agents),

a BNE is an extreme point of BCE  $\Leftrightarrow$  it is pure

- Even minimal uncertainty—e.g., a single binary state—can be sufficient to render mixed BNE non-extreme
  - Contrast with complete information games, where two agents can mix without losing extremality

**Intuition:** Randomness in the state and private types provides more degrees of freedom for correlation to be beneficial

# Coarse Correlated Equilibria

---

► skip



## Coarse Correlated Equilibria

Consider situations where agents commit **ex-ante** to a correlating device, before receiving recommendations. For example,

- firms entering binding collusive agreements (McAfee and McMillan, 1992)
- users opting in to algorithmic recommendation systems

# Coarse Correlated Equilibria

Consider situations where agents commit **ex-ante** to a correlating device, before receiving recommendations. For example,

- firms entering binding collusive agreements (McAfee and McMillan, 1992)
- users opting in to algorithmic recommendation systems

**Definition (Hannan, 1957; Moulin and Vial, 1978)**

A distribution  $\mu \in \Delta(A)$  is a coarse correlated equilibrium (CCE) if, for all  $i \in N$ ,

$$\sum_{a \in A} \mu(a) u_i(a) \geq \max_{a'_i \in A_i} \sum_{a \in A} u_i(a'_i, a_{-i}) \mu(a),$$

i.e., the expected payoff from following the correlated strategy is at least as high as the best deterministic deviation

- $\text{CCE} \supseteq \text{CE} \supseteq \text{NE}$

### **Proposition**

In a generic game, a NE is an extreme point of the set of CCE  $\Leftrightarrow$  it is pure

## Proposition

In a generic game, a NE is an extreme point of the set of CCE  $\Leftrightarrow$  it is pure

- The tension between randomness and optimality is even stronger for CCE than for CE: any randomness allows for improvement
- Consistent with the prevalence of examples where CCE improves over NE, even in two-player games (Moulin and Vial, 1978; Gérard-Varet and Moulin, 1978; Moulin, Ray, and Gupta, 2014; Dokka, Moulin, Ray, and SenGupta, 2023)

# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a general tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Main takeaway: inherent suboptimality of equilibria with a lot of mixing

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a general tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Main takeaway: inherent suboptimality of equilibria with a lot of mixing

# Thank you!

## Key Lemmas

---



## Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If  $\mu$  is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |S_i| \cdot (|S_i| - 1)$$

## Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium,  $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ :

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$


## Regularity of Generic games (**Harsanyi, 1973**)

In a generic game, any Nash equilibrium is regular


# Simulations

---



# Simulations



(a)  $NE/2u$



(b) Fraction of CE  $> NE^*$



# Extreme Symmetric CE with Any Number of Players

Consider  $n$  players with  $m$  actions each

## Proposition

Any extreme symmetric CE can be obtained as follows:

- there are  $M$  urns, each with  $n$  balls labeled by actions

$$1 \leq M \leq m(m-1) + 1$$

- an urn is selected at random according to  $p \in \Delta_M$ , secretly from players
- players draw balls sequentially without replacement
- $i$ 's action = her ball's label, no incentive to deviate

**Remark:** If  $n$  is large, sampling without replacement can be approximated by i.i.d.

## References

---

- Arieli, I., Y. Babichenko, R. Smorodinsky, and T. Yamashita (2023). Optimal persuasion via bi-pooling. *Theoretical Economics* 18(1), 15–36.
- Ashlagi, I., D. Monderer, and M. Tennenholtz (2008). On the value of correlation. *Journal of Artificial Intelligence Research* 33, 575–613.
- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Aumann, R. J. (1987). Correlated equilibrium as an expression of bayesian rationality. *Econometrica: Journal of the Econometric Society*, 1–18.
- Ball, I. (2023). Bauer’s maximum principle for quasiconvex functions. *arXiv preprint arXiv:2305.04893*.
- Bárány, I. (1992). Fair distribution protocols or how the players replace fortune. *Mathematics of Operations Research* 17(2), 327–340.
- Ben-Porath, E. (1998). Correlation without mediation: Expanding the set of equilibrium outcomes by “cheap” pre-play procedures. *Journal of Economic Theory* 80(1), 108–122.
- Bergemann, D. and S. Morris (2019). Information design: A unified perspective. *Journal of Economic Literature* 57(1), 44–95.
- Calvó-Armengol, A. (2006). The set of correlated equilibria of 2x2 games. *mimeo*.
- Canovas, S. G., P. Hansen, and B. Jaumard (1999). Nash equilibria from the correlated equilibria viewpoint. *International Game Theory Review* 1(01), 33–44.

- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Dokka, T., H. Moulin, I. Ray, and S. SenGupta (2023). Equilibrium design in an n-player quadratic game. *Review of economic design* 27(2), 419–438.
- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Evangelista, F. S. and T. Raghavan (1996). A note on correlated equilibrium. *International Journal of Game Theory* 25, 35–41.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings* 12, pp. 131–144. Springer.
- Foster, D. P. and R. V. Vohra (1997). Calibrated learning and correlated equilibrium. *Games and Economic Behavior* 21(1-2), 40–55.
- Fudenberg, D. and D. K. Levine (1999). Conditional universal consistency. *Games and Economic Behavior* 29(1-2), 104–130.
- Gérard-Varet, L.-A. and H. Moulin (1978). Correlation and duopoly. *Journal of economic theory* 19(1), 123–149.
- Gerardi, D. (2004). Unmediated communication in games with complete and incomplete information. *Journal of Economic Theory* 114(1), 104–131.

- Hannan, J. (1957). Approximation to bayes risk in repeated play. *Contributions to the Theory of Games* 3(2), 97–139.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.
- Hart, S. and A. Mas-Colell (2000). A simple adaptive procedure leading to correlated equilibrium. *Econometrica* 68(5), 1127–1150.
- Kleiner, A., B. Moldovanu, and P. Strack (2021). Extreme points and majorization: Economic applications. *Econometrica* 89(4), 1557–1593.
- Kleiner, A., B. Moldovanu, P. Strack, and M. Whitmeyer (2024). The extreme points of fusions. *arXiv preprint arXiv:2409.10779*.
- Lahr, P. and A. Niemyer (2024). Extreme points in multi-dimensional screening. *arXiv preprint arXiv:2412.00649*.
- Lehrer, E. and S. Sorin (1997). One-shot public mediated talk. *Games and Economic Behavior* 20(2), 131–148.
- Manelli, A. M. and D. R. Vincent (2007). Multidimensional mechanism design: Revenue maximization and the multiple-good monopoly. *Journal of Economic theory* 137(1), 153–185.
- McAfee, R. P. and J. McMillan (1992). Bidding rings. *The American Economic Review*, 579–599.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.



- Moulin, H., I. Ray, and S. S. Gupta (2014). Coarse correlated equilibria in an abatement game. Technical report, Cardiff Economics Working Papers.
- Moulin, H. and J. P. Vial (1978). Strategically zero-sum games: the class of games whose completely mixed equilibria cannot be improved upon. *International Journal of Game Theory* 7, 201–221.
- Nash, J. F. (1950). Non-cooperative games.
- Nau, R., S. G. Canovas, and P. Hansen (2004). On the geometry of nash equilibria and correlated equilibria. *International Journal of Game Theory* 32, 443–453.
- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Nikzad, A. (2022). Constrained majorization: Applications in mechanism design. In *Proceedings of the 23rd ACM Conference on Economics and Computation*, pp. 330–331.
- Palfrey, T. R. and H. Rosenthal (1983). A strategic calculus of voting. *Public Choice* 41(1), 7–53.
- Papadimitriou, C. H. and T. Roughgarden (2008). Computing correlated equilibria in multi-player games. *Journal of the ACM (JACM)* 55(3), 1–29.
- Peeters, R. and J. Potters (1999). On the structure of the set of correlated equilibria in two-by-two bimatrix games.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.

Viossat, Y. (2010). Properties and applications of dual reduction. *Economic theory* 44, 53–68.

Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.

Yang, K. H. and A. K. Zentefis (2024). Monotone function intervals: Theory and applications. *American Economic Review* 114(8), 2239–2270.