

# Beckmann's approach to multi-item multi-bidder auctions

arXiv:2203.06837

---

Alexander V. Kolesnikov (HSE)

Fedor Sandomirskiy (Caltech, [fsandomi@caltech.edu](mailto:fsandomi@caltech.edu), <https://www.fedors.info/>)

Aleh Tsyvinski (Yale)

Alexander P. Zimin (MIT)

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$		
$n \geq 2:$		

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	<b>Single-item monopolist</b> Myerson (1981): posted price is optimal	
$n \geq 2:$		

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1:$	$m \geq 2:$
$n = 1:$	<b>Single-item monopolist</b> Myerson (1981): posted price is optimal	
$n \geq 2:$	<b>Classic auctions</b> Myerson (1981): 2nd-price auction with reserve is optimal	

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1$ :	$m \geq 2$ :
$n = 1$ :	<b>Single-item monopolist</b> Myerson (1981): posted price is optimal	<b>Multi-item monopolist</b> <ul style="list-style-type: none"><li>• optimal mechanisms known in particular cases</li><li>• connection to optimal transport</li></ul>
$n \geq 2$ :	<b>Classic auctions</b> Myerson (1981): 2nd-price auction with reserve is optimal	

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1$ :	$m \geq 2$ :
$n = 1$ :	<b>Single-item monopolist</b>  Myerson (1981): posted price is optimal	<b>Multi-item monopolist</b> <ul style="list-style-type: none"><li>• optimal mechanisms known in particular cases</li><li>• connection to optimal transport<sup>a</sup></li></ul> <hr/> <sup>a</sup> C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) <b>Strong duality for a multiple-good monopolist</b> Econometrica
$n \geq 2$ :	<b>Classic auctions</b>  Myerson (1981): 2nd-price auction with reserve is optimal	

# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1$ :	$m \geq 2$ :
$n = 1$ :	<b>Single-item monopolist</b>  Myerson (1981): posted price is optimal	<b>Multi-item monopolist</b> <ul style="list-style-type: none"><li>• optimal mechanisms known in particular cases</li><li>• connection to optimal transport<sup>a</sup></li></ul> <hr/> <sup>a</sup> C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) <b>Strong duality for a multiple-good monopolist</b> Econometrica
$n \geq 2$ :	<b>Classic auctions</b>  Myerson (1981): 2nd-price auction with reserve is optimal	<b>Multi-item auctions</b> <ul style="list-style-type: none"><li>• almost nothing known about optimal mechanisms<sup>a</sup></li><li>• <b>Our paper:</b> connection to optimal transport</li></ul> <hr/> <sup>a</sup> Combined obstacles of multidimensional and multi-agent mechanism design



# Most profitable way to sell $m$ unrelated items to $n$ agents

**Bayesian setting:** independent private values, seller knows distribution

## What is known?

	$m = 1$ :	$m \geq 2$ :
$n = 1$ :	<b>Single-item monopolist</b>  Myerson (1981): posted price is optimal	<b>Multi-item monopolist</b> <ul style="list-style-type: none"><li>• optimal mechanisms known in particular cases</li><li>• connection to optimal transport<sup>a</sup></li></ul> <hr/> <sup>a</sup> C.Daskalakis, A.Deckelbaum, C.Tzamos (2017) <b>Strong duality for a multiple-good monopolist</b> Econometrica
$n \geq 2$ :	<b>Classic auctions</b>  Myerson (1981): 2nd-price auction with reserve is optimal	<b>Multi-item auctions</b> <ul style="list-style-type: none"><li>• almost nothing known about optimal mechanisms<sup>a</sup></li><li>• <b>Our paper:</b> connection to optimal transport</li></ul> <hr/> <sup>a</sup> Combined obstacles of multidimensional and multi-agent mechanism design

# What will we see?

## Strong duality (informal)

For  $n \geq 1$  bidders with additive utilities over  $m \geq 1$  items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive  $\Rightarrow$  discuss the right-hand side

# What will we see?

## Strong duality (informal)

For  $n \geq 1$  bidders with additive utilities over  $m \geq 1$  items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive  $\Rightarrow$  discuss the right-hand side

# What will we see?

## Strong duality (informal)

For  $n \geq 1$  bidders with additive utilities over  $m \geq 1$  items

$$\max_{\text{BIC IR mechanisms}} \text{Revenue} = \min_{\text{transport flows}} \text{Cost}$$

- formal statement later
- left-hand side is intuitive  $\Rightarrow$  discuss the right-hand side

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$



# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

## Beckmann's problem<sup>a</sup>

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

<sup>a</sup>M. Beckmann (1952) **A continuous model of transportation** *Econometrica*

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\substack{\gamma \in \Delta(X \times X) : \\ \text{marginals } \pi_+, \pi_-}} \int c(x, y) d\gamma(x, y)$$

## Beckmann's problem<sup>a</sup>

**Given:**

$\pi_+, \pi_-$ , costs  $\Phi(x)$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

<sup>a</sup>M. Beckmann (1952) **A continuous model of transportation** *Econometrica*

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

## Beckmann's problem<sup>a</sup>

**Given:**

$\pi_+, \pi_-$ , costs  $\Phi(x)$

**Find:**

**flow**  $f: X \rightarrow \mathbb{R}^n$  solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

<sup>a</sup>M.Beckmann (1952) **A continuous model of transportation** Econometrica

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

## Beckmann's problem<sup>a</sup>

**Given:**

$\pi_+, \pi_-$ , costs  $\Phi(x)$

**Find:**

**flow**  $f: X \rightarrow \mathbb{R}^n$  solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

<sup>a</sup>M.Beckmann (1952) **A continuous model of transportation** Econometrica

# Transportation problem

For given geographical distribution of production  $\pi_+ \in \Delta(X)$  and consumption  $\pi_- \in \Delta(X)$ ,  $X \subset \mathbb{R}^d$ , find least costly way of transportation

## Monge-Kantorovich problem

**Given:**

$\pi_+, \pi_-$  and transport costs  $c(x, y)$

**Find:**

transportation plan  $\gamma$  solving

$$\min_{\gamma \in \Delta(X \times X) : \text{marginals } \pi_+, \pi_-} \int c(x, y) d\gamma(x, y)$$

- archetypal coupling problem, many econ applications
- appears for  $n = 1$  bidder
- transportation is immediate

## Beckmann's problem<sup>a</sup>

**Given:**

$\pi_+, \pi_-$ , costs  $\Phi(x)$

**Find:**

**flow**  $f: X \rightarrow \mathbb{R}^n$  solving

$$\min_{f \text{ s.t.}} \int \Phi(f(x)) dx$$
$$\text{div}[f] = \pi_+ - \pi_-$$

- transportation is continuous
- surprisingly, we get Beckmann not Monge-Kantorovich

<sup>a</sup>M.Beckmann (1952) **A continuous model of transportation** Econometrica



- **Econ applications of optimal transport**
  - Monge-Kantorovich: Daskalakis et al. (2017), Kleiner, Manelli (2019), Boerma et al.(2021), Chiapporiet et al. (2010), Galichon (2021), Steinerberger, Tsyvinski (2019), Gensbittel (2015), Arieli et al.,(2022), Guo, Shmaya (2021)
  - Beckmann: Fajgelbaum, Schaal (2020), Allenand, Arkolakis (2014), Santambrogio (2015)
- **Non-transport duality in auction design** Giannakopoulos, Koutsoupas (2018), Cai et al. (2019), Bergemann et al. (2016)
- **Simple mechanisms with good revenue guarantees** Hart, Reny (2019), Haghpasand, Hartline (2021), Babaioff et al. (2020,2021), Hart, Nisan (2017), Jehiel et al. (2007), Yao (2017), and many more...
- **Majorization in economics** Hart and Reny (2015), Kleiner et al. (2021), Arieli et al. (2019), Candogan, Strack (2021), Nikzad (2022)

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\text{optimal revenue} = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left( \langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv$$

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left( \langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv = \\ & \left[ \text{integrating by parts} \right] \end{aligned}$$

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\text{optimal revenue} = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} \left( \langle \partial u(v), v \rangle - u(v) \right) \rho(v) dv =$$

[integrating by parts]

$$= \max_{\substack{\text{convex } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi,$$

where  $d\psi = ((m+1)\rho(v) + \sum_{j=1}^m v_j \partial_{v_j} \rho) dv$  (signed measure!)

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$



## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

What is the dual?

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

### What is the dual?

**Definition:** 2nd-order stochastic dominance aka **majorization**

$$\mu \succeq \nu \iff \int g d\mu \geq \int g d\nu \text{ for any convex monotone } g$$

## Known results: $m \geq 2$ goods, $n = 1$ agent

- agent with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$  and additive utilities
- **Goal:** maximize revenue over BIC IR mechanisms
- **Rochet-Chone approach:** mechanisms  $\Leftrightarrow$  interim utility functions

### Theorem (Rochet and Chone (1998))

$$\begin{aligned} \text{optimal revenue} = & \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ 1\text{-Lipshitz}}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

**Definition:** 2nd-order stochastic dominance aka **majorization**

$$\mu \succeq \nu \iff \int g d\mu \geq \int g d\nu \text{ for any convex monotone } g$$

### Theorem (Daskalakis et al (2017))

$$\begin{aligned} \text{optimal revenue} = & \min_{\substack{\text{positive measures } \gamma \\ \text{on } \mathbb{R}_+^m \times \mathbb{R}_+^m \\ \gamma_1 - \gamma_2 \succeq \psi}} \int_{\mathbb{R}_+^m \times \mathbb{R}_+^m} \|v - v'\|_1 d\gamma(v, v') \end{aligned}$$

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

### Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

### Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} &= n \cdot \max_{\text{convex monotone } u} \int_{\mathbb{R}_+^m} u(v) d\psi \\ &\quad u(0) = 0, \\ &\quad \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ &\quad z \sim \text{Uniform}([0, 1]) \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

### Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution
- **Ingredients:**
  - **reduction:**  $n$ -agent mechanism  $\rightarrow$  1-agent reduced form
  - characterization of feasible reduced forms via majorization:

$m = 1$  proved by Hart and Reny<sup>1</sup>, equivalent to Border's theorem

<sup>1</sup>S.Hart, P.Reny (2015) *Implementation of reduced form mechanisms* ET Bulletin



## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

### Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

- non-local non-linear majorization constraint on gradient's distribution
- **Ingredients:**
  - **reduction:**  $n$ -agent mechanism  $\rightarrow$  1-agent reduced form
  - **characterization of feasible reduced forms via majorization:**  
 $m = 1$  proved by Hart and Reny<sup>1</sup>, equivalent to Border's theorem

<sup>1</sup>S.Hart, P.Reny (2015) **Implementation of reduced form mechanisms** ET Bulletin

## Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

- $n$  i.i.d agents with values  $v = (v_1, \dots, v_m) \sim \rho(v) dv$
- **Goal:** maximize revenue over BIC IR mechanisms

### Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) d\psi \end{aligned}$$

What is the dual?

# Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

## Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) \, d\psi \end{aligned}$$

What is the dual?

$$\text{Beckmann: } B_\rho(\pi, \Phi) = \min_{f: \operatorname{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv$$

# Multi-bidder case: $m \geq 2$ goods, $n \geq 1$ agents

## Multi-bidder extension of Rochet-Chone representation

$$\begin{aligned} \text{optimal revenue} = n \cdot \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \\ \partial_{v_i} u(v) \preceq z^{n-1} \forall i \\ z \sim \text{Uniform}([0, 1])}} \int_{\mathbb{R}_+^m} u(v) \, d\psi \end{aligned}$$

$$\text{Beckmann: } B_\rho(\pi, \Phi) = \min_{f: \text{div}[\rho \cdot f] + \pi = 0} \int_{\mathbb{R}_+^m} \Phi(f(v)) \cdot \rho(v) \, dv$$

## Theorem (strong duality)

$$\begin{aligned} \text{optimal revenue} = n \cdot \min_{\substack{\pi \succeq \psi \\ \varphi_i \text{ on } \mathbb{R}_+ \text{ s.t.} \\ \text{convex, monotone, } \varphi_i(0) = 0}} & \left[ B_\rho(\pi, \Phi) + \sum_{i=1}^m \int_0^1 \varphi_i(z^{n-1}) \, dz \right], \end{aligned}$$

$$\text{where } \Phi(f) = \sum_{i=1}^m \varphi_i^*(|f_i|) \quad \text{and} \quad \varphi_i^*(y) = \sup_x \langle x, y \rangle - \varphi_i(x)$$

## The case for $n = 1$ agent

**Question:** How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

## The case for $n = 1$ agent

**Question:** How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

**Beckmann's dual simplifies:**

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_{\rho} \left( \pi, \| \cdot \|_1 \right)$$

## The case for $n = 1$ agent

**Question:** How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

**Beckmann's dual simplifies:**

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_{\rho}(\pi, \|\cdot\|_1)$$

**Theorem (Santambrogio (2015))**

$$B_{\rho}(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

## The case for $n = 1$ agent

**Question:** How can it be that seller's problem admits two duals: Monge-Kantorovich and Beckmann?

**Beckmann's dual simplifies:**

$$\text{optimal revenue} = \min_{\pi \succeq \psi} B_\rho(\pi, \|\cdot\|_1)$$

**Theorem (Santambrogio (2015))**

$$B_\rho(\pi, \|\cdot\|_1) = \min_{\substack{\text{positive measures } \gamma \\ \text{with marginals } \pi_+, \pi_-}} \int \|v - v'\|_1 d\gamma(v, v')$$

**Corollary:** duality by Daskalakis et al. (2017)



## Strong duality $\Rightarrow$ complementary slackness conditions

- allow to **disprove** optimality
  - **Example:** For  $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$ , selling separately is never optimal<sup>1</sup>
- help to **guess** an explicit solution and to **prove optimality**
  - **Example:** For  $n = 1$  and  $m = 2$  i.i.d. uniform items, selling each for  $\frac{2}{3}$  or both for  $\frac{4-\sqrt{2}}{3}$  is optimal<sup>2</sup>

---

<sup>1</sup>P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

<sup>2</sup>A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

## Strong duality $\Rightarrow$ complementary slackness conditions

- allow to **disprove** optimality
  - **Example:** For  $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$ , selling separately is never optimal<sup>1</sup>
- help to **guess** an explicit solution and to **prove optimality**
  - **Example:** For  $n = 1$  and  $m = 2$  i.i.d. uniform items, selling each for  $\frac{2}{3}$  or both for  $\frac{4-\sqrt{2}}{3}$  is optimal<sup>2</sup>

---

<sup>1</sup>P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

<sup>2</sup>A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

## Strong duality $\Rightarrow$ complementary slackness conditions

- allow to **disprove** optimality
  - **Example:** For  $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$ , selling separately is never optimal<sup>1</sup>
- help to **guess** an explicit solution and to **prove optimality**
  - **Example:** For  $n = 1$  and  $m = 2$  i.i.d. uniform items, selling each for  $\frac{2}{3}$  or both for  $\frac{4-\sqrt{2}}{3}$  is optimal<sup>2</sup>

---

<sup>1</sup>P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

<sup>2</sup>A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

## Strong duality $\Rightarrow$ complementary slackness conditions

- allow to **disprove** optimality
  - **Example:** For  $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$ , selling separately is never optimal<sup>1</sup>
- help to **guess** an explicit solution and to **prove optimality**
  - **Example:** For  $n = 1$  and  $m = 2$  i.i.d. uniform items, selling each for  $\frac{2}{3}$  or both for  $\frac{4-\sqrt{2}}{3}$  is optimal<sup>2</sup>

**Question:** Any hope for an explicit solution with  $n \geq 2$  and  $m = 2$  i.i.d. uniform items?

---

<sup>1</sup>P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

<sup>2</sup>A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET

## Strong duality $\Rightarrow$ complementary slackness conditions

- allow to **disprove** optimality
  - **Example:** For  $\rho(v) = \rho_1(v_1) \cdot \dots \cdot \rho_m(v_m)$ , selling separately is never optimal<sup>1</sup>
- help to **guess** an explicit solution and to **prove optimality**
  - **Example:** For  $n = 1$  and  $m = 2$  i.i.d. uniform items, selling each for  $\frac{2}{3}$  or both for  $\frac{4-\sqrt{2}}{3}$  is optimal<sup>2</sup>

**Question:** Any hope for an explicit solution with  $n \geq 2$  and  $m = 2$  i.i.d. uniform items? **Perhaps, not**


---

<sup>1</sup>P. Jehiel, M.Meyer-Ter-Vehn, B.Moldovanu (2007) **Mixed bundling auctions** JET

<sup>2</sup>A.Manelli, D.Vincent (2007) **Multidimensional Mechanism Design** JET


# Pictures for dessert: 2 bidders, 2 i.i.d. uniform items

Probability to receive the first item as a function of bidder's values  $(v_1, v_2)$  in the optimal auction (about algorithm):



# Pictures for dessert: 2 bidders, 2 i.i.d. uniform items

Probability to receive the first item as a function of bidder's values  $(v_1, v_2)$  in the optimal auction ( [about algorithm](#) ):



Thank you!

Optimal  $u^{\text{opt}}$ , functions  $\varphi_i^{\text{opt}}$ , measure  $\pi^{\text{opt}}$ , and vector field  $f^{\text{opt}}$  satisfy:

$$\int u^{\text{opt}}(v) \, d\psi(v) = \int u^{\text{opt}}(v) \, d\pi^{\text{opt}}(v)$$

$$f_i^{\text{opt}}(v) \in \partial \varphi_i^{\text{opt}} \left( \frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right)$$

$$\int \varphi_i^{\text{opt}} \left( \frac{\partial u^{\text{opt}}}{\partial v_i}(v) \right) \rho(v) \, dv = \int_0^1 \varphi_i^{\text{opt}}(z^{n-1}) \, dz$$



- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\text{convex monotone } u} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

$$u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)



- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**

$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$  [revenue curve](#)

- **Automated mechanism design:** revenue maximization is an LP, let's feed it to an LP solver; Sandholm (2003)
- **Curse of dimensionality:** If each of  $n$  agents can have  $q$  different values for each of  $m$  items  $\Rightarrow$  the dimension  $\sim (q^n)^m$ 
  - intractable for  $(m = 2, q = 100, n = 2)$  or for  $(m = 2, q = 10, n = 4)$
  - deep neural networks improve the bounds; Dutting et al. (2019)
- **How to avoid:**


$$R_{n,m}(\rho) = \max_{\substack{\text{convex monotone } u \\ u(0) = 0, \partial_{v_i} u(v) \preceq z^{n-1}}} n \cdot \int_{\mathbb{R}_+^m} u(v) d\psi(v)$$

- Pros: dependence on  $n$  is killed; Cai et al.(2012), Alaei et al. (2019)
- Cons: non-linear program
- **Linearization via transport:**
  - $\mu$  on  $[0, 1]$  majorizes  $\nu$  if and only if there is  $\gamma$  on  $[0, 1]^2$  with marginals  $\mu$  on  $y$  and  $\nu$  on  $x$  and such that  $\int y d\gamma(y | x) \geq x$  for  $\gamma$ -almost all  $x$
  - solve for  $(u, \gamma)$
- **Performance:** algorithm handles  $(m = 2, q = 100, n = 10)$



Revenue as a function of the number of bidders  $n$  for two items with i.i.d. values uniform on  $[0, 1]$ . Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for  $n \rightarrow \infty$  (the dashed line).

**Remark:** For  $n = 2$ , selling optimally improves upon selling separately by 5%



Revenue as a function of the number of bidders  $n$  for two items with i.i.d. values uniform on  $[0, 1]$ . Graphs from bottom to top: selling separately (light-green), selling optimally (blue), full surplus extraction (red), limit for  $n \rightarrow \infty$  (the dashed line).

**Remark:** For  $n = 2$ , selling optimally improves upon selling separately by 5%