

# Improvable Equilibria

---

**Kirill Rudov** - UC Berkeley

**Fedor Sandomirskiy** - Princeton

**Leeat Yariv** - Princeton

Columbia Conference in Economic Theory, September 13, 2024

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

**This project:** When is there potential value in correlation?

## Normal-form game

$$\Gamma = \left( N, (A_i)_{i \in N}, (u_i: A \rightarrow \mathbb{R})_{i \in N} \right)$$

- $N = \{1, \dots, n\}$  is finite set of players
- $A_i$  is a finite set of actions of player  $i$
- $A = \prod_{i \in N} A_i$  is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$  is utility of player  $i$

# Correlated Equilibria (CE)

## Definition

A distribution  $\mu \in \Delta(A)$  is a correlated equilibrium if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all  $i \in N$  and all  $a_i, a'_i \in A_i$

**Interpretation:**  $\mu$  generated by a mediator and players best respond by adhering

**Remark:** Nash Equilibria (NE) are CE of the form  $\mu = \mu_1 \times \dots \times \mu_n$

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

## Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

## Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

**Our Question:** When is a Nash equilibrium extreme?



# Improvability of non-extreme equilibria

Maximization of a linear objective—e.g., utilitarian welfare—over a polytope  $P$ :




Two cases:

- If the optimum is unique, it is an extreme point
  - We call objectives with a unique optimum **non-degenerate**
  - Utilitarian welfare is non-degenerate, as we will see
- In knife-edge cases, the whole face of  $P$  can be optimal

# Improvability of non-extreme equilibria

Maximization of a linear objective—e.g., utilitarian welfare—over a polytope  $P$ :



Two cases:

- If the optimum is unique, it is an extreme point
  - We call objectives with a unique optimum **non-degenerate**
  - Utilitarian welfare is non-degenerate, as we will see
- In knife-edge cases, the whole face of  $P$  can be optimal

## Observation

NE is non-extreme  $\iff$  any non-degenerate linear objective can be improved

### Bauer's Maximum Principle

Any non-degenerate linear or (quasi-)convex objective attains its maximum at an extreme point

- $\Rightarrow$  Non-extreme equilibria are improvable **no matter** the objective
- A conservative notion, agnostic to the designer's objective

- Extreme Nash equilibria = vertices of the set of CE
- Non-extreme Nash are improvable for **any** non-degenerate linear or convex objective
- This improvability notion seems very demanding
- Are NE often extreme?

## Conditions for Extremality

---

## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)



## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme (Cripps, 1995)

## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme (Cripps, 1995)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness

## Theorem 1

In a generic  $n$ -player game, a mixed NE is extreme  $\iff \leq 2$  players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme (Cripps, 1995)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness  
 $\Rightarrow$  2-player games not representative

Genericity can be dropped in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

Genericity can be dropped in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize

Genericity can be dropped in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)

Genericity can be dropped in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small payoff perturbations

## Theorem 1'

In any game, a regular mixed NE is extreme  $\iff \leq 2$  players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)
- Hence, Theorem 1'  $\Rightarrow$  Theorem 1


## **Example: 2 Players vs 3 Players**

---



## Example: 2-Player Games

A version of the Game of Chicken by **Aumann (1974)**:



	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9

## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players

Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players  
Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)

## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players  
Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**


## Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	$p$	$1 - p$

- Mixed NE:  $(1/2, 1/2)$  for both players  
Solves linear equation:  $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**
- Indeed, it is the optimum for a non-degenerate objective


weight of (Risky, Risky) & (Safe, Safe)  $\rightarrow \max$

## Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

## Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10


## Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10




## Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	
				$p$	$1 - p$	$1 - p$

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation on  $p \Rightarrow$  irrational weights (Nash, 1950)


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	
				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation on  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates


## Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
Safe				$p$	$1 - p$	

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation on  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates
- The mixed NE is **not extreme**

## Example: 3-Player Games



		Safe		Risky			
		Risky	Safe	Risky	Safe		
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	$p$	$1 - p$
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10		
				$p$	$1 - p$		

- Symmetric Mixed NE:  $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$  for each player
- Non-linear equation on  $p \Rightarrow$  irrational weights (Nash, 1950)
- However, extreme CE solve a linear system  $\Rightarrow$  have rational coordinates
- The mixed NE is **not extreme**

More than 2 players mixing makes a difference...

## General Proof Intuition

---

## How the Actual Proof Goes

**Idea:** When many players randomize, there are too many ways to correlate their actions, one must be beneficial

Focus on a particular example to illustrate



## How the Actual Proof Goes

- Game with  $n$  players, each with 2 actions

## How the Actual Proof Goes

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

## How the Actual Proof Goes

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints

## How the Actual Proof Goes

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq k + 1$

## How the Actual Proof Goes

- Game with  $n$  players, each with 2 actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq k + 1$
- $\Rightarrow$  support of an extreme CE  $\mu$  is bounded by  $2n + 1$

## How the Actual Proof Goes

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing

## How the Actual Proof Goes

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles

## How the Actual Proof Goes

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$



## How the Actual Proof Goes

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- At most  $\log_2(2n + 1)$  out of  $n$  players can randomize

## How the Actual Proof Goes

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  players mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be extreme,

$$2^k \leq 2n + 1$$

- At most  $\log_2(2n + 1)$  out of  $n$  players can randomize

**Conclusion:** NE with too much randomness cannot be extreme

- In fact, 3 mixing agents is already too much [▶ details](#)

## Extreme Points in Payoff Space

---

- The set of CE  $\subset \Delta(A)$  subset of a space of dimension  $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in  $\mathbb{R}^n$

## Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

- The set of CE  $\subset \Delta(A)$  subset of a space of dimension  $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in  $\mathbb{R}^n$

## Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

**Question:** What can we say about payoff-extreme equilibria?

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

- Projection of an extreme point *need not* be an extreme point of a projection



## Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection  $\subset$  projection of extreme points

## Corollary

In a generic game, a Nash equilibrium with  $\geq 3$  players randomizing is **not payoff-extreme**

- Projection of an extreme point *need not* be an extreme point of a projection
- $\Rightarrow$  pure NE and NE with 2 mixers *need not* be payoff-extreme
  - e.g, the mixed NE in the Game of Chicken

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case  $\alpha_1 = \dots = \alpha_n = 1$  corresponds to the **utilitarian welfare**

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case  $\alpha_1 = \dots = \alpha_n = 1$  corresponds to the **utilitarian welfare**
- Non-degeneracy means unique optimum

- NE is not payoff-extreme  $\Rightarrow$  any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case  $\alpha_1 = \dots = \alpha_n = 1$  corresponds to the **utilitarian welfare**
- Non-degeneracy means unique optimum

## Proposition

In a generic game, utilitarian welfare is non-degenerate

## **Applications to Particular Classes of Games**

---

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$



## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

**Other Applications:** games where players want to mismatch actions of others

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are not extreme

**Other Applications:** games where players want to mismatch actions of others

- e.g., network games (with substitutes), congestion games, all-pay auctions, Boston matching mechanism

# Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is *not extreme* in the (smaller!) set of *symmetric CE*

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is *not extreme* in the (smaller!) set of *symmetric CE*

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

## Theorem 2

In any symmetric game with  $n \geq 3$  players, a completely mixed symmetric NE is *not extreme* in the (smaller!) set of *symmetric CE*

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

**Take-away:** caution when focusing on symmetric mixed equilibria in symmetric games



# Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$  robustness to incomplete information about payoffs (Einy et al., 2022)

# Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$  robustness to incomplete information about payoffs (Einy et al., 2022)

## Corollary

If a game has a unique correlated equilibrium  $\nu$ , then  $\nu$  is either:

- A pure Nash equilibrium, or
- A Nash equilibrium where exactly two players randomize

# Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$  robustness to incomplete information about payoffs (Einy et al., 2022)

## Corollary

If a game has a unique correlated equilibrium  $\nu$ , then  $\nu$  is either:

- A pure Nash equilibrium, or
  - A Nash equilibrium where exactly two players randomize
- 
- No genericity assumption needed thanks to the open-set property

## What Extreme CE Look Like

---

► skip

## What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?

# What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?

- For general games, we only know that extreme CE have small support
- For symmetric games and symmetric CE, we can say more

**Observation:**

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable

**Observation:**

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known



## Observation:

- For a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known

## Theorem (de Finetti)

Any infinite exchangeable sequence  $a_1, a_2, a_3 \dots$  is a mixture of i.i.d. distributions

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### **Proposition 2**

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

# Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parameter family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

- A version of Proposition 2 holds: sampling without replacement instead of i.i.d.

# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)



# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design

# Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:


- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Ongoing:


- Incomplete information
- “Correlated implementation” in mechanism design

# Thank you!



# Simulations



(a)  $NE/2u$



(b) Fraction of CE  $> NE^*$



## Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If  $\mu$  is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |S_i| \cdot (|S_i| - 1)$$

## Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium,  $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ :

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

## Regularity of Generic games (**Harsanyi, 1973**)

In a generic game, any Nash equilibrium is regular

# General linear objectives

- Consider a NE  $\nu$
- For simplicity,  $\nu$  has full support
- By Farkas lemma, a linear objective  $L$  can be improved for  $\nu \iff L$  **cannot** be expressed as

$$L(\mu) = C + \sum_{i, a_i, a'_i, a_{-i}} \mu(a) \cdot \lambda_i(a_i, a'_i) \cdot (u_i(a_i, a_{-i}) - u_i(a'_i, a_{-i}))$$

for some  $\lambda_i(a_i, a'_i) \geq 0$ .

- For non-extreme NE  $\nu$ , “bad”  $L$  form a lower-dimensional subspace

# Extreme Symmetric CE with Any Number of Players

Consider  $n$  players with  $m$  actions each

## Proposition

Any extreme symmetric CE can be obtained as follows:

- there are  $M$  urns, each with  $n$  balls labeled by actions

$$1 \leq M \leq m(m-1) + 1$$

- an urn is selected at random according to  $p \in \Delta_M$ , secretly from players
- players draw balls sequentially without replacement
- $i$ 's action = her ball's label, no incentive to deviate

**Remark:** If  $n$  is large, sampling without replacement can be approximated by i.i.d.

## Bayesian games

---



## Bayesian game

$$\mathcal{B} = \left( N, (A_i)_{i \in N}, (T_i)_{i \in N}, \tau \in \Delta(T), (u_i: A \times T_i \rightarrow \mathbb{R})_{i \in N} \right)$$

- Each player  $i \in N$  has a type  $t_i \in T_i$
- Profile of types  $(t_1, \dots, t_n) \in T$  sampled from  $\tau$
- Each player  $i$  observes her realized type
- Utility  $u_i: A \times T_i \rightarrow \mathbb{R}$  depends on the action profile and  $i$ 's type

**Technical assumption:** sets of types  $T_i$  are finite

# Bayesian Correlated Equilibria (BCE)

## Definition

A joint distribution  $\mu \in \Delta(A \times T)$  is a Bayesian correlated equilibrium if

- The marginal on  $T$  coincides with  $\tau$
- For each player  $i$ , type  $t_i$ , recommended action  $a_i$ , and deviation  $a'_i$ ,

$$\sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a_i, t_i, a_{-i}) \geq \sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a'_i, t_i, a_{-i})$$

**Interpretation:** a mediator having access to realized types recommends actions to each player. Two aspects:

1. *Ex-ante coordination:* a source of correlated randomness (as in CE)
2. *Information sharing:* providing  $i$  more info about  $t_{-i}$  than contained in  $t_i$

**Remark:** Bergemann and Morris (2016) allow for a broader class of BCE, where player  $i$  observes a noisy signal about her type

# Induced Complete Information Game

We can associate a complete information normal form game  $\Gamma_{\mathcal{B}}$  with  $\mathcal{B}$ :

- Replace  $A_i$  with set of functions  $\sigma_i : T_i \rightarrow A_i$
- $\Sigma_i$  is the set of all such  $\sigma_i$
- Utility  $v_i : \Sigma \rightarrow \mathbb{R}$  is given by

$$v_i(\sigma) = \sum_{t \in T} \tau(t) \cdot u_i((\sigma_1(t_1), \dots, \sigma_n(t_n)), t_i)$$

## Induced Complete Information Game

$$\Gamma_{\mathcal{B}} = (N, (\Sigma_i)_{i \in N}, (v_i)_{i \in N})$$

**Question:** What is a relation between CE of  $\Gamma_{\mathcal{B}}$  and BCE of  $\mathcal{B}$ ?

# Induced complete information game

## Relationship between equilibria in $\Gamma_{\mathcal{B}}$ and $\mathcal{B}$

CE in  $\Gamma_{\mathcal{B}} \Leftrightarrow$  ex-ante coordination in  $\mathcal{B}$  with no information sharing

- i.e., BCE such that  $a_i$  is independent of  $t_{-i}$  conditionally on  $t_i$

Nash in  $\Gamma_{\mathcal{B}} \Leftrightarrow$  Bayes-Nash in  $\mathcal{B}$

**Observation:** Generic  $\mathcal{B}$  leads to generic  $\Gamma_{\mathcal{B}}$

- $\Rightarrow$  we can apply our theorem to  $\Gamma_{\mathcal{B}}$  to learn about generic  $\mathcal{B}$

## Corollary

For a generic Bayesian game, a Bayes-Nash equilibrium is improvable via ex-ante coordination  $\iff$  at least 3 players randomize

Applies to Bayesian games where players randomize in equilibrium, e.g., costly voting with private types (Feddersen and Pesendorfer, 1997) and contests (Baranski and Goel, 2024)

## References

---

- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Baranski, A. and S. Goel (2024). Contest design with a finite type-space. *to appear*.
- Bergemann, D. and S. Morris (2016). Bayes correlated equilibrium and the comparison of information structures in games. *Theoretical Economics* 11(2), 487–522.
- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Feddersen, T. and W. Pesendorfer (1997). Voting behavior and information aggregation in elections with private information. *Econometrica: Journal of the Econometric Society*, 1029–1058.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings* 12, pp. 131–144. Springer.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.

- Nash, J. F. (1950). Non-cooperative games.
- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Palfrey, T. R. and H. Rosenthal (1983). A strategic calculus of voting. *Public Choice* 41(1), 7–53.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.
- Viossat, Y. (2010). Properties and applications of dual reduction. *Economic theory* 44, 53–68.
- Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.