

Extreme Equilibria:

The Benefits of Correlation

Kirill Rudov - Analysis Group

Fedor Sandomirskiy - Princeton

Leeat Yariv - Princeton

OSU, October 3, 2025

Correlated Equilibria (Aumann, 1974) generalize Nash equilibria to allow correlation

- capture pre-play communication, intermediation, and joint randomization
Bárány (1992), Ben-Porath (1998), Gerardi (2004), Lehrer and Sorin (1997)
- result from natural learning dynamics
Foster and Vohra (1997), Fudenberg and Levine (1999), Hart and Mas-Colell (2000)
- efficiently computable
Papadimitriou and Roughgarden (2008)

Correlated Equilibria (Aumann, 1974) generalize Nash equilibria to allow correlation

- capture pre-play communication, intermediation, and joint randomization
Bárány (1992), Ben-Porath (1998), Gerardi (2004), Lehrer and Sorin (1997)
- result from natural learning dynamics
Foster and Vohra (1997), Fudenberg and Levine (1999), Hart and Mas-Colell (2000)
- efficiently computable
Papadimitriou and Roughgarden (2008)

Broad question: When is there potential value in correlation?

CE \simeq adding a recommendation system on top of the existing interaction

- \implies What interactions can be improved by a recommendation system?

CE \simeq adding a recommendation system on top of the existing interaction

- \implies What interactions can be improved by a recommendation system?

CE \simeq outcomes of arbitrary pre-play communication protocols

- \implies What strategic interactions are susceptible to communication / collusion?

Normal-form game

$$\Gamma = \left(N, (A_i)_{i \in N}, (u_i)_{i \in N} \right)$$

- $N = \{1, \dots, n\}$ is finite set of players
- A_i is a finite set of actions of player i
- $A = \prod_{i \in N} A_i$ is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$ is utility of player i

Definition (Aumann, 1974)

A distribution $\mu \in \Delta(A)$ is a CE if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all $i \in N$ and all $a_i, a'_i \in A_i$

Interpretation: μ generated by a mediator and players best respond by adhering

Remark: Nash Equilibria (NE) are CE of the form $\mu = \mu_1 \times \dots \times \mu_n$

Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is the convex hull of its vertices, aka **extreme points**

Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is the convex hull of its vertices, aka **extreme points**

Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

Formalizing the Question

- The set of correlated equilibria is a convex polytope
- A polytope is the convex hull of its vertices, aka **extreme points**


Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

Our Question: When is a Nash equilibrium extreme?


Improvability of non-extreme equilibria

Maximization of a linear objective over a polytope P :



Improvability of non-extreme equilibria

Maximization of a linear objective over a polytope P :




Bauer's Maximum Principle

Generically, any linear or convex objective attains its unique maximum at an extreme point

- \Rightarrow Non-extreme equilibria are generically improvable

Improvability of non-extreme equilibria

Maximization of a linear objective over a polytope P :




Bauer's Maximum Principle

Generically, any linear or convex objective attains its unique maximum at an extreme point

- \Rightarrow Non-extreme equilibria are generically improvable
- A conservative notion, agnostic to the designer's objective
- Might seem too demanding as it includes contrived objectives

Improvability of non-extreme equilibria

Maximization of a linear objective over a polytope P :



Bauer's Maximum Principle

Generically, any linear or convex objective attains its unique maximum at an extreme point

- \Rightarrow Non-extreme equilibria are generically improvable
- A conservative notion, agnostic to the designer's objective
- Might seem too demanding as it includes contrived objectives

Main Insight: Improvable, or **non-extreme**, NE are prevalent

- **Value of correlation in 2-player games:** Cripps (1995), Evangelista and Raghavan (1996), Canovas, Hansen, and Jaumard (1999), Nau, Canovas, and Hansen (2004), Peeters and Potters (1999), Calvó-Armengol (2006), Ashlagi, Monderer, and Tennenholtz (2008)
- **Extreme-point approach in info & mech. design:** Manelli and Vincent (2007), Kleiner, Moldovanu, and Strack (2021), Arieli, Babichenko, Smorodinsky, and Yamashita (2023), Yang and Zentefis (2024), Nikzad (2022), Kleiner, Moldovanu, Strack, and Whitmeyer (2024), Lahr and Niemeyer (2024)

- **Conditions for extremality:**
in the space of action distributions and payoff space
- **Particular classes of games:**
symmetric, having unique CE

Conditions for Extremality

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness

Theorem

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness
 \Rightarrow 2-player games not representative

Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...

Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)

Genericity Assumption

- A **generic** game is a game from an open everywhere dense set with the complement of 0 Lebesgue measure
- But no given game is generic...
- A version of the theorem applies to **all games** and **regular** NE

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'


In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)
- Hence, Theorem 1' \Rightarrow Theorem 1

Example: 2 Players vs 3 Players

Example: 2-Player Games

A version of the Game of Chicken by **Aumann (1974)**:



	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players

Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players
Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players

Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**


Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players
Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**
- Indeed, it is the optimum for a non-degenerate objective


weight of (Risky, Risky) & (Safe, Safe) $\rightarrow \max$

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
Safe				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
Safe				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
Safe				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	
				p	$1 - p$	$1 - p$

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates
- The mixed NE is **not extreme**

Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates
- The mixed NE is **not extreme**

More than 2 players mixing makes a difference...

General Proof Intuition

High-level idea: When many players randomize, there are too many ways to correlate their actions \implies one must be beneficial

Focus on a particular example to illustrate

- Game with n players, each with 2 actions

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints
- **Winkler (1988)**: if k linear constraints are imposed on the set of all distributions $\Delta(A)$, extreme distributions have support $\leq k + 1$

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints
- **Winkler (1988)**: if k linear constraints are imposed on the set of all distributions $\Delta(A)$, extreme distributions have support $\leq k + 1$
- \Rightarrow support of an extreme CE μ is bounded by $2n + 1$

- Suppose ν is a Nash equilibrium with the k players mixing

General Proof Intuition

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles

General Proof Intuition

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

- The same argument applies to equilibria where players mix over the same number of pure strategies

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

- The same argument applies to equilibria where players mix over the same number of pure strategies
- The main difficulty is handling very asymmetric equilibria [▶ details](#)

Utilitarian and Pareto Improvements

Particular Objectives

- So far, agnostic perspective: improvability for generic objectives
- What about specific goals like **utilitarian welfare** or **Pareto efficiency**?

Particular Objectives

- So far, agnostic perspective: improvability for generic objectives
- What about specific goals like **utilitarian welfare** or **Pareto efficiency**?
- Relevant information is captured in the **payoff space**
⇒ represent equilibria via payoff vectors in \mathbb{R}^n , not distributions in $\Delta(A)$

Particular Objectives

- So far, agnostic perspective: improvability for generic objectives
- What about specific goals like **utilitarian welfare** or **Pareto efficiency**?
- Relevant information is captured in the **payoff space**
⇒ represent equilibria via payoff vectors in \mathbb{R}^n , not distributions in $\Delta(A)$

Proposition

In a generic game, any NE ν with three or more agents randomizing:

- not extreme in the payoff space
- its utilitarian welfare $\sum_i u_i(\nu)$ can be strictly improved

Particular Objectives

- So far, agnostic perspective: improvability for generic objectives
- What about specific goals like **utilitarian welfare** or **Pareto efficiency**?
- Relevant information is captured in the **payoff space**
⇒ represent equilibria via payoff vectors in \mathbb{R}^n , not distributions in $\Delta(A)$

Proposition

In a generic game, any NE ν with three or more agents randomizing:

- not extreme in the payoff space
- its utilitarian welfare $\sum_i u_i(\nu)$ can be strictly improved


Remark: For 2 agents mixing, the NE may or may not be extreme

- Example: the game of chicken

- CE payoffs = projection of CE to a lower-dimensional space


Geometric Intuition

- CE payoffs = projection of CE to a lower-dimensional space



Geometric Intuition


- CE payoffs = projection of CE to a lower-dimensional space



- For a generic projection, extreme points only originate from extreme points

Geometric Intuition


- CE payoffs = projection of CE to a lower-dimensional space



- For a generic projection, extreme points only originate from extreme points
- Projections to the payoff space \simeq generic projection

Geometric Intuition

- CE payoffs = projection of CE to a lower-dimensional space



- For a generic projection, extreme points only originate from extreme points
- Projections to the payoff space \simeq generic projection
- \Rightarrow NE with ≥ 3 mixers cannot lead to extreme payoffs


Proposition

In a generic game, any NE with at least $9 + \log_2(n + 1)$ mixers is Pareto dominated

Proposition

In a generic game, any NE with at least $9 + \log_2(n + 1)$ mixers is Pareto dominated


Intuition: For a generic projection of a polytope to an n -dimensional space, the interior of n -dimensional faces maps to the interior of the image.



Proposition

In a generic game, any NE with at least $9 + \log_2(n + 1)$ mixers is Pareto dominated

Intuition: For a generic projection of a polytope to an n -dimensional space, the interior of n -dimensional faces maps to the interior of the image.



$\geq 9 + \log_2(n + 1)$ agents randomizing \Rightarrow in a generic game, $\text{NE} \in$ a face of dimension at least n of the CE polytope.

What Extreme CE Look Like

► skip

What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

Question: What is the structure of extreme CE?

What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

Question: What is the structure of extreme CE?

- For general games, we only know that extreme CE have small support
- For symmetric games, we can say more

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is not extreme in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

Take-away: symmetric mixed equilibria in symmetric games are inherently suboptimal

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable
- If $n \rightarrow \infty$, the structure of exchangeable distributions is well-known

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable
- If $n \rightarrow \infty$, the structure of exchangeable distributions is well-known

Theorem (de Finetti)

Any infinite exchangeable sequence $a_1, a_2, a_3 \dots$ is a mixture of i.i.d. distributions

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Question: What if we want the exact result, not an approximation?

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Question: What if we want the exact result, not an approximation?

- A version of Proposition 2 holds: sampling without replacement instead of i.i.d.

Example: Symmetric Binary Action Games

- n agents, actions from $\{0, 1\}$
 - Take route A or route B, vote or not, protest or not

Example: Symmetric Binary Action Games

- n agents, actions from $\{0, 1\}$
 - Take route A or route B, vote or not, protest or not
- Normalize payoff from action 0 to 0

Example: Symmetric Binary Action Games

- n agents, actions from $\{0, 1\}$
 - Take route A or route B, vote or not, protest or not
- Normalize payoff from action 0 to 0
- Payoff for player i :

$$u_i(a_i, a_{-i}) = \begin{cases} f\left(\frac{|\{j \in N: a_j=1\}|}{n}\right), & a_i = 1 \\ 0, & a_i = 0 \end{cases}$$

where f is continuous, takes both positive and negative values, and $f(1) < 0$

Example: Symmetric Binary Action Games

- n agents, actions from $\{0, 1\}$
 - Take route A or route B, vote or not, protest or not
- Normalize payoff from action 0 to 0
- Payoff for player i :

$$u_i(a_i, a_{-i}) = \begin{cases} f\left(\frac{|\{j \in N: a_j=1\}|}{n}\right), & a_i = 1 \\ 0, & a_i = 0 \end{cases}$$

where f is continuous, takes both positive and negative values, and $f(1) < 0$

- Focus on large-population behavior

Nash equilibrium characterization


Proposition

All agents' equilibrium payoffs at all Nash equilibria converge to 0 as $n \rightarrow \infty$

Nash equilibrium characterization

Proposition

All agents' equilibrium payoffs at all Nash equilibria converge to 0 as $n \rightarrow \infty$



- In shaded areas, incentive to deviate from 0 to 1
- In blank areas, incentive to deviate from 1 to 0

Finding Optimal Correlated Equilibrium

De Finetti's: A symmetric CE \simeq a mixture of i.i.d. distributions

- Denote X is the (random) fraction of agents choosing 1

Finding Optimal Correlated Equilibrium

De Finetti's: A symmetric CE \simeq a mixture of i.i.d. distributions

- Denote X is the (random) fraction of agents choosing 1
- Per-capita utilitarian welfare is

$$W = \mathbb{E}[Xf(X)]$$

Finding Optimal Correlated Equilibrium

De Finetti's: A symmetric CE \simeq a mixture of i.i.d. distributions

- Denote X is the (random) fraction of agents choosing 1
- Per-capita utilitarian welfare is

$$W = \mathbb{E}[Xf(X)]$$

- An agent prescribed the action 0 should abide $\Rightarrow \mathbb{E}[(1 - X)f(X)] \leq 0$

Finding Optimal Correlated Equilibrium

De Finetti's: A symmetric CE \simeq a mixture of i.i.d. distributions

- Denote X is the (random) fraction of agents choosing 1
- Per-capita utilitarian welfare is

$$W = \mathbb{E}[Xf(X)]$$

- An agent prescribed the action 0 should abide $\Rightarrow \mathbb{E}[(1 - X)f(X)] \leq 0$
- An agent prescribed the action 1 should abide $\Rightarrow \mathbb{E}[Xf(X)] = W \geq 0$
 - satisfied automatically

Finding Optimal Correlated Equilibrium

De Finetti's: A symmetric CE \simeq a mixture of i.i.d. distributions

- Denote X is the (random) fraction of agents choosing 1
- Per-capita utilitarian welfare is

$$W = \mathbb{E}[Xf(X)]$$

- An agent prescribed the action 0 should abide $\Rightarrow \mathbb{E}[(1 - X)f(X)] \leq 0$
- An agent prescribed the action 1 should abide $\Rightarrow \mathbb{E}[Xf(X)] = W \geq 0$
 - satisfied automatically

Optimization Problem for Utilitarian Optimal CE

$\max W = \mathbb{E}[Xf(X)]$ over distributions on $[0, 1]$ subject to $IC = \mathbb{E}[(1 - X)f(X)] \leq 0$

Optimization Problem for Utilitarian Optimal CE

$\max W = \mathbb{E}[Xf(X)]$ over distributions on $[0, 1]$ subject to $IC = \mathbb{E}[(1 - X)f(X)] \leq 0$

Optimization Problem for Utilitarian Optimal CE

$\max W = \mathbb{E}[Xf(X)]$ over distributions on $[0, 1]$ subject to $IC = \mathbb{E}[(1 - X)f(X)] \leq 0$

- Define $\varphi(x) = (\underbrace{xf(x)}_{W(x)}, \underbrace{(1-x)f(x)}_{IC(x)})$ and consider its graph

$$\Phi = \left\{ \varphi(x), x \in [0, 1] \right\} \subset \mathbb{R}^2$$

- Φ describes feasible (W, IC) arising from a point-mass distribution at some x

Optimization Problem for Utilitarian Optimal CE

$\max W = \mathbb{E}[Xf(X)]$ over distributions on $[0, 1]$ subject to $IC = \mathbb{E}[(1 - X)f(X)] \leq 0$

- Define $\varphi(x) = (\underbrace{xf(x)}_{W(x)}, \underbrace{(1-x)f(x)}_{IC(x)})$ and consider its graph

$$\Phi = \left\{ \varphi(x), x \in [0, 1] \right\} \subset \mathbb{R}^2$$

- Φ describes feasible (W, IC) arising from a point-mass distribution at some x
- In general, (W, IC) is feasible if and only if $(W, IC) \in \text{conv}[\Phi]$

Optimization Problem for Utilitarian Optimal CE

$\max W = \mathbb{E}[Xf(X)]$ over distributions on $[0, 1]$ subject to $IC = \mathbb{E}[(1 - X)f(X)] \leq 0$

- Define $\varphi(x) = (\underbrace{xf(x)}_{W(x)}, \underbrace{(1-x)f(x)}_{IC(x)})$ and consider its graph

$$\Phi = \left\{ \varphi(x), x \in [0, 1] \right\} \subset \mathbb{R}^2$$

- Φ describes feasible (W, IC) arising from a point-mass distribution at some x
- In general, (W, IC) is feasible if and only if $(W, IC) \in \text{conv}[\Phi]$

Optimization Problem for Utilitarian Optimal CE 2


$$\max W \quad \text{over} \quad (W, IC) \in \text{conv}[\Phi], \quad IC \leq 0$$

Optimization Problem for Utilitarian Optimal CE 2

$$\max W \quad \text{over} \quad (W, IC) \in \text{conv}[\Phi], \quad IC \leq 0$$

Assume:

- f symmetric around $1/2$
- $f(1/2) > 0$
- $f(0) = f(1) < 0$




Optimization Problem for Utilitarian Optimal CE 2

$$\max W \quad \text{over} \quad (W, IC) \in \text{conv}[\Phi], \quad IC \leq 0$$

Assume:

- f symmetric around $1/2$
- $f(1/2) > 0$
- $f(0) = f(1) < 0$



Optimum:

randomize between $x = 0$ and some $x = x^* > 1/2$ with weights making IC bind

Games with Unique Correlated Equilibrium

► skip

Games with Unique Correlated Equilibrium

- Unique CE \implies CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

Games with Unique Correlated Equilibrium

- Unique CE \implies CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

Corollary

If a game has a unique CE, then it is either:

- a pure Nash, or
- a Nash where exactly two players randomize

Games with Unique Correlated Equilibrium

- Unique CE \implies CE=NE
- Such NE is robust to communication/collusion, incomplete information about payoffs, and can be computed without fixed points (Einy et al., 2022)
- **Examples:** games with dominant strategies, some congestion games, Cournot competition

Corollary

If a game has a unique CE, then it is either:

- a pure Nash, or
 - a Nash where exactly two players randomize
-
- No genericity assumption since games with a unique CE form an open set (Viossat, 2010)

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a general tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Main takeaway: inherent suboptimality of equilibria with a lot of mixing

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a general tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Main takeaway: inherent suboptimality of equilibria with a lot of mixing

Thank you!

Coarse Correlated Equilibria

► skip

Coarse Correlated Equilibria

Consider situations where agents commit **ex-ante** to a correlating device, before receiving recommendations. For example,

- firms entering binding collusive agreements (McAfee and McMillan, 1992)
- users opting in to algorithmic recommendation systems

Coarse Correlated Equilibria

Consider situations where agents commit **ex-ante** to a correlating device, before receiving recommendations. For example,

- firms entering binding collusive agreements (McAfee and McMillan, 1992)
- users opting in to algorithmic recommendation systems

Definition (Hannan, 1957; Moulin and Vial, 1978)

A distribution $\mu \in \Delta(A)$ is a coarse correlated equilibrium (CCE) if, for all $i \in N$,

$$\sum_{a \in A} \mu(a) u_i(a) \geq \max_{a'_i \in A_i} \sum_{a \in A} u_i(a'_i, a_{-i}) \mu(a),$$

i.e., the expected payoff from following the correlated strategy is at least as high as the best deterministic deviation

- $\text{CCE} \supseteq \text{CE} \supseteq \text{NE}$

Proposition

A NE an extreme point of the set of CCE \Leftrightarrow

- it is pure
- or 2 players randomize over 2 actions each and this subgame is strategically equivalent to “matching pennies”

Proposition

A NE an extreme point of the set of CCE \Leftrightarrow

- it is pure
 - or 2 players randomize over 2 actions each and this subgame is strategically equivalent to “matching pennies”
-
- No genericity assumption
 - The tension between randomness and optimality is even stronger for CCE than for CE
 - Consistent with the prevalence of examples where CCE improves over NE, even in two-player games (Moulin and Vial, 1978; Gérard-Varet and Moulin, 1978; Moulin, Ray, and Gupta, 2014; Dokka, Moulin, Ray, and SenGupta, 2023)

Key Lemmas

Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If μ is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1)$$

Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium, $\nu = (\nu_1, \nu_2, \dots, \nu_n)$:


$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

Regularity of Generic games (**Harsanyi, 1973**)


In a generic game, any Nash equilibrium is regular

Simulations



Simulations



(a) $NE/2u$



(b) Fraction of CE $>$ NE*



Extreme Symmetric CE with Any Number of Players

Consider n players with m actions each

Proposition

Any extreme symmetric CE can be obtained as follows:

- there are M urns, each with n balls labeled by actions

$$1 \leq M \leq m(m-1) + 1$$

- an urn is selected at random according to $p \in \Delta_M$, secretly from players
- players draw balls sequentially without replacement
- i 's action = her ball's label, no incentive to deviate

Remark: If n is large, sampling without replacement can be approximated by i.i.d.

Bayesian Correlated Equilibria

► skip

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states Θ
- Private information: finite sets of types T_i
- Prior $\pi \in \Delta(\Theta \times T)$

Bayesian Correlated Equilibria

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states Θ
- Private information: finite sets of types T_i
- Prior $\pi \in \Delta(\Theta \times T)$

Definition

A distribution $\psi \in \Delta(A \times \Theta \times T)$ is a BCE if

- its marginal on $\Theta \times T$ coincides with π
- no agent can gain by deviating from a recommended action a_i to another action a'_i , given her private type t_i

$$\sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a, \theta, t) \geq \sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a'_i, a_{-i}, \theta, t)$$

Bayesian Correlated Equilibria

General games with incomplete information (Bergemann and Morris, 2019):

- Common payoff uncertainty: a finite set of states Θ
- Private information: finite sets of types T_i
- Prior $\pi \in \Delta(\Theta \times T)$

Definition

A distribution $\psi \in \Delta(A \times \Theta \times T)$ is a BCE if

- its marginal on $\Theta \times T$ coincides with π
- no agent can gain by deviating from a recommended action a_i to another action a'_i , given her private type t_i

$$\sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a, \theta, t) \geq \sum_{a_{-i} \in A_{-i}, \theta \in \Theta, t_{-i} \in T_{-i}} \psi(a, \theta, t) u_i(a'_i, a_{-i}, \theta, t)$$

A Bayesian Nash equilibrium (BNE) is a BCE where a_i is independent of (θ, a_{-i}, t_{-i}) conditional on t_i for each agent i

Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ($|\Theta| \geq 2$), or
- non-trivial private information ($|T_i| \geq 2$ for at least 3 agents),

a BNE is an extreme point of BCE \Leftrightarrow it is pure

Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ($|\Theta| \geq 2$), or
- non-trivial private information ($|T_i| \geq 2$ for at least 3 agents),

a BNE is an extreme point of BCE \Leftrightarrow it is pure

- Even minimal uncertainty—e.g., a single binary state—can be sufficient to render mixed BNE non-extreme
 - Contrast with complete information games, where two agents can mix without losing extremality

Theorem

For a generic game with either:

- non-trivial common payoff uncertainty ($|\Theta| \geq 2$), or
- non-trivial private information ($|T_i| \geq 2$ for at least 3 agents),

a BNE is an extreme point of BCE \Leftrightarrow it is pure

- Even minimal uncertainty—e.g., a single binary state—can be sufficient to render mixed BNE non-extreme
 - Contrast with complete information games, where two agents can mix without losing extremality

Intuition: Randomness in the state and private types provides more degrees of freedom for correlation to be beneficial

References

- Arieli, I., Y. Babichenko, R. Smorodinsky, and T. Yamashita (2023). Optimal persuasion via bi-pooling. *Theoretical Economics* 18(1), 15–36.
- Ashlagi, I., D. Monderer, and M. Tennenholtz (2008). On the value of correlation. *Journal of Artificial Intelligence Research* 33, 575–613.
- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Bárány, I. (1992). Fair distribution protocols or how the players replace fortune. *Mathematics of Operations Research* 17(2), 327–340.
- Ben-Porath, E. (1998). Correlation without mediation: Expanding the set of equilibrium outcomes by “cheap” pre-play procedures. *Journal of Economic Theory* 80(1), 108–122.
- Bergemann, D. and S. Morris (2019). Information design: A unified perspective. *Journal of Economic Literature* 57(1), 44–95.
- Calvo-Armengol, A. (2006). The set of correlated equilibria of 2x2 games. *mimeo*.
- Canovas, S. G., P. Hansen, and B. Jaumard (1999). Nash equilibria from the correlated equilibria viewpoint. *International Game Theory Review* 1(01), 33–44.
- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Dokka, T., H. Moulin, I. Ray, and S. SenGupta (2023). Equilibrium design in an n-player quadratic game. *Review of economic design* 27(2), 419–438.

- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Evangelista, F. S. and T. Raghavan (1996). A note on correlated equilibrium. *International Journal of Game Theory* 25, 35–41.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings* 12, pp. 131–144. Springer.
- Foster, D. P. and R. V. Vohra (1997). Calibrated learning and correlated equilibrium. *Games and Economic Behavior* 21(1-2), 40–55.
- Fudenberg, D. and D. K. Levine (1999). Conditional universal consistency. *Games and Economic Behavior* 29(1-2), 104–130.
- Gérard-Varet, L.-A. and H. Moulin (1978). Correlation and duopoly. *Journal of economic theory* 19(1), 123–149.
- Gerardi, D. (2004). Unmediated communication in games with complete and incomplete information. *Journal of Economic Theory* 114(1), 104–131.
- Hannan, J. (1957). Approximation to bayes risk in repeated play. *Contributions to the Theory of Games* 3(2), 97–139.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.

- Hart, S. and A. Mas-Colell (2000). A simple adaptive procedure leading to correlated equilibrium. *Econometrica* 68(5), 1127–1150.
- Kleiner, A., B. Moldovanu, and P. Strack (2021). Extreme points and majorization: Economic applications. *Econometrica* 89(4), 1557–1593.
- Kleiner, A., B. Moldovanu, P. Strack, and M. Whitmeyer (2024). The extreme points of fusions. *arXiv preprint arXiv:2409.10779*.
- Lahr, P. and A. Niemeyer (2024). Extreme points in multi-dimensional screening. *arXiv preprint arXiv:2412.00649*.
- Lehrer, E. and S. Sorin (1997). One-shot public mediated talk. *Games and Economic Behavior* 20(2), 131–148.
- Manelli, A. M. and D. R. Vincent (2007). Multidimensional mechanism design: Revenue maximization and the multiple-good monopoly. *Journal of Economic theory* 137(1), 153–185.
- McAfee, R. P. and J. McMillan (1992). Bidding rings. *The American Economic Review*, 579–599.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.
- Moulin, H., I. Ray, and S. S. Gupta (2014). Coarse correlated equilibria in an abatement game. Technical report, Cardiff Economics Working Papers.

- Moulin, H. and J. P. Vial (1978). Strategically zero-sum games: the class of games whose completely mixed equilibria cannot be improved upon. *International Journal of Game Theory* 7, 201–221.
- Nash, J. F. (1950). Non-cooperative games.
- Nau, R., S. G. Canovas, and P. Hansen (2004). On the geometry of nash equilibria and correlated equilibria. *International Journal of Game Theory* 32, 443–453.
- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Nikzad, A. (2022). Constrained majorization: Applications in mechanism design. In *Proceedings of the 23rd ACM Conference on Economics and Computation*, pp. 330–331.
- Papadimitriou, C. H. and T. Roughgarden (2008). Computing correlated equilibria in multi-player games. *Journal of the ACM (JACM)* 55(3), 1–29.
- Peeters, R. and J. Potters (1999). On the structure of the set of correlated equilibria in two-by-two bimatrix games.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.
- Viossat, Y. (2010). Properties and applications of dual reduction. *Economic theory* 44, 53–68.
- Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.

Yang, K. H. and A. K. Zentefis (2024). Monotone function intervals: Theory and applications. *American Economic Review* 114(8), 2239–2270.