

Improvable Equilibria

Kirill Rudov - UC Berkeley

Fedor Sandomirskiy - Princeton

Leeat Yariv - Princeton

Penn State, September 27, 2024

Communication or intermediation

- precede many interactions: voting, matching, product adoption, etc.
- a possible channel for collusion by auction bidders, market competitors, and the like

Broad question: What strategic interactions are susceptible to communication influences or collusion?

Correlated equilibria (Aumann, 1974) generalize Nash equilibria to allow correlation

- Can be implemented via communication, as well as mediation or joint randomization
- Capture outcomes of arbitrary communication protocols without explicitly modeling the communication phase

Correlated equilibria (Aumann, 1974) generalize Nash equilibria to allow correlation

- Can be implemented via communication, as well as mediation or joint randomization
- Capture outcomes of arbitrary communication protocols without explicitly modeling the communication phase

This project: When is there potential value in correlation?

Normal-form game

$$\Gamma = \left(N, (A_i)_{i \in N}, (u_i: A \rightarrow \mathbb{R})_{i \in N} \right)$$

- $N = \{1, \dots, n\}$ is finite set of players
- A_i is a finite set of actions of player i
- $A = \prod_{i \in N} A_i$ is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$ is utility of player i

Correlated Equilibria (CE)

Definition

A distribution $\mu \in \Delta(A)$ is a correlated equilibrium if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all $i \in N$ and all $a_i, a'_i \in A_i$

Interpretation: μ generated by a mediator and players best respond by adhering

Remark: Nash Equilibria (NE) are CE of the form $\mu = \mu_1 \times \dots \times \mu_n$

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**

Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

- The set of correlated equilibria is a convex polytope
- A polytope is a convex hull of its vertices, aka **extreme points**


Definition

A Nash equilibrium is **extreme** if it is an extreme point of the set of CE

Our Question: When is a Nash equilibrium extreme?

Improvability of non-extreme equilibria

Maximization of a linear objective—e.g., utilitarian welfare—over a polytope P :




Two cases:

- If the optimum is unique, it is an extreme point
 - We call objectives with a unique optimum **non-degenerate**
 - Utilitarian welfare is non-degenerate, as we will see
- In knife-edge cases, the whole face of P can be optimal

Improvability of non-extreme equilibria

Maximization of a linear objective—e.g., utilitarian welfare—over a polytope P :



Two cases:


- If the optimum is unique, it is an extreme point
 - We call objectives with a unique optimum **non-degenerate**
 - Utilitarian welfare is non-degenerate, as we will see
- In knife-edge cases, the whole face of P can be optimal

Observation

NE is non-extreme \iff any non-degenerate linear objective can be improved

Improvability of non-extreme equilibria

Maximization of a linear objective—e.g., utilitarian welfare—over a polytope P :



Two cases:

- If the optimum is unique, it is an extreme point
 - We call objectives with a unique optimum **non-degenerate**
 - Utilitarian welfare is non-degenerate, as we will see
- In knife-edge cases, the whole face of P can be optimal

Observation

NE is non-extreme \iff any non-degenerate linear objective can be improved

Remark: linear in probabilities, not in actions \Rightarrow a broad class of objectives

Bauer's Maximum Principle

Any non-degenerate linear or (quasi-)convex objective attains its maximum at an extreme point

- \Rightarrow Non-extreme equilibria are improvable **no matter** the objective

Bauer's Maximum Principle

Any non-degenerate linear or (quasi-)convex objective attains its maximum at an extreme point

- \Rightarrow Non-extreme equilibria are improvable **no matter** the objective
- A conservative notion, agnostic to the designer's objective
- Usually, assess outcomes for a **given** objective

Improvability of non-extreme equilibria 2

Bauer's Maximum Principle

Any non-degenerate linear or (quasi-)convex objective attains its maximum at an extreme point

- \Rightarrow Non-extreme equilibria are improvable **no matter** the objective
- A conservative notion, agnostic to the designer's objective
- Usually, assess outcomes for a **given** objective

Main Insight

Despite restrictiveness of improvability notion, **many** equilibria are improvable, i.e., **non-extreme**

- **Value of correlation in 2-player games:** Cripps (1995), Evangelista and Raghavan (1996), Canovas et al. (1999), Nau et al. (2004), Peeters and Potters (1999), Calvó-Armengol (2006), Ashlagi et al. (2008)
- **Communication \Leftrightarrow correlation:** Forges (2020), Bárány (1992), Ben-Porath (1998), Gerardi (2004), Lehrer and Sorin (1997)
- **Communication & collusion in specific contexts:**
 - Bargaining: Crawford (1990), Agranov and Tergiman (2014), Baranski and Kagel (2015)
 - Auctions: McAfee and McMillan (1992), Lopomo et al. (2011), Feldman et al. (2016), Agranov and Yariv (2018), Pavlov (2023)
 - Voting: Gerardi and Yariv (2007), Goeree and Yariv (2011)
 - Matching: Beyhaghi and Tardos (2018), Echenique et al. (2022)
- **Extreme-point approach in info & mech. design:** Kleiner et al. (2021), Arieli et al. (2023), Yang and Zentefis (2024), Kleiner et al. (2024)

- **Part 1**

- Conditions for extremality/improvability
- Translation to payoffs
- Applications

- **Part 2**

- Proof idea
- Simple description of extreme CE

Conditions for Extremality

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness

Theorem 1

In a generic n -player game, a mixed NE is extreme $\iff \leq 2$ players randomize

Complete detail-free characterization of extreme Nash equilibria

- Pure equilibria are extreme (trivial)
- Equilibria with exactly 2 randomizing players are extreme
(Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)
- If 3 or more players randomize, *any* non-degenerate objective can be improved, either by introducing correlation, or by reducing randomness
 \Rightarrow 2-player games not representative

Genericity can be dropped in any game, by considering **regular** NE only

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Genericity can be dropped in any game, by considering **regular** NE only

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

Genericity can be dropped in any game, by considering **regular** NE only

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)

Genericity can be dropped in any game, by considering **regular** NE only

Definition (informal): a NE is regular if it is stable under small payoff perturbations

Theorem 1'


In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- In a generic game, any NE is regular (Harsanyi, 1973)
- Hence, Theorem 1' \Rightarrow Theorem 1

Example: 2 Players vs 3 Players

Example: 2-Player Games

A version of the Game of Chicken by **Aumann (1974)**:



	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players

Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players

Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from $(6,6)$

Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players

Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$

- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**


Example: 2-Player Games

	Risky	Safe
Risky	6, 6	10, 7
Safe	7, 10	9, 9
	p	$1 - p$

- Mixed NE: $(1/2, 1/2)$ for both players
Solves linear equation: $6p + 10(1 - p) = 7p + 9(1 - p) \implies p = 1/2$
- **Aumann (1974)**: CE can increase utilitarian welfare by shifting weight from (6,6)
- However, the mixed NE is an **extreme point**
- Indeed, it is the optimum for a non-degenerate objective


weight of (Risky, Risky) & (Safe, Safe) $\rightarrow \max$

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games




		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10

Example: 3-Player Games



		Safe		Risky	
		Risky	Safe	Risky	Safe
Risky	Safe	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6
		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates


Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
Risky		6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
Safe		7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates
- The mixed NE is **not extreme**

Example: 3-Player Games



		Safe		Risky		
		Risky	Safe	Risky	Safe	
	Risky	6, 6, 5	10, 7, 7	0, 0, 0	6, 5, 6	p
	Safe	7, 10, 7	9, 9, 9	5, 6, 6	7, 7, 10	$1 - p$
				p	$1 - p$	

- Symmetric Mixed NE: $(\sqrt{3/2} - 1, 2 - \sqrt{3/2})$ for each player
- Non-linear equation in $p \Rightarrow$ irrational weights (Nash, 1950)
- However, extreme CE solve a linear system \Rightarrow have rational coordinates
- The mixed NE is **not extreme**

More than 2 players mixing makes a difference...

Extreme Points in Payoff Space

- The set of CE $\subset \Delta(A)$ subset of a space of dimension $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in \mathbb{R}^n

Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

- The set of CE $\subset \Delta(A)$ subset of a space of dimension $|A_1| \cdot \dots \cdot |A_n|$
- Equilibria are often represented via payoffs in \mathbb{R}^n

Definition

A Nash equilibrium is **payoff-extreme** if its payoff vector is an extreme point of the set of CE payoffs

Question: What can we say about payoff-extreme equilibria?

Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection \subset projection of extreme points

Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection \subset projection of extreme points

Corollary

In a generic game, a Nash equilibrium with ≥ 3 players randomizing is **not payoff-extreme**

Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection \subset projection of extreme points

Corollary

In a generic game, a Nash equilibrium with ≥ 3 players randomizing is **not payoff-extreme**

- Projection of an extreme point **need not** be an extreme point of a projection

Observations:

- CE payoffs = projection of CE to a lower-dimensional space
- Extreme points of a projection \subset projection of extreme points

Corollary

In a generic game, a Nash equilibrium with ≥ 3 players randomizing is **not payoff-extreme**

- Projection of an extreme point **need not** be an extreme point of a projection
- \Rightarrow pure NE and NE with 2 mixers **need not** be payoff-extreme
 - e.g, the mixed NE in the Game of Chicken

- NE is not payoff-extreme \Rightarrow any non-degenerate linear objective in the space of payoffs can be improved

- NE is not payoff-extreme \Rightarrow any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- NE is not payoff-extreme \Rightarrow any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case $\alpha_1 = \dots = \alpha_n = 1$ corresponds to the **utilitarian welfare**

- NE is not payoff-extreme \Rightarrow any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case $\alpha_1 = \dots = \alpha_n = 1$ corresponds to the **utilitarian welfare**
- Non-degeneracy means unique optimum

- NE is not payoff-extreme \Rightarrow any non-degenerate linear objective in the space of payoffs can be improved
- Linear objective in payoffs = weighted welfare

$$W(\mu) = \sum_{i \in N} \alpha_i \sum_{s \in S} u_i(s) \mu(s) \rightarrow \max$$

- The case $\alpha_1 = \dots = \alpha_n = 1$ corresponds to the **utilitarian welfare**
- Non-degeneracy means unique optimum

Proposition

In a generic game, utilitarian welfare is non-degenerate

Applications to Particular Classes of Games

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0 otherwise
- Voters in R get utility of 1 if r -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0 otherwise
- Voters in R get utility of 1 if r -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0 otherwise
- Voters in R get utility of 1 if r -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are not extreme

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0 otherwise
- Voters in R get utility of 1 if r -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are not extreme

Other Applications: games where players want to mismatch actions of others

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0 otherwise
- Voters in R get utility of 1 if r -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are not extreme

Other Applications: games where players want to mismatch actions of others

- e.g., network games (with substitutes), congestion games, all-pay auctions, Boston matching mechanism

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is **not extreme** in the (smaller!) set of **symmetric CE**

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is **not extreme** in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria extreme?

Theorem 2

In any symmetric game with $n \geq 3$ players, a completely mixed symmetric NE is **not extreme** in the (smaller!) set of **symmetric CE**

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

Take-away: caution when focusing on symmetric mixed equilibria in symmetric games

Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$ robustness to incomplete information about payoffs (Einy et al., 2022)

Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$ robustness to incomplete information about payoffs (Einy et al., 2022)

Corollary

If a game has a unique correlated equilibrium ν , then ν is either:

- A pure Nash equilibrium, or
- A Nash equilibrium where exactly two players randomize

Games with Unique Correlated Equilibrium

- Games with a unique CE form an open set (Viossat, 2010)
- $NE=CE \Rightarrow$ robustness to incomplete information about payoffs (Einy et al., 2022)

Corollary

If a game has a unique correlated equilibrium ν , then ν is either:

- A pure Nash equilibrium, or
 - A Nash equilibrium where exactly two players randomize
-
- No genericity assumption needed thanks to the open-set property

PART II

How to Prove Theorem 1

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- We've seen intuition based on the possibility of having irrational NE for $n \geq 3$ mixers

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- We've seen intuition based on the possibility of having irrational NE for $n \geq 3$ mixers
- This is not how the actual proof goes since not all NE with $n = 3$ mixers are irrational

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- We've seen intuition based on the possibility of having irrational NE for $n \geq 3$ mixers
- This is not how the actual proof goes since not all NE with $n = 3$ mixers are irrational

Idea: When many players randomize, there are too many ways to correlate their actions, one must be beneficial

Theorem 1'

In any game, a regular mixed NE is extreme $\iff \leq 2$ players randomize

- We've seen intuition based on the possibility of having irrational NE for $n \geq 3$ mixers
- This is not how the actual proof goes since not all NE with $n = 3$ mixers are irrational

Idea: When many players randomize, there are too many ways to correlate their actions, one must be beneficial

Focus on a particular example to illustrate

How the Proof Goes: Example

- Game with n players, each with 2 actions

How the Proof Goes: Example

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

How the Proof Goes: Example

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints

How the Proof Goes: Example

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints
- **Winkler (1988)**: if k linear constraints are imposed on the set of all distributions $\Delta(A)$, extreme distributions have support $\leq k + 1$

How the Proof Goes: Example

- Game with n players, each with 2 actions
- If μ is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints
- **Winkler (1988)**: if k linear constraints are imposed on the set of all distributions $\Delta(A)$, extreme distributions have support $\leq k + 1$
- \Rightarrow support of an extreme CE μ is bounded by $2n + 1$

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

- The same argument applies to equilibria, where players mix over the **same number of pure strategies**

How the Proof Goes: Example

- Suppose ν is a Nash equilibrium with the k players mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be extreme,

$$2^k \leq 2n + 1$$

- We can replace $2n + 1$ with $2k + 1$ by eliminating non-randomizing agents. Thus

$$2^k \leq 2k + 1$$

Conclusion: NE with $k \geq 3$ mixing agents cannot be extreme

- The same argument applies to equilibria, where players mix over the **same number of pure strategies**
- The main difficulty is handling very asymmetric equilibria

Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If μ is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1)$$

Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium, $\nu = (\nu_1, \nu_2, \dots, \nu_n)$:

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

Key Lemmas for the General Proof

Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If μ is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1)$$

Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium, $\nu = (\nu_1, \nu_2, \dots, \nu_n)$:

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

Let's combine these two observations

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant
- \Rightarrow w.l.o.g., ν is fully mixed and all $|A_i| \geq 2$

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant
- \Rightarrow w.l.o.g., ν is fully mixed and all $|A_i| \geq 2$

By the lemmas from the previous slide:

$$\prod_{i=1}^n |A_i| \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1) \quad \Leftarrow \text{the bound on the support of extreme CE}$$

$$|A_i| - 1 \leq \sum_{j \neq i} (|A_j| - 1), \quad \forall i \quad \Leftarrow \text{McKelvey and McLennan (1997)}$$

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant
- \Rightarrow w.l.o.g., ν is fully mixed and all $|A_i| \geq 2$

By the lemmas from the previous slide:

$$\prod_{i=1}^n |A_i| \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1) \quad \Leftarrow \text{the bound on the support of extreme CE}$$

$$|A_i| - 1 \leq \sum_{j \neq i} (|A_j| - 1), \quad \forall i \quad \Leftarrow \text{McKelvey and McLennan (1997)}$$

Proposition

These inequalities can only hold for some integral $|A_i| \geq 2, i = 1 \dots, n$, if $n \leq 2$

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant
- \Rightarrow w.l.o.g., ν is fully mixed and all $|A_i| \geq 2$

By the lemmas from the previous slide:

$$\prod_{i=1}^n |A_i| \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1) \quad \Leftarrow \text{the bound on the support of extreme CE}$$

$$|A_i| - 1 \leq \sum_{j \neq i} (|A_j| - 1), \quad \forall i \quad \Leftarrow \text{McKelvey and McLennan (1997)}$$

Proposition

These inequalities can only hold for some integral $|A_i| \geq 2, i = 1 \dots, n$, if $n \leq 2$

- $\Rightarrow \nu$ with $n \geq 3$ mixers cannot be extreme

How the Proof Goes

Consider a game $\Gamma = (A, u)$ and a non-pure **extreme** regular Nash equilibrium ν

- Since ν is regular, incentive constraints outside of $\text{supp}(\nu)$ are inactive
- \Rightarrow pure strategies outside $\text{supp}(\nu)$ and non-mixing players are irrelevant
- \Rightarrow w.l.o.g., ν is fully mixed and all $|A_i| \geq 2$

By the lemmas from the previous slide:

$$\prod_{i=1}^n |A_i| \leq 1 + \sum_{i \in N} |A_i| \cdot (|A_i| - 1) \quad \Leftarrow \text{the bound on the support of extreme CE}$$

$$|A_i| - 1 \leq \sum_{j \neq i} (|A_j| - 1), \quad \forall i \quad \Leftarrow \text{McKelvey and McLennan (1997)}$$

Proposition

These inequalities can only hold for some integral $|A_i| \geq 2, i = 1 \dots, n$, if $n \leq 2$

- $\Rightarrow \nu$ with $n \geq 3$ mixers cannot be extreme
- The proposition is proved via majorization & Schur convexity

What Extreme CE Look Like

What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

Question: What is the structure of extreme CE?

What Extreme CE Look Like

For a non-extreme NE, any non-degenerate objective can be strictly improved by switching to an extreme CE

Question: What is the structure of extreme CE?

- For general games, we only know that extreme CE have small support
- For symmetric games and symmetric CE, we can say more

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable
- If $n \rightarrow \infty$, the structure of exchangeable distributions is well-known

Observation:

- For a symmetric CE, the random variables a_1, \dots, a_n are exchangeable
- If $n \rightarrow \infty$, the structure of exchangeable distributions is well-known

Theorem (de Finetti)

Any infinite exchangeable sequence $a_1, a_2, a_3 \dots$ is a mixture of i.i.d. distributions

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Question: What if we want the exact result, not an approximation?

Extreme Symmetric CE with Many Players

- Consider a symmetric game with m actions per player
- Assume the number of players n is large

Proposition 2

Any extreme symmetric CE can be approximated by a mixture of $m(m - 1) + 1$ i.i.d. distributions

- For $m = 2$, a mixture of 3 i.i.d. distributions \Rightarrow 5-parameter family of extreme CE
- A radical dimension reduction

Question: What if we want the exact result, not an approximation?

- A version of Proposition 2 holds: sampling without replacement instead of i.i.d.

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:

- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design

Conclusions

Several papers effectively show extremality of NE in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995; Evangelista and Raghavan, 1996; Canovas et al., 1999)

Our paper:


- a tension between equilibrium randomness and extremality
- detail-free criterion for extremality in various settings

Ongoing:


- Incomplete information
- “Correlated implementation” in mechanism design

Thank you!



Simulations



(a) $NE/2u$



(b) Fraction of CE $>$ NE*



General linear objectives

- Consider a NE ν
- For simplicity, ν has full support
- By Farkas lemma, a linear objective L can be improved for $\nu \iff L$ **cannot** be expressed as

$$L(\mu) = C + \sum_{i, a_i, a'_i, a_{-i}} \mu(a) \cdot \lambda_i(a_i, a'_i) \cdot (u_i(a_i, a_{-i}) - u_i(a'_i, a_{-i}))$$

for some $\lambda_i(a_i, a'_i) \geq 0$.

- For non-extreme NE ν , “bad” L form a lower-dimensional subspace

Extreme Symmetric CE with Any Number of Players

Consider n players with m actions each

Proposition

Any extreme symmetric CE can be obtained as follows:

- there are M urns, each with n balls labeled by actions

$$1 \leq M \leq m(m-1) + 1$$

- an urn is selected at random according to $p \in \Delta_M$, secretly from players
- players draw balls sequentially without replacement
- i 's action = her ball's label, no incentive to deviate

Remark: If n is large, sampling without replacement can be approximated by i.i.d.

Bayesian games

Bayesian game

$$\mathcal{B} = \left(N, (A_i)_{i \in N}, (T_i)_{i \in N}, \tau \in \Delta(T), (u_i : A \times T_i \rightarrow \mathbb{R})_{i \in N} \right)$$

- Each player $i \in N$ has a type $t_i \in T_i$
- Profile of types $(t_1, \dots, t_n) \in T$ sampled from τ
- Each player i observes her realized type
- Utility $u_i : A \times T_i \rightarrow \mathbb{R}$ depends on the action profile and i 's type

Technical assumption: sets of types T_i are finite

Bayesian Correlated Equilibria (BCE)

Definition

A joint distribution $\mu \in \Delta(A \times T)$ is a Bayesian correlated equilibrium if

- The marginal on T coincides with τ
- For each player i , type t_i , recommended action a_i , and deviation a'_i ,

$$\sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a_i, t_i, a_{-i}) \geq \sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a'_i, t_i, a_{-i})$$

Interpretation: a mediator having access to realized types recommends actions to each player. Two aspects:

1. **Ex-ante coordination:** a source of correlated randomness (as in CE)
2. **Information sharing:** providing i more info about t_{-i} than contained in t_i

Remark: Bergemann and Morris (2016) allow for a broader class of BCE, where player i observes a noisy signal about her type

Induced Complete Information Game

We can associate a complete information normal form game $\Gamma_{\mathcal{B}}$ with \mathcal{B} :

- Replace A_i with set of functions $\sigma_i : T_i \rightarrow A_i$
- Σ_i is the set of all such σ_i
- Utility $v_i : \Sigma \rightarrow \mathbb{R}$ is given by

$$v_i(\sigma) = \sum_{t \in T} \tau(t) \cdot u_i((\sigma_1(t_1), \dots, \sigma_n(t_n)), t_i)$$

Induced Complete Information Game

$$\Gamma_{\mathcal{B}} = (N, (\Sigma_i)_{i \in N}, (v_i)_{i \in N})$$

Question: What is a relation between CE of $\Gamma_{\mathcal{B}}$ and BCE of \mathcal{B} ?

Induced complete information game

Relationship between equilibria in $\Gamma_{\mathcal{B}}$ and \mathcal{B}

CE in $\Gamma_{\mathcal{B}} \Leftrightarrow$ ex-ante coordination in \mathcal{B} with no information sharing

- i.e., BCE such that a_i is independent of t_{-i} conditionally on t_i

Nash in $\Gamma_{\mathcal{B}} \Leftrightarrow$ Bayes-Nash in \mathcal{B}

Observation: Generic \mathcal{B} leads to generic $\Gamma_{\mathcal{B}}$

- \Rightarrow we can apply our theorem to $\Gamma_{\mathcal{B}}$ to learn about generic \mathcal{B}

Corollary

For a generic Bayesian game, a Bayes-Nash equilibrium is improvable via ex-ante coordination \iff at least 3 players randomize

Applies to Bayesian games where players randomize in equilibrium, e.g., costly voting with private types (Feddersen and Pesendorfer, 1997) and contests (Baranski and Goel, 2024)

References

- Agranov, M. and C. Tergiman (2014). Communication in multilateral bargaining. *Journal of Public Economics* 118, 75–85.
- Agranov, M. and L. Yariv (2018). Collusion through communication in auctions. *Games and Economic Behavior* 107, 93–108.
- Arieli, I., Y. Babichenko, R. Smorodinsky, and T. Yamashita (2023). Optimal persuasion via bi-pooling. *Theoretical Economics* 18(1), 15–36.
- Ashlagi, I., D. Monderer, and M. Tennenholtz (2008). On the value of correlation. *Journal of Artificial Intelligence Research* 33, 575–613.
- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Baranski, A. and S. Goel (2024). Contest design with a finite type-space. *to appear*.
- Baranski, A. and J. H. Kagel (2015). Communication in legislative bargaining. *Journal of the Economic science Association* 1, 59–71.
- Bárány, I. (1992). Fair distribution protocols or how the players replace fortune. *Mathematics of Operations Research* 17(2), 327–340.
- Ben-Porath, E. (1998). Correlation without mediation: Expanding the set of equilibrium outcomes by “cheap” pre-play procedures. *Journal of Economic Theory* 80(1), 108–122.
- Bergemann, D. and S. Morris (2016). Bayes correlated equilibrium and the comparison of information structures in games. *Theoretical Economics* 11(2), 487–522.

- Beyhaghi, H. and E. Tardos (2018). Two-sided matching with limited interviews. Technical report, Mimeo, Cornell University.
- Calvó-Armengol, A. (2006). The set of correlated equilibria of 2x2 games. *mimeo*.
- Canovas, S. G., P. Hansen, and B. Jaumard (1999). Nash equilibria from the correlated equilibria viewpoint. *International Game Theory Review* 1(01), 33–44.
- Crawford, V. P. (1990). Explicit communication and bargaining outcome. *The American Economic Review* 80(2), 213–219.
- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Echenique, F., R. Gonzalez, A. J. Wilson, and L. Yariv (2022). Top of the batch: Interviews and the match. *American Economic Review: Insights* 4(2), 223–238.
- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Evangelista, F. S. and T. Raghavan (1996). A note on correlated equilibrium. *International Journal of Game Theory* 25, 35–41.
- Feddersen, T. and W. Pesendorfer (1997). Voting behavior and information aggregation in elections with private information. *Econometrica: Journal of the Econometric Society*, 1029–1058.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International*

Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings 12, pp. 131–144. Springer.

- Forges, F. (2020). Correlated equilibria and communication in games. *Complex Social and Behavioral Systems: Game Theory and Agent-Based Models*, 107–118.
- Gerardi, D. (2004). Unmediated communication in games with complete and incomplete information. *Journal of Economic Theory* 114(1), 104–131.
- Gerardi, D. and L. Yariv (2007). Deliberative voting. *Journal of Economic theory* 134(1), 317–338.
- Goeree, J. K. and L. Yariv (2011). An experimental study of collective deliberation. *Econometrica* 79(3), 893–921.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.
- Kleiner, A., B. Moldovanu, and P. Strack (2021). Extreme points and majorization: Economic applications. *Econometrica* 89(4), 1557–1593.
- Kleiner, A., B. Moldovanu, P. Strack, and M. Whitmeyer (2024). The extreme points of fusions. *arXiv preprint arXiv:2409.10779*.
- Lehrer, E. and S. Sorin (1997). One-shot public mediated talk. *Games and Economic Behavior* 20(2), 131–148.
- Lopomo, G., L. M. Marx, and P. Sun (2011). Bidder collusion at first-price auctions. *Review of Economic Design* 15(3), 177–211.

- McAfee, R. P. and J. McMillan (1992). Bidding rings. *The American Economic Review*, 579–599.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.
- Nash, J. F. (1950). Non-cooperative games.
- Nau, R., S. G. Canovas, and P. Hansen (2004). On the geometry of nash equilibria and correlated equilibria. *International Journal of Game Theory* 32, 443–453.
- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Palfrey, T. R. and H. Rosenthal (1983). A strategic calculus of voting. *Public Choice* 41(1), 7–53.
- Pavlov, G. (2023). Correlated equilibria and communication equilibria in all-pay auctions. *Review of Economic Design*, 1–33.
- Peeters, R. and J. Potters (1999). On the structure of the set of correlated equilibria in two-by-two bimatrix games.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.
- Viossat, Y. (2010). Properties and applications of dual reduction. *Economic theory* 44, 53–68.

Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.

Yang, K. H. and A. K. Zentefis (2024). Monotone function intervals: Theory and applications. *American Economic Review* 114(8), 2239–2270.