

# Improvable Equilibria

---

**Kirill Rudov** - UC Berkeley

**Fedor Sandomirskiy** - Princeton

**Leeat Yariv** - Princeton

Becker Friedman Institute, August 22, 2024

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

**This project:** When is there potential value in correlation?

## Normal-form game

$$\Gamma = \left( N, (A_i)_{i \in N}, (u_i: A \rightarrow \mathbb{R})_{i \in N} \right)$$

- $N = \{1, \dots, n\}$  is finite set of players
- $A_i$  is a finite set of actions of player  $i$
- $A = \prod_{i \in N} A_i$  is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$  is utility of player  $i$

# Correlated Equilibria (CE)

## Definition

A distribution  $\mu \in \Delta(A)$  is a correlated equilibrium if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all  $i \in N$  and all  $a_i, a'_i \in A_i$

**Interpretation:**  $\mu$  as generated by a mediator, where agents best respond by adhering

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?

## Definition

A Nash equilibrium is **improvable** if it is not an extreme point of the set of correlated equilibria



- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?

## Definition

A Nash equilibrium is **improvable** if it is not an extreme point of the set of correlated equilibria

- Improvable  $\Leftrightarrow$  **any** strictly convex objective can be strictly improved

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?

## Definition

A Nash equilibrium is **improvable** if it is not an extreme point of the set of correlated equilibria

- Improvable  $\Leftrightarrow$  **any** strictly convex objective can be strictly improved
- Agnostic about designer's objective

## Example: Improving Welfare of Nash

The Game of Chicken by [Aumann \(1974\)](#):



	Dare	Chicken
Dare	0, 0	4, 1
Chicken	1, 4	3, 3

## Example: Improving Welfare of Nash

	Dare	Chicken
Dare	0, 0	4, 1
Chicken	1, 4	3, 3

- Mixed Nash:  $(1/2, 1/2)$  for both players


## Example: Improving Welfare of Nash

	Dare	Chicken
Dare	0, 0	4, 1
Chicken	1, 4	3, 3

- Mixed Nash:  $(1/2, 1/2)$  for both players
- Correlated equilibrium can increase welfare by shifting weight from  $(0,0)$

## Example: Improving Welfare of Nash

		Dare	Chicken
Dare	0, 0	4, 1	
Chicken	1, 4	3, 3	



- Mixed Nash:  $(1/2, 1/2)$  for both players
- Correlated equilibrium can increase welfare by shifting weight from  $(0,0)$
- But the mixed Nash is **not improvable**
  - It is a unique optimum for the objective corresponding to weight on diagonal entries,  $(\text{Dare}, \text{Dare})$  and  $(\text{Chicken}, \text{Chicken})$

- Welfare maximization is just one objective
- An objective can maximize likelihood of certain profiles, minimize entropy of the joint distribution over profiles, etc.
- An improvable NE can be improved **no matter the objective**
- Improvability is very demanding

- Conditions for improvability
- Linear objectives
- Applications
- Symmetric games



## Conditions for Improvability

---

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

Complete detail-free characterization of improvable Nash equilibria

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved, either by introducing correlation, or by reducing randomness

## Theorem 1

In a generic  $n$ -player game, a mixed NE is improvable  $\iff \geq 3$  agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved, either by introducing correlation, or by reducing randomness  
 $\Rightarrow$  2-agent games not representative

When can the genericity assumption be dropped?



When can the genericity assumption be dropped?

- in games with 2 actions per player
- in any game, by considering *regular* NE only

When can the genericity assumption be dropped?

- in games with 2 actions per player
- in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small perturbation of payoffs

When can the genericity assumption be dropped?

- in games with 2 actions per player
- in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small perturbation of payoffs

## Theorem 1'

In any game, a regular mixed NE is improvable  $\iff \geq 3$  agents randomize

When can the genericity assumption be dropped?

- in games with 2 actions per player
- in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small perturbation of payoffs

## Theorem 1'

In any game, a regular mixed NE is improvable  $\iff \geq 3$  agents randomize

- In a generic game, any NE is regular (Harsanyi, 1973)

When can the genericity assumption be dropped?

- in games with 2 actions per player
- in any game, by considering *regular* NE only

**Definition** (informal): a NE is regular if it is stable under small perturbation of payoffs

## Theorem 1'

In any game, a regular mixed NE is improvable  $\iff \geq 3$  agents randomize

- In a generic game, any NE is regular (Harsanyi, 1973)
- Hence, Theorem 1'  $\Rightarrow$  Theorem 1

**Rough intuition:** When many agents randomize, there are too many ways to correlate their actions, one must be beneficial

Focus on a particular example to illustrate

## Why Theorems Hold

- Game with  $n$  agents, each with two actions

## Why Theorems Hold

- Game with  $n$  agents, each with two actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$



# Why Theorems Hold

- Game with  $n$  agents, each with two actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints

- Game with  $n$  agents, each with two actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq 1 + k$

- Game with  $n$  agents, each with two actions
- If  $\mu$  is a CE, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$  constraints
- **Winkler (1988)**: if  $k$  linear constraints are imposed on the set of all distributions  $\Delta(A)$ , extreme distributions have support  $\leq 1 + k$
- $\Rightarrow$  support of an extreme CE  $\mu$  is bounded by  $1 + 2n$

## Why Theorems Hold

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  agents mixing

## Why Theorems Hold

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  agents mixing
- The support of  $\nu$  contains  $2^k$  action profiles

## Why Theorems Hold

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  agents mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be non-improvable,

$$2^k \leq 1 + 2n$$

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  agents mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be non-improvable,

$$2^k \leq 1 + 2n$$

- At most  $\log_2(1 + 2n)$  out of  $n$  agents can randomize

## Why Theorems Hold

- Suppose  $\nu$  is a Nash equilibrium with the  $k$  agents mixing
- The support of  $\nu$  contains  $2^k$  action profiles
- $\Rightarrow$  For  $\nu$  to be non-improvable,

$$2^k \leq 1 + 2n$$

- At most  $\log_2(1 + 2n)$  out of  $n$  agents can randomize
- In fact, only 2 agents can randomize

► details



## Linear Objectives

---

Utilitarian welfare is a common objective, which is linear:

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

# Linear Objectives and Utilitarian Welfare

Utilitarian welfare is a common objective, which is linear:

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

In [Aumann \(1974\)](#)'s example, we saw a NE whose utilitarian welfare could be strictly improved

- But that was a  $2 \times 2$  game...

# Linear Objectives and Utilitarian Welfare

Utilitarian welfare is a common objective, which is linear:

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$


In [Aumann \(1974\)](#)'s example, we saw a NE whose utilitarian welfare could be strictly improved

- But that was a  $2 \times 2$  game...

**Question:** When is a NE improvable in terms of utilitarian welfare, or other linear objectives?

# Improving Linear Objectives

Maximization of a linear objective over a polytope  $P$ :




Two cases:

- For a generic objective, the unique optimum is at an extreme point
- In the knife-edge case, the whole face of  $P$  can be optimal

# Improving Linear Objectives

Maximization of a linear objective over a polytope  $P$ :



Two cases:

- For a generic objective, the unique optimum is at an extreme point
- In the knife-edge case, the whole face of  $P$  can be optimal

## Corollary

NE is improvable  $\implies$  a generic linear objective can be strictly improved

**Question:** When can the welfare objective

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

be strictly improved?

**Question:** When can the welfare objective

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

be strictly improved?

## **Proposition 1**

In generic game  $\Gamma$ , welfare of an improvable NE can be strictly improved



**Question:** When can the welfare objective

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

be strictly improved?

## Proposition 1

In generic game  $\Gamma$ , welfare of an improvable NE can be strictly improved

**Intuition:** Perturb utilities  $u'_i(s) = u_i(s) + \delta_i(s_{-i})$ . This

- does not affect the incentive constraints
- affects  $W$

$\Rightarrow$  the hyperplane  $W = \text{const}$  cannot be parallel to a face of  $CE(\Gamma)$  in a generic  $\Gamma$

**Question:** When can the welfare objective

$$W(\mu) = \sum_{i \in N} \sum_{s \in S} u_i(s) \mu(s)$$

be strictly improved?

## Proposition 1

In generic game  $\Gamma$ , welfare of an improvable NE can be strictly improved

**Intuition:** Perturb utilities  $u'_i(s) = u_i(s) + \delta_i(s_{-i})$ . This

- does not affect the incentive constraints
- affects  $W$

$\Rightarrow$  the hyperplane  $W = \text{const}$  cannot be parallel to a face of  $CE(\Gamma)$  in a generic  $\Gamma$

**Other linear objectives:** conditions for strict improvability from Farkas lemma [details](#)

## **Sample Applications**

---

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are improvable

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are improvable

**Other Applications:** games where agents want to mismatch actions of others

## Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters:  $D$  and  $R$ ,  $|R| > |D|$
- Voters in  $D$  get utility of 1 if  $d$ -candidate wins and 0 otherwise
- Voters in  $R$  get utility of 1 if  $r$ -candidate wins and 0 otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation:  $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of  $c$ , all equilibria involve at least one group all mixing

- $\Rightarrow$  These equilibria are improvable

**Other Applications:** games where agents want to mismatch actions of others

- e.g., network games (with substitutes), congestion games, all-pay auctions, Boston matching mechanism



# Symmetric Games

---

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

## **Theorem 2**

In any symmetric game with  $n \geq 3$  players, any completely mixed symmetric NE is improvable

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

## **Theorem 2**

In any symmetric game with  $n \geq 3$  players, any completely mixed symmetric NE is improvable

- No genericity or regularity assumptions
- Any pure strategy must be played with a positive probability

# Symmetric Games: Highlights

- The result is, in fact, stronger:

For  $n \geq 3$ , completely mixed symmetric NE cannot be extreme within the (smaller!) set of *symmetric CE*

# Symmetric Games: Highlights

- The result is, in fact, stronger:

For  $n \geq 3$ , completely mixed symmetric NE cannot be extreme within the (smaller!) set of *symmetric CE*

- $\Rightarrow$  a generic *symmetric* linear objective can be strictly improved, e.g., utilitarian welfare

# Symmetric Games: Highlights

- The result is, in fact, stronger:

For  $n \geq 3$ , completely mixed symmetric NE cannot be extreme within the (smaller!) set of *symmetric CE*

- $\Rightarrow$  a generic *symmetric* linear objective can be strictly improved, e.g., utilitarian welfare

**Take-away:** caution when focusing on symmetric equilibria in symmetric games

## What Improvements Look Like

For an improvable NE, any strictly convex objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?



# What Improvements Look Like

For an improvable NE, any strictly convex objective can be strictly improved by switching to an extreme CE

**Question:** What is the structure of extreme CE?

- For general games, we only know that extreme CE have small support
- For symmetric games and symmetric CE, we can say more

- **Observation:** for a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable

- **Observation:** for a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known

- **Observation:** for a symmetric CE, the random variables  $a_1, \dots, a_n$  are exchangeable
- If  $n \rightarrow \infty$ , the structure of exchangeable distributions is well-known

## Theorem (de Finetti)

Any infinite exchangeable sequence  $a_1, a_2, a_3 \dots$  is a mixture of i.i.d.

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### **Proposition 2**

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parametric family of extreme CE

# Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parametric family of extreme CE
- A radical dimension reduction



## Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

### Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parametric family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

# Extreme Symmetric CE with Many Players

- Consider a symmetric game with  $m$  actions per player
- Assume the number of players  $n$  is large

## Proposition 2

Any extreme symmetric CE can be approximated by a mixture of  $m(m - 1) + 1$  i.i.d. distributions

- For  $m = 2$ , a mixture of 3 i.i.d. distributions  $\Rightarrow$  5-parametric family of extreme CE
- A radical dimension reduction

**Question:** What if we want the exact result, not an approximation?

- A version of Proposition 2 holds: sampling without replacement instead of i.i.d.

# Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

# Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- a tension between equilibrium randomness and non-improvability
- detail-free criterion for non-improvability in various settings

# Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- a tension between equilibrium randomness and non-improvability
- detail-free criterion for non-improvability in various settings

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design

# Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- a tension between equilibrium randomness and non-improvability
- detail-free criterion for non-improvability in various settings

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design


# Thank you!

## Corollary


If a generic game has a unique correlated equilibrium  $\nu$ , then  $\nu$  is either:

- A pure Nash equilibrium, or
  - A Nash equilibrium where exactly two agents randomize
- 
- Examples: some congestion games, Cournot competition



# Simulations



(a)  $NE/2u$



(b) Fraction of CE  $> NE^*$





## Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If  $\mu$  is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |S_i| \cdot (|S_i| - 1)$$

## Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium,  $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ :

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

## Regularity of Generic games (**Harsanyi, 1973**)

In a generic game, any Nash equilibrium is regular

# General linear objectives

- Consider a NE  $\nu$
- For simplicity,  $\nu$  has full support
- By Farkas lemma, a linear objective  $L$  can be improved for  $\nu \iff L$  **cannot** be expressed as

$$L(\mu) = C + \sum_{i, a_i, a'_i, a_{-i}} \mu(a) \cdot \lambda_i(a_i, a'_i) \cdot (u_i(a_i, a_{-i}) - u_i(a'_i, a_{-i}))$$

for some  $\lambda_i(a_i, a'_i) \geq 0$ .

- For improvable NE  $\nu$ , “bad”  $L$  form a lower-dimensional subspace

# Extreme Symmetric CE with Any Number of Players

Consider  $n$  players with  $m$  actions each

## Proposition

Any extreme symmetric CE can be obtained as follows:

- there are  $M$  urns, each with  $n$  balls labeled by actions

$$1 \leq M \leq m(m-1) + 1$$

- an urn is selected at random according to  $p \in \Delta_M$ , secretly from players
- players draw balls sequentially without replacement
- $i$ 's action = her ball's label, no incentive to deviate

**Remark:** If  $n$  is large, sampling without replacement can be approximated by i.i.d.

## Bayesian games

---

## Bayesian game

$$\mathcal{B} = \left( N, (A_i)_{i \in N}, (T_i)_{i \in N}, \tau \in \Delta(T), (u_i: A \times T_i \rightarrow \mathbb{R})_{i \in N} \right)$$

- Each player  $i \in N$  has a type  $t_i \in T_i$
- Profile of types  $(t_1, \dots, t_n) \in T$  sampled from  $\tau$
- Each player  $i$  observes her realized type
- Utility  $u_i: A \times T_i \rightarrow \mathbb{R}$  depends on the action profile and  $i$ 's type

**Technical assumption:** sets of types  $T_i$  are finite

# Bayesian Correlated Equilibria (BCE)

## Definition

A joint distribution  $\mu \in \Delta(A \times T)$  is a Bayesian correlated equilibrium if

- The marginal on  $T$  coincides with  $\tau$
- For each player  $i$ , type  $t_i$ , recommended action  $a_i$ , and deviation  $a'_i$ ,

$$\sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a_i, t_i, a_{-i}) \geq \sum_{(a_{-i}, t_{-i})} \mu((a_i, t_i), (a_{-i}, t_{-i})) u_i(a'_i, t_i, a_{-i})$$

**Interpretation:** a mediator having access to realized types recommends actions to each player. Two aspects:

1. *Ex-ante coordination:* a source of correlated randomness (as in CE)
2. *Information sharing:* providing  $i$  more info about  $t_{-i}$  than contained in  $t_i$

**Remark:** Bergemann and Morris (2016) allow for a broader class of BCE, where player  $i$  observes a noisy signal about her type

# Induced Complete Information Game

We can associate a complete information normal form game  $\Gamma_{\mathcal{B}}$  with  $\mathcal{B}$ :

- Replace  $A_i$  with set of functions  $\sigma_i : T_i \rightarrow A_i$
- $\Sigma_i$  is the set of all such  $\sigma_i$
- Utility  $v_i : \Sigma \rightarrow \mathbb{R}$  is given by

$$v_i(\sigma) = \sum_{t \in T} \tau(t) \cdot u_i((\sigma_1(t_1), \dots, \sigma_n(t_n)), t_i)$$

## Induced Complete Information Game

$$\Gamma_{\mathcal{B}} = (N, (\Sigma_i)_{i \in N}, (v_i)_{i \in N})$$

**Question:** What is a relation between CE of  $\Gamma_{\mathcal{B}}$  and BCE of  $\mathcal{B}$ ?

# Induced complete information game

## Relationship between equilibria in $\Gamma_{\mathcal{B}}$ and $\mathcal{B}$

CE in  $\Gamma_{\mathcal{B}} \Leftrightarrow$  ex-ante coordination in  $\mathcal{B}$  with no information sharing

- i.e., BCE such that  $a_i$  is independent of  $t_{-i}$  conditionally on  $t_i$

Nash in  $\Gamma_{\mathcal{B}} \Leftrightarrow$  Bayes-Nash in  $\mathcal{B}$

**Observation:** Generic  $\mathcal{B}$  leads to generic  $\Gamma_{\mathcal{B}}$

- $\Rightarrow$  we can apply our theorem to  $\Gamma_{\mathcal{B}}$  to learn about generic  $\mathcal{B}$

## Corollary

For a generic Bayesian game, a Bayes-Nash equilibrium is improvable via ex-ante coordination  $\Leftrightarrow$  at least 3 players randomize

Applies to Bayesian games where agents randomize in equilibrium, e.g., costly voting with private types (Feddersen and Pesendorfer, 1997) and contests (Baranski and Goel, 2024)



## References

---

- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Baranski, A. and S. Goel (2024). Contest design with a finite type-space. *to appear*.
- Bergemann, D. and S. Morris (2016). Bayes correlated equilibrium and the comparison of information structures in games. *Theoretical Economics* 11(2), 487–522.
- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Feddersen, T. and W. Pesendorfer (1997). Voting behavior and information aggregation in elections with private information. *Econometrica: Journal of the Econometric Society*, 1029–1058.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings* 12, pp. 131–144. Springer.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.

- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Palfrey, T. R. and H. Rosenthal (1983). A strategic calculus of voting. *Public Choice* 41(1), 7–53.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.
- Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.