

Improvable Equilibria

Kirill Rudov - UC Berkeley

Fedor Sandomirskiy - Princeton

Leeat Yariv - Princeton

Stony Brook Game Theory Festival, July 15, 2024

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

Correlated equilibrium (Aumann, 1974) generalizes Nash equilibrium to allow correlation

- Implementable via mediation, communication, joint randomization, etc.

This project: When is there potential value in correlation?

Games on a Shoestring

Normal-form game is given by $(N, (A_i)_{i \in N}, (u_i)_{i \in N})$, where

- $N = \{1, \dots, n\}$ is finite set of players
- A_i is a finite set of actions of agent i
- $A = \prod_{i \in N} A_i$ is the set of action profiles
- $u_i: A \rightarrow \mathbb{R}$ is utility of agent i

Definition

A distribution $\mu \in \Delta(A)$ is a **correlated equilibrium** if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all $i \in N$ and all $a_i, a'_i \in A_i$

Definition

A distribution $\mu \in \Delta(A)$ is a **correlated equilibrium** if

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

for all $i \in N$ and all $a_i, a'_i \in A_i$

Interpretation: μ as generated by a mediator, where agents best respond by adhering

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?
- If not, *no matter the objective*, we could always benefit from introducing correlation

- The set of correlated equilibria is a convex set
- **Bauer's Maximum Principle:** Any linear or convex objective attains its maximum at an extreme point (uniquely with strict convexity)
- **Our Question:** When is a Nash equilibrium an extreme point of the set of correlated equilibria?
- If not, *no matter the objective*, we could always benefit from introducing correlation

Definition

A Nash equilibrium is **improvable** if it is not an extreme point of the set of correlated equilibria

Improvability of Nash Equilibria

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Improvability of Nash Equilibria

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

Improvability of Nash Equilibria

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved either by introducing correlation or by reducing randomness

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved either by introducing correlation or by reducing randomness
 \Rightarrow 2-agent games not representative

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved either by introducing correlation or by reducing randomness
 \Rightarrow 2-agent games not representative

The genericity assumption can be dropped

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved either by introducing correlation or by reducing randomness
 \Rightarrow 2-agent games not representative

The genericity assumption can be dropped

- in games with 2 actions per player

Theorem 1

In a generic n -player game, a mixed **Nash equilibrium is improvable** \iff
three or more agents randomize

Complete detail-free characterization of improvable Nash equilibria

- Pure equilibria are non-improvable (trivial)
- Equilibria with exactly 2 randomizing agents are non-improvable (Cripps, 1995)
- If 3 or more agents randomize, *any* objective can be improved either by introducing correlation or by reducing randomness
 \Rightarrow 2-agent games not representative

The genericity assumption can be dropped

- in games with 2 actions per player
- in any game, by considering *regular* Nash equilibria only

Rough intuition: When many agents randomize, there are too many ways to correlate their actions, one must be beneficial

Rough intuition: When many agents randomize, there are too many ways to correlate their actions, one must be beneficial

Focus on particular example to illustrate

Intuition through Example

- Game with n agents, each with two actions

Intuition through Example

- Game with n agents, each with two actions
- If μ is a correlated equilibrium, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

Intuition through Example

- Game with n agents, each with two actions
- If μ is a correlated equilibrium, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints

Intuition through Example

- Game with n agents, each with two actions
- If μ is a correlated equilibrium, must satisfy incentive constraints

$$\sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a_i, a_{-i}) \geq \sum_{a_{-i} \in A_{-i}} \mu(a_i, a_{-i}) u_i(a'_i, a_{-i})$$

- $2n$ constraints
- **Winkler (1988)** \Rightarrow support of any extreme μ is bounded by $1 + 2n$

- Suppose ν is a Nash equilibrium with the k agents mixing

Intuition through Example

- Suppose ν is a Nash equilibrium with the k agents mixing
- The support of ν contains 2^k action profiles

Intuition through Example

- Suppose ν is a Nash equilibrium with the k agents mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be non-improvable,

$$2^k \leq 1 + 2n$$

Intuition through Example

- Suppose ν is a Nash equilibrium with the k agents mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be non-improvable,

$$2^k \leq 1 + 2n$$

- At most $\log_2(1 + 2n)$ out of n agents can randomize

Intuition through Example

- Suppose ν is a Nash equilibrium with the k agents mixing
- The support of ν contains 2^k action profiles
- \Rightarrow For ν to be non-improvable,

$$2^k \leq 1 + 2n$$

- At most $\log_2(1 + 2n)$ out of n agents can randomize
- In fact, only 2 agents can randomize (requires more careful analysis [▶ details](#))

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0, otherwise
- Voters in R get utility of 1 if r -candidate wins and 0, otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0, otherwise
- Voters in R get utility of 1 if r -candidate wins and 0, otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0, otherwise
- Voters in R get utility of 1 if r -candidate wins and 0, otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are improvable

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0, otherwise
- Voters in R get utility of 1 if r -candidate wins and 0, otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are improvable

Other Applications: games where agents want to mismatch actions of others

Costly voting model of Palfrey and Rosenthal (1983):

- Two finite groups of voters: D and R , $|R| > |D|$
- Voters in D get utility of 1 if d -candidate wins and 0, otherwise
- Voters in R get utility of 1 if r -candidate wins and 0, otherwise
- Majority voting (among those who participate), ties broken randomly
- Costly participation: $c > 0$

Palfrey and Rosenthal (1983): For intermediate values of c , all equilibria involve at least one group all mixing

- \Rightarrow These equilibria are improvable

Other Applications: games where agents want to mismatch actions of others

- e.g., network games (with substitutes), congestion games, all-pay auctions, Boston matching mechanism

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

Corollary

In a generic symmetric game with $n \geq 3$ players, any non-pure symmetric Nash equilibrium is improvable

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

Corollary

In a generic symmetric game with $n \geq 3$ players, any non-pure symmetric Nash equilibrium is improvable

What if we insist that improvement be done via a *symmetric* correlated equilibrium?

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

Corollary

In a generic symmetric game with $n \geq 3$ players, any non-pure symmetric Nash equilibrium is improvable

What if we insist that improvement be done via a *symmetric* correlated equilibrium?

- **Theorem 2:** for $n \geq 3$, non-pure symmetric equilibria remain improvable even within the set of symmetric correlated equilibria

Symmetric Games

- In many applications, strategic interactions are symmetric
- When are symmetric equilibria improvable?

Corollary

In a generic symmetric game with $n \geq 3$ players, any non-pure symmetric Nash equilibrium is improvable

What if we insist that improvement be done via a *symmetric* correlated equilibrium?

- **Theorem 2:** for $n \geq 3$, non-pure symmetric equilibria remain improvable even within the set of symmetric correlated equilibria

Take-away: caution when focusing on symmetric equilibria in symmetric games

Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- identifies a tension between equilibrium randomness and non-improvability
- offers detail-free criterion for non-improvability: “almost pureness”

Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- identifies a tension between equilibrium randomness and non-improvability
- offers detail-free criterion for non-improvability: “almost pureness”

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design

Conclusions

Several papers effectively show non-improvability in specific contexts:

- Tullock contests, Cournot and Bertrand, patent races, location games (Einy, Haimanko, and Lagziel, 2022)
- First-price auctions (Feldman, Lucier, and Nisan, 2016)
- Convex potential games (Neyman, 1997; Ui, 2008)
- Two-player normal-form games (Cripps, 1995)

Our paper:

- identifies a tension between equilibrium randomness and non-improvability
- offers detail-free criterion for non-improvability: “almost pureness”

Ongoing:

- Incomplete information
- “Correlated implementation” in mechanism design

Thank you!

Corollary

If a generic game has a unique correlated equilibrium ν , then ν is either:


- A pure Nash equilibrium, or
- A Nash equilibrium where exactly two agents randomize

Corollary


If a generic game has a unique correlated equilibrium ν , then ν is either:

- A pure Nash equilibrium, or
 - A Nash equilibrium where exactly two agents randomize
-
- Examples: some congestion games, Cournot competition



Simulations



(a) $NE/2u$



(b) Fraction of CE $> NE^*$



Support Size of Extreme Correlated Equilibria (follows from **Winkler (1988)**)

If μ is an extreme correlated equilibrium, then

$$\text{supp}(\mu) \leq 1 + \sum_{i \in N} |S_i| \cdot (|S_i| - 1)$$

Support Size of Regular Nash Equilibria (**McKelvey and McLennan, 1997**)

For a regular Nash equilibrium, $\nu = (\nu_1, \nu_2, \dots, \nu_n)$:

$$\text{supp}(\nu_i) - 1 \leq \sum_{j \neq i} (\text{supp}(\nu_j) - 1), \quad \text{for any player } i$$

Regularity of Generic games (**Harsanyi, 1973**)

In a generic game, any Nash equilibrium is regular

References

- Aumann, R. J. (1974). Subjectivity and correlation in randomized strategies. *Journal of mathematical Economics* 1(1), 67–96.
- Cripps, M. (1995). Extreme correlated and nash equilibria in two-person games.
- Einy, E., O. Haimanko, and D. Lagziel (2022). Strong robustness to incomplete information and the uniqueness of a correlated equilibrium. *Economic Theory* 73(1), 91–119.
- Feldman, M., B. Lucier, and N. Nisan (2016). Correlated and coarse equilibria of single-item auctions. In *Web and Internet Economics: 12th International Conference, WINE 2016, Montreal, Canada, December 11-14, 2016, Proceedings* 12, pp. 131–144. Springer.
- Harsanyi, J. C. (1973). Oddness of the number of equilibrium points: a new proof. *International Journal of Game Theory* 2, 235–250.
- McKelvey, R. D. and A. McLennan (1997). The maximal number of regular totally mixed nash equilibria. *Journal of Economic Theory* 72(2), 411–425.
- Neyman, A. (1997). Correlated equilibrium and potential games. *International Journal of Game Theory* 26, 223–227.
- Palfrey, T. R. and H. Rosenthal (1983). A strategic calculus of voting. *Public Choice* 41(1), 7–53.
- Ui, T. (2008). Correlated equilibrium and concave games. *International Journal of Game Theory* 37, 1–13.

Winkler, G. (1988). Extreme points of moment sets. *Mathematics of Operations Research* 13(4), 581–587.