

Extension to time dependent problems

Christophe Prud'homme

prudhomme@unistra.fr

February 13, 2023

CeMosis - <http://www.cemosis.fr>

IRMA

Université de Strasbourg

Linear Parabolic Problems

New ingredients/challenges:

- Simultaneous dependence on both time and parameters.
 - "Time" as an additional (albeit special) parameter. //
- Output, $s = s(t; \mu)$, is a function of time (and parameter).
 - Important for applications, e.g., control.
 - A posteriori error bounds (no "compliance" \Rightarrow dual problem). /
- Sampling procedure.
 - Greedy algorithm for parameter-time case.
 - Unknown "control" input.
- Dimension N of RB space.
 - (• Advection-dominated problems.

Problem statement

Figure

Given $\mu \in \mathcal{D} \subset \mathbb{R}^P$, evaluate $t \in (0, t_f]$

$$s^e(t; \mu) = \ell(u^e(x; t; \mu); \mu) \quad (1)$$

where $u^e(x; t; \mu) \in L^2(0, t_f; X^e(\Omega)) \cup C^0([0, t_f]; L^2(\Omega))$ satisfies

$$m \left(\frac{\partial u^e}{\partial t}(x; t; \mu), v; \mu \right) + \underline{a}(u^e(x; t; \mu), v; \mu) \quad (2)$$

$$= \underline{f(v; \mu)g(t)}, \quad \forall v \in X^e \text{ control}$$

with initial condition $u_0 = 0$. (Note: extension to nonzero initial conditions are briefly discussed below).

$$u^e(t=0) = \underline{u_0} = 0$$

- μ : input parameter $-\mu = (\mu_1, \mu_2, \dots, \mu_P)$; P -tuple
- \mathcal{D} : parameter domain in \mathbb{R}^P ;
- Ω : spatial domain in \mathbb{R}^d ;
- s^e : output;
- ℓ : output functional;
- u^e : field variable;
- X^e : function space $(H_0^1(\Omega))^\nu \subset X^e \subset (H^1(\Omega))^\nu$ we assume here $\nu = 1$, with inner product $(\mathbf{w}, \mathbf{v})_{X^e}, \forall \mathbf{w}, \mathbf{v} \in X^e$, and induced norm $\|\mathbf{w}\|_{X^e} = \sqrt{(\mathbf{w}, \mathbf{w})_{X^e}}, \quad \forall \mathbf{w} \in X^e$.

$\nu = 1$

Note Ω is parameter-independent: the reduced basis requires a common spatial configuration, i.e., a reference domain Ω_{ref} . Introduce a piecewise affine mapping $\mathcal{T}(\cdot; \mu) : \Omega \rightarrow \Omega_o(\mu)$

$$a_o(w_o, v_o; \mu) \text{ over } \Omega_o(\mu)$$

$$\Downarrow$$

$$\mathcal{T}(\cdot; \mu)^{-1} : \Omega_o(\mu) \rightarrow \Omega_{\text{ref}} \equiv \Omega \quad (\Omega_{\text{ref}} = \Omega_o(\mu_{\text{ref}}))$$

$$\Downarrow$$

$$a(w, v; \mu) \text{ over } \Omega$$

$$\text{where } a(w, v; \mu) = a_o(w_o \circ \mathcal{T}_\mu, v_o \circ \mathcal{T}_\mu; \mu)$$

We henceforth assume that the problem is already mapped to the reference domain.

Linear forms and functions

$$\begin{aligned} f(\cdot; \mu) &: \text{linear, affine in } \mu, \\ &: \mathbf{X}^e\text{-bounded, } \forall \mu \in \mathcal{D} \\ g(\cdot) &: L^2(0, t_f) \quad \text{"control" input} \\ \ell(\cdot; \mu) &: \text{linear, affine in } \mu, \\ &: L^2(\Omega)\text{-bounded, } \forall \mu \in \mathcal{D} \end{aligned}$$

$$\begin{aligned} a(\cdot, \cdot; \mu) &: \text{bilinear, affine in } \mu, \\ &\quad \left(\text{symmetric,} \right) \\ &\quad X^e\text{-continuous,} \\ &\quad X^e\text{-coercive form,} \quad \forall \mu \in \mathcal{D}; \\ m(\cdot, \cdot; \mu) &: \text{bilinear, affine in } \mu, \\ &\quad \text{symmetric,} \\ &\quad L^2(\Omega)\text{-continuous,} \\ &\quad L^2(\Omega)\text{-coercive form,} \quad \forall \mu \in \mathcal{D}; \end{aligned} \tag{3}$$

Note: a may satisfy a weak coercivity condition.

Require

also $\ell(v; \mu), f(v; \mu)$

Figure

$$a(w, v; \mu) = \sum_{q=1}^{Q_a} \Theta_a^q(\mu) a^q(w, v)$$

$$m(w, v; \mu) = \sum_{q=1}^{Q_m} \Theta_m^q(\mu) m^q(w, v)$$

where


- $\Theta_{a,m}^q : \mathcal{D} \rightarrow \mathbb{R}$, μ -dependent functions; representing coefficients, geometry, ...
- a^q and m^q μ -independent forms.

Note: affine assumption may be relaxed.

Temporal Discretization: Finite Difference

$$\left. \frac{\partial u}{\partial t}(t^k; \mu) \approx \frac{u(t^k; \mu) - u(t^{k-1}; \mu)}{\Delta t} \right\}$$

- Euler Backward
- Crank-Nicolson (advection-dominated problems)



C. Prud'homme

Extension to time dependent problems

$$m\left(\frac{\partial u}{\partial t}, v; \mu\right) + a(u, v; \mu) = f(v; \mu) g(t) \quad (1)$$

$$\frac{\partial u(t^k)}{\partial t} \approx \frac{u(t^{k+1}) - u(t^k)}{\Delta t}$$

et $t = t^k$ avec DF progressive

$$m\left(\frac{u(t^{k+1}) - u(t^k)}{\Delta t}, v; \mu\right) + a(u(t^k), v; \mu) = f(v; \mu) g(t^k)$$

Problem Statement

Given $\mu \in \mathcal{D} \subset \mathbb{R}^P$, evaluate

$$\forall k \in \mathbb{K} \quad \triangle$$

$$s^k(\mu) \equiv s(t^k; \mu) = \ell(u(t^k; \mu); \mu)$$

where $u^k(\mu) \equiv u(t^k; \mu) \in X$ satisfies $u_0 = 0$

$$m\left(\frac{u(t^k; \mu) - u(t^{k-1}; \mu)}{\Delta t}, v; \mu\right) + a(u(t^k; \mu), v; \mu) = f(v; \mu)g(t^k), \quad \forall v \in X$$

Note: We directly drop the superscript \mathcal{N} , i.e., $\mathbf{X} = \mathbf{X}^{\mathcal{N}}$,

$$u(t^k; \mu) = u^{\mathcal{N}}(t^k; \mu), s(t^k; \mu) = s^{\mathcal{N}}(t^k; \mu)$$

C. Prud'homme

Extension to time dependent problems

$$\frac{1}{\Delta t} m(u(t^k), v; \mu) + a(u(t^k), v; \mu) = f(v; \mu)g(t^k)$$

$$\left(\frac{1}{\Delta t} M + A\right) U^k = F g(t^k) + \frac{1}{\Delta t} M U^{k-1} + \frac{m}{\Delta t}(u(t^{k-1}), v; \mu)$$

$$U^k = \begin{pmatrix} u_1(t^k) \\ \vdots \\ u_{\mathcal{CP}}(t^k) \end{pmatrix} \uparrow \mathcal{CP} \quad u(t^k) = \sum_{i=1}^{\mathcal{CP}} u_i(t^k) \varphi_i(x)$$

(*) $k = 1, \dots, K$

We shall

- **build** our reduced basis approximation upon "truth" solutions $u(t^k; \mu) \in X$
- **measure** the error in the reduced basis approximation relative to the "truth" solution $u(t^k; \mu) \in X$ (and $s(t^k; \mu)$);
 $(\Rightarrow \mathbf{u}(t^k; \mu)$ is a **calculable surrogate** for $\mathbf{u}^e(t; \mu)$).

Reduced Basis Approximation

We assume


- the form a is continuous and coercive (or inf-sup stable); and
- the form m is continuous and coercive;
- and the $\Theta_{m,a}^q(\mu)$, $1 \leq q \leq Q_{m,a}$, are smooth; then

$$\hookrightarrow \mathcal{M}^{\mathcal{N}K} \equiv \{u(t^k; \mu) \mid 1 \leq k \leq K, \forall \mu \in \mathcal{D}\}$$

lies on a smooth $P+1$ -dimensional manifold in \mathbf{X} .

To approximate $u(t^k; \mu)$, and hence $s(t^k; \mu)$,

we need not represent every possible function in $\mathcal{X}^{\mathcal{N}}$.




$$\mathcal{X}_N \subset \text{span} \{ u(t^k; \mu^m), 1 \leq k \leq K, 1 \leq m \leq M \} \quad (4)$$

To approximate $u(t^k; \mu)$, and hence $s(t^k; \mu)$,

Figure

we need not represent every possible function in $\mathbf{X}^{\mathcal{N}}$.




① solutions ED
utilisées
pour construire
la base
réduite

LOCALIZATION

To approximate $u(t^k; \mu)$, and hence $s(t^k; \mu)$,

Figure

we need not represent every possible function in $\mathcal{X}^{\mathcal{N}}$.



SMOOTHNESS

We define the Lagrangian RB space

$$X_N = \text{span} \{ \zeta^n, 1 \leq n \leq N \}, \quad 1 \leq N \leq N_{\max}$$

with mutually $(\cdot, \cdot)_X$ -orthonormal basis functions

$$\zeta^n \in X, \quad 1 \leq n \leq N_{\max}$$

We thus obtain

$$\mathbf{X}_N \subset X, \quad \dim(\mathbf{X}_N) = N, \quad 1 \leq N \leq N_{\max}$$

and

hierarchical spaces

$$X_1 \subset X_2 \subset \dots \subset X_{N_{\max}-1} \subset X_{N_{\max}} (\subset X)$$

The basis functions are constructed using a POD-Greedy algorithm outlined below.

Given $\mu \in \mathcal{D} \subset \mathbb{R}^P$, evaluate $\forall k \in \mathbb{K}$

$$s_N^k(\mu) \equiv s_N(t^k; \mu) = \ell(u_N(t^k; \mu); \mu)$$

where $u_N^k(\mu) \equiv u_N(t^k; \mu) \in X_N$ satisfies $u_{N,0} = 0$

$$m\left(\frac{u_N(t^k; \mu) - u_N(t^{k-1}; \mu)}{\Delta t}, v; \mu\right) + a(u_N(t^k; \mu), v; \mu) = f(v; \mu)g(t^k), \quad \forall v \in X_N.$$

\Rightarrow reduced basis inherits the **fixed** truth temporal discretization.

C. Prud'homme

Extension to time dependent problems

$$\underbrace{\left(\frac{1}{\Delta t} M_N + A_N\right)}_{N \times N} \underbrace{U_N^k}_N = \underbrace{F_N g(t^k)}_N + \underbrace{\frac{1}{\Delta t} M_N U_N^{k-1}}_{N \times N}$$

- Similar to elliptic case
 - Additional terms due to m
 - Time-dependence: LU-decomposition
- Affine parameter dependence of (bi)linear forms
- Hierarchical RB space
 - Arrays for $N \leq N_{\max}$ are principal subarrays of arrays for $N = N_{\max}$.

We expand $u_N^k(\mu) = \sum_{j=1}^N u_{Nj}^k(\mu) \zeta^j$ and obtain

$$\begin{aligned}
 & a(u_N^k(\mu), v; \mu) + \frac{1}{\Delta t} m(u_N^k(\mu), v; \mu) = \dots \\
 & \sum_{j=1}^N \left[a(\zeta^j, \zeta^i; \mu) + \frac{1}{\Delta t} m(\zeta^j, \zeta^i; \mu) \right] u_{Nj}^k(\mu) = \dots \\
 & \underbrace{\sum_{j=1}^N \left[\sum_{q=1}^{Q_a} \Theta_a^q(\mu) \underbrace{a^q(\zeta^j, \zeta^i)}_{\text{OFFLINE: } O(\mathcal{N})} \right]}_{\text{ONLINE: } O(Q_a N^2)} + \frac{1}{\Delta t} \underbrace{\sum_{q=1}^{Q_m} \Theta_m^q(\mu) \underbrace{m^q(\zeta^j, \zeta^i)}_{\text{OFFLINE: } O(\mathcal{N})}}_{\text{ONLINE: } O(Q_m N^2)} u_{Nj}^k(\mu) = \dots
 \end{aligned}$$

and

$$v = \zeta^i, 1 \leq i \leq N$$

Figure

$$\dots = \frac{1}{\Delta t} m(u_N^{k-1}(\mu), v; \mu) + f(v; \mu) g(t^k)$$

$$\dots = \sum_{j=1}^N \frac{1}{\Delta t} m(\zeta^j, \zeta^i; \mu) u_{Nj}^{k-1}(\mu) + f(\zeta^i; \mu) g(t^k)$$

$$\dots = \sum_{j=1}^N \frac{1}{\Delta t} \underbrace{\sum_m^{Q_m} \Theta_m^q(\mu) \underbrace{m^q(\zeta^j, \zeta^i)}_{\text{OFFLINE: } O(\mathcal{N})} u_{Nj}^{k-1}(\mu)}_{\text{ONLINE: } O(Q_m N^2)} + \underbrace{\sum_{q=1}^{Q_f} \Theta_f^q(\mu) \underbrace{f^q(\zeta^i)}_{\text{OFFLINE: } O(\mathcal{N})} g(t^k)}_{\text{ONLINE: } O(Q_f N)}$$

$$\Rightarrow \left\{ \begin{array}{l} \text{solve for } u_{Nj}^k(\mu), 1 \leq j \leq N, 1 \leq k \leq K. \end{array} \right.$$

$O(N^3 + KN^2)$ \rightarrow resolution
factorisation

Given $u_{Nj}^k(\mu), 1 \leq j \leq N$, evaluate the output from $\forall k \in \mathbb{K}$

$$\begin{aligned} s_N^k(\mu) &= \ell(u_N^k(\mu); \mu) = \sum_{j=1}^N u_{Nj}^k(\mu) \ell(\zeta^j; \mu) \\ &= \sum_{j=1}^N \underbrace{u_{Nj}^k(\mu)}_{\text{ONLINE: } O(N)} \underbrace{\sum_{q=1}^{Q_\ell} \Theta_\ell^q(\mu)}_{\text{ONLINE: } O(Q_\ell N)} \underbrace{\ell^q(\zeta^j)}_{\text{OFFLINE: } O(N)} \\ &\Rightarrow \text{solve for } s_N^k(\mu), 1 \leq k \leq K, \text{ in } O(KN) \end{aligned}$$

Summary computational cost:

$$(Q = Q_a + Q_m)$$

- OFFLINE - once, parameter independent

$$\underbrace{O(KN_{\max}\mathcal{N}^\bullet)}_{\text{solve for } \zeta_n} + \underbrace{O(QN_{\max}^2\mathcal{N})}_{\text{form } \mu\text{-independent quantities}}$$

- ONLINE - many times, parameter dependent

 μ^{new}

$$\underbrace{O(QN^2)}_{\text{form RB matrices}} + \underbrace{O(N^3 + KN^2)}_{\text{solve for } u_{N_j}^k(\mu)} + \underbrace{O(KN)}_{\text{evaluate output}}$$

Online cost is independent of \mathcal{N} .

Evaluation of RB Stiffness Matrix $\underline{A}_N \in \mathbb{R}^{N \times N}$: Parameter-independent matrices $\underline{A}_N^q \in \mathbb{R}^{N \times N}$, $1 \leq q \leq Q_a$:

$$\begin{aligned}\underline{A}_{Nnm}^q &= a^q(\zeta^m, \zeta^n) \\ &= \sum_{i=1}^{\mathcal{N}} \sum_{j=1}^{\mathcal{N}} \zeta_i^m a^q(\varphi_i^{\mathcal{N}}, \varphi_j^{\mathcal{N}}) \zeta_j^n, \quad 1 \leq n, m \leq N,\end{aligned}$$

thus

$$\underline{A}_N^q = \underline{Z}_N^T \underline{A}^{\mathcal{N}q} \underline{Z}_N.$$

We finally assemble

$$\underline{A}_N = \sum_{q=1}^{Q_a} \Theta_a^q(\mu) \underline{A}_N^q.$$

Here, $\underline{Z}_N = [\zeta^1 | \zeta^2 | \dots | \zeta^N] \in \mathbb{R}^{\mathcal{N} \times N}$.

Fig

Evaluation of RB Mass Matrix $\underline{M}_N \in \mathbb{R}^{N \times N}$: Parameter-independent matrices $\underline{M}_N^q \in \mathbb{R}^{N \times N}, 1 \leq q \leq Q_m$:

$$\begin{aligned}\underline{M}_{Nnm}^q &= m^q(\zeta^m, \zeta^n) \\ &= \sum_{i=1}^{\mathcal{N}} \sum_{j=1}^{\mathcal{N}} \zeta_i^m m^q(\varphi_i^{\mathcal{N}}, \varphi_j^{\mathcal{N}}) \zeta_j^n, \quad 1 \leq n, m \leq N\end{aligned}$$

thus

$$\underline{M}_N^q = \underline{Z}_N^T \underline{M}^{\mathcal{N}q} \underline{Z}_N.$$

We finally assemble

$$\underline{M}_N = \sum_{q=1}^{Q_m} \Theta_m^q(\mu) \underline{M}_N^q.$$

Evaluation of RB Load/Source Vector $\underline{F}_N \in \mathbb{R}^N$: Parameter-independent vectors $\mathbb{F}_N^q \in \mathbb{R}^N, 1 \leq q \leq Q_f$:

$$\begin{aligned}\mathbb{F}_{Nn}^q &= f^q(\zeta^n) \\ &= \sum_{i=1}^{\mathcal{N}} \zeta_i^m f^q(\varphi_i^N), \quad 1 \leq n \leq N\end{aligned}$$

thus

$$\underline{\mathbb{F}}_N^q = \mathbb{Z}_N^T \underline{\mathbb{F}}^{\mathcal{N}\Pi}.$$

We finally assemble

$$\underline{F}_N = \sum_{q=1}^{Q_f} \Theta_f^q(\mu) \underline{\mathbb{F}}_N^q.$$

Evaluation of RB Output Vector $\underline{L}_N \in \mathbb{R}^N$: Parameter-independent vectors $\underline{L}_N^q \in \mathbb{R}^N, 1 \leq q \leq Q_\ell$:

$$\begin{aligned}\mathbb{L}_{Nn}^q &= \ell^q(\zeta^n) \\ &= \sum_{i=1}^{\mathcal{N}} \zeta_i^m \ell^q(\varphi_i^N), \quad 1 \leq n \leq N\end{aligned}$$

thus

$$\underline{L}_N^q = \mathbb{Z}_N^T \underline{L}^{\mathcal{N}q}.$$

We finally assemble

$$\underline{L}_N = \sum_{q=1}^{Q_\ell} \Theta_\ell^q(\mu) \underline{L}_N^q.$$

Given $\mu \in \mathcal{D}$, evaluate $\forall k \in \mathbb{K}$

$$s_N^k(\mu) = \underline{L}_N^T(\mu) \underline{u}_N^k(\mu)$$


where $\underline{u}_N^k(\mu) \in \mathbb{R}^N$ satisfies $u_{N,0}(\mu) = 0$

$$\left(\underline{A}_N(\mu) + \frac{1}{\Delta t} \underline{M}_N(\mu) \right) \underline{u}_N^k(\mu) = \frac{1}{\Delta t} \underline{M}_N(\mu) \underline{u}_N^{k-1}(\mu) + \underline{F}_N(\mu) g(t^k).$$

- LU-decomposition: $\underline{A}_N(\mu) + \frac{1}{\Delta t} \underline{M}_N(\mu)$
- Forward/Back Substitution: $\underline{u}_N^k(\mu), \forall k \in \mathbb{K}$

Example: Concrete Delamination - Results i

Fiber reinforced plastic



- Input (parameter) : $\mu \equiv \left(\underbrace{w_{del} / 2}, \kappa \equiv \tilde{k}_{FRP} / \tilde{k}_C \right) \subset \mathcal{D}$ where $\mathcal{D} \equiv [1, 10] \times [0.4, 1.8]$.
- "Truth": $\mathcal{N} = 5601, K = 200$.

Example: Concrete Delamination - Results ii

| N | $\epsilon_{\max, \text{rel}}^u$ | $\epsilon_{\max, \text{rel}}^s$ |
|-----|---------------------------------|---------------------------------|
| 20 | 8.09E - 02 | 6.76E - 01 |
| 40 | 2.71E - 02 | 1.44E - 02 |
| 60 | 1.02E - 02 | 3.34E - 03 |
| 80 | 5.02E - 03 | 1.43E - 03 |
| 120 | 7.40E - 04 | 9.81E - 05 |
| 160 | 2.13E - 04 | 2.34E - 05 |
| 200 | 9.55E - 05 | 6.02E - 06 |

Figure

- Maximum relative error:

$$\epsilon_{\max, \text{rel}}^u = \max_{\mu \in \Xi_{\text{test}}} \frac{\|e^K\|_{\mu}}{\|u^K(\mu)\|}, \quad \mu_u = \arg \max_{\mu \in \Xi_{\text{test}}} \|u^K(\mu)\|$$

- Maximum relative output error:

$$\epsilon_{\max, \text{rel}}^s = \max_{\mu \in \Xi_{\text{test}}} \max_{k \in \mathbb{K}} \frac{|s^k(\mu) - s_N^k(\mu)|}{s_{\max}}, \quad s_{\max} = \max_{\mu \in \Xi_{\text{test}}} \max_{k \in \mathbb{K}} |s^k(\mu)|$$

How do we choose N ?

A posteriori error estimation

- How do we know that $\mathbf{u}_N^k(\mu), s_N^k(\mu)$ are accurate? ONLINE

$$\| \mathbf{u}^k(\mu) - \mathbf{u}_N^k(\mu) \|_{\mu} \leq \epsilon_{\text{tol}, \min}, \quad \forall k \in \mathbb{K}, \forall \mu \in \mathcal{D}$$

$$|s^k(\mu) - s_N^k(\mu)| \leq \epsilon_{\text{tol}, \min}^s, \quad \forall k \in \mathbb{K}, \forall \mu \in \mathcal{D}$$

- How do we know what value of N to take? ONLINE/OFFLINE
 - N too large \Rightarrow computational inefficiency
 - N too small \Rightarrow unacceptable uncertainty

born d'uncert

- How do we choose the sample S_N optimally? OFFLINE
 - RB space has to approximate manifold \mathcal{M} well, but
 - RB matrices need to be "well-conditioned."

Our a posteriori error bounds, $\Delta_N^k(\mu)$ and $\Delta_N^{sk}(\mu)$, must be

- rigorous

$$1 \leq N \leq N_{\max}$$

$$\begin{aligned} |||u^k(\mu) - u_N^k(\mu)||| &\leq \Delta_N^k(\mu), \quad \boxed{\forall k \in \mathbb{K}}, \underbrace{\forall \mu \in \mathcal{D}}, \\ |s^k(\mu) - s_N^k(\mu)| &\leq \Delta_N^{sk}(\mu), \quad \boxed{\forall k \in \mathbb{K}}, \underbrace{\forall \mu \in \mathcal{D}}. \end{aligned}$$

- reasonably sharp

$$\frac{\Delta_N^k(\mu)}{|||u^k(\mu) - u_N^k(\mu)|||} \leq \frac{C}{\varepsilon}, \quad \frac{\Delta_N^{sk}(\mu)}{|s^k(\mu) - s_N^k(\mu)|} \leq \frac{C}{\varepsilon} \quad (5)$$

where $C \approx 1$

- efficient

\Rightarrow Online cost depends on \mathbf{N} , \mathbf{Q} , and \mathbf{K} , but not on \mathcal{N} .

- X -inner product and induced norm (parameter-independent)

$$(w, v)_X \equiv a(w, v; \bar{\mu}), \quad \forall w, v \in X$$

$$\|w\|_X \equiv \sqrt{(w, w)_X}, \quad \forall w \in X$$

- L^2 -inner product and induced norm (parameter-independent)

$$(w, v) \equiv m(w, v; \bar{\mu}), \quad \forall w, v \in X$$

$$\|w\| \equiv \sqrt{(w, w)}, \quad \forall w \in X$$

Milane

- "Spatio-temporal" energy norm (parameter-dependent) $1 \leq k \leq K$

$$\left(\left((w^k, v^k) \right) \right) = m(w^k, v^k; \mu) + \sum_{k'=1}^k \Delta t a(w^{k'}, v^{k'}; \mu)$$

$$\left(\|w^k\| \right) = \left(m(w^k, w^k; \mu) + \sum_{k'=1}^k \Delta t a(w^{k'}, w^{k'}; \mu) \right)^{1/2}$$

We also define

- - Coercivity constants

$$\alpha(\mu) \equiv \inf_{w \in X} \frac{a(w, w; \mu)}{\|w\|_X^2}; \quad \sigma(\mu) \equiv \inf_{w \in X} \frac{m(w, w; \mu)}{\|w\|^2};$$

- Continuity constants

$$\gamma_a(\mu) \equiv \sup_{w \in X} \sup_{v \in X} \frac{a(w, v; \mu)}{\|w\|_X \|v\|_X}.$$
$$\gamma_m(\mu) \equiv \sup_{w \in X} \sup_{v \in X} \frac{m(w, v; \mu)}{\|w\| \|v\|}.$$

Preliminaries: Coercivity lower bound

Figur

We require a positive lower bound for the coercivity constant

$$\alpha_{\text{LB}} : \mathcal{D} \rightarrow \mathbb{R}$$

$$0 < \alpha_{\text{LB}}(\mu) \leq \mu(\mu), \quad \forall \mu \in \mathcal{D}$$

$$\sigma_{\text{LB}} : \mathcal{D} \rightarrow \mathbb{R}$$

$$0 < \sigma_{\text{LB}}(\mu) \leq \mu(\mu), \quad \forall \mu \in \mathcal{D}$$

This bound can be calculated using the

- "min Θ " Approach (if a is parametrically coercive), or
- Successive Constraint Method

exactly as in elliptic case.

Prerequisites: Dual Norm of Residual

We define the residual, $\forall \mathbf{k} \in \mathbb{K}$

$$r^k(v; \mu) \equiv f(v; \mu)g(t^k) - m \left(\frac{u_N(t^k; \mu) - u_N(t^{k-1}; \mu)}{\Delta t}, v; \mu \right) - a(u_N(t^k; \mu), v; \mu), \quad \forall v \in X$$

Dual norm of Residual

Given $\mu \in \mathcal{D}$, the dual norm of $r^k(v; \mu)$ is defined as

$$\begin{aligned} \|r^k(\cdot; \mu)\|_{X'} &\equiv \sup_{v \in X} \frac{r^k(v; \mu)}{\|v\|_X} \\ &= \|\hat{e}^k(\mu)\|_X, \end{aligned}$$

where $\hat{e}^k(\mu) \in X$ satisfies

$$(\hat{e}^k(\mu), v)_X = r^k(v; \mu), \quad \forall v \in X$$

We define the error bound, $\Delta_N^k(\mu) = \Delta_N(t^k; \mu)$, $1 \leq k \leq K$, as

$$\Delta_N^k(\mu) = \alpha_{\text{LB}}^{-1/2}(\mu) \left(\sum_{k'=1}^k \Delta t \left\| \hat{e}^{k'}(\mu) \right\|_X^2 \right)^{1/2}.$$

We can then prove

Proposition (Energy Error Bound)

For any $N = 1, \dots, N_{\max}$, the error in the field variable,

$e^k(\mu) = u^k(\mu) - u_N^k(\mu)$, is bounded by

$$||| e^k(\mu) ||| < \Delta_N^k(\mu), \quad \forall \mu \in \mathcal{D}, \forall k \in \mathbb{K}.$$

We define the output error bound, $\Delta_N^{sk}(\mu) = \Delta_N^s(t^k; \mu)$ $1 \leq k \leq K$, as

$$\Delta_N^{sk}(\mu) \equiv \sigma_{\text{LB}}^{-1}(\mu) \left(\sup_{v \in X} \frac{\ell(v; \mu)}{\|v\|} \right) \Delta_N^k(\mu)$$

Proposition (Simple Output Error Bound)

For any $N = 1, \dots, N_{\max}$, the error in the output is bounded by

$$|s^k(\mu) - s_N^k(\mu)| \leq \Delta_N^{sk}(\mu), \quad \forall \mu \in \mathcal{D}, \forall k \in \mathbb{K}$$

Remarks

- The error bounds are rigorous upper bounds for the reduced basis error for any $N = 1, \dots, N_{\max}$, for all $\mu \in \mathcal{D}$, and for all $k \in \mathbb{K}$.
- Define: $s_N^{\pm}(t^k; \mu) = s_N(t^k; \mu) \pm \Delta^s(t^k; \mu)$, then

$$\Rightarrow s_N^{-}(t^k; \mu) \leq s(t^k; \mu) \leq s_N^{+}(t^k; \mu)$$

- We may also consider other norms than $||| \cdot |||_{\mu}$, i.e., $L^2(\Omega)$
- Results for energy norm and output bound directly extend to nonsymmetric problems
 - if we choose an appropriate definition for the energy norm

Crucial ingredient: Dual norm of residual $\|\hat{e}^k(\mu)\|_X, \forall k \in \mathbb{K}$.

Computational procedure follows directly from the elliptic case with added complexity due to mass term and time dependence.

- Expand $u_N(\mu) = \sum_{j=1}^N u_{Nj}^k(\mu) \zeta^j$
- Riesz representation:

$$(\hat{e}^k(\mu), v)_X = r^k(v; \mu)$$

$$\|\hat{e}^k\|_X = \|a^k\|_X$$

- Affine decomposition
- Linear superposition

Summary of computational cost:

$$Q = Q_a + Q_m$$

OFFLINE -

$$\begin{array}{ll} O(QN_{\max}\mathcal{N}^\bullet) + & O(Q^2N_{\max}^2\mathcal{N}) \\ \text{solve Poisson problems} & \text{form } \mu\text{-independent inner products} \end{array} ;$$

ONLINE -

$$\begin{array}{l} O(KQ^2N^2) \\ \text{evaluate } \|\hat{e}^k(\mu)\|_X \text{-sum for } 1 \leq k \leq K \end{array}$$

Online cost is independent of \mathcal{N} .

Example: Concrete Delamination - Results i

| N | $\epsilon_{\max, \text{rel}}^u$ | $\Delta_{\max, \text{rel}}^u$ | $\bar{\eta}^u$ |
|-----|---------------------------------|-------------------------------|----------------|
| 20 | 8.09E - 02 | 3.18E - 01 | 2.74 |
| 40 | 2.71E - 02 | 8.01E - 02 | 2.77 |
| 60 | 1.02E - 02 | 2.01E - 02 | 2.58 |
| 80 | 5.02E - 03 | 8.40E - 03 | 2.83 |
| 120 | 7.40E - 04 | 1.71E - 03 | 2.45 |
| 160 | 2.13E - 04 | 4.84E - 04 | 2.21 |
| 200 | 9.55E - 05 | 2.70E - 04 | 2.20 |

Table 1: Convergence energy norm error and bound

- Maximum relative error bound:

$$\Delta_{\max, \text{rel}}^y = \max_{\mu \in \Xi_{\text{test}}} \frac{\Delta_N^\alpha(\mu)}{\|u^K(\mu)\|}, \quad \mu_u = \arg \max_{\mu \in \Xi_{\text{test}}} \|u^K(\mu)\|$$

- Average effectivity:

$$\left(\bar{\eta}^u = \frac{1}{n_{\text{train}} K} \sum_{\mu \in \Xi_{\text{test}}} \sum_{k \in \mathbb{K}} \frac{\Delta_N^k(\mu)}{\|e^k(\mu)\|} \right)$$

Example: Concrete Delamination - Results ii

| N | $\epsilon_{\max}^s, \text{rel}$ | $\Delta_{\max}^s, \text{rel}$ | $\bar{\eta}^s$ |
|-----|---------------------------------|-------------------------------|----------------|
| 20 | 6.76E - 02 | 2.58E + 01 | 211 |
| 40 | 1.44E - 02 | 6.24E + 00 | 341 |
| 60 | 3.34E - 03 | 1.46E + 00 | 363 |
| 80 | 1.43E - 03 | 4.73E - 01 | 379 |
| 120 | 9.81E - 05 | 1.24E - 01 | 604 |
| 160 | 2.34E - 05 | 2.88E - 02 | 674 |
| 200 | 6.02E - 06 | 9.18E - 03 | 1117 |

Table 2: Convergence output error and bound

- Maximum relative output bound:

$$\Delta_{\max, \text{rel}}^s = \max_{\mu \in \Xi_{\text{test}}} \frac{\Delta_N^{sK}(\mu)}{|s_{\max}|}$$

- Average output effectivity:

$$\bar{\eta}^s = \frac{1}{n_{\text{train}}} \sum_{\mu \in \Xi_{\text{test}}} \frac{\Delta_N^{s k_{\eta}(\mu)}(\mu)}{|s_{\eta}^{k_{\eta}(\mu)}(\mu) - s_N^{k_{\eta}(\mu)}(\mu)|}, \quad k_{\eta}(\mu) = \arg \max_{k \in \mathbb{K}} |s^k(\mu) - s_N^k(\mu)|$$

The notion "compliance" does not exist in the parabolic context.

Thus similar to the noncompliant elliptic problem, we consider a primal-dual formulation for the parabolic problem

Goal:

- - Faster convergence of output error & bound.
output error = primal error (N_{pr}) \times dual error (N_{du})
- Improved effectivities for output error estimation.

Sampling Strategy

We extend the Greedy Algorithm to a POD(t)-Greedy (μ) sampling procedure, combining a

- small POD in time, with \Rightarrow optimally captures causality of time variation
- (exhaustive) Greedy search in parameter space \mathcal{D} . \Rightarrow (sub-)optimal selection for high-dimensional \mathcal{D} (large n_{train}).

We define

- Desired error tolerance $\varepsilon_{\text{tol},\min}$.
- Train sample $\Xi_{\text{train}} \equiv \{\mu_{\text{train}}^1, \dots, \mu_{\text{train}}^{n_{\text{train}}}\} \subset \mathcal{D}$, with
- Cardinality (size) $|\Xi_{\text{train}}| = n_{\text{train}}$. $\Rightarrow \Xi_{\text{train}}$ serves as our (finite) surrogate for \mathcal{D} .

Proper Orthogonal Decomposition (POD) in time:

- Let

$$\text{POD}_X(\{\underbrace{u^k(\mu), 1 \leq k \leq K}_{\text{Ric}}, R\})$$

return the R largest POD modes, $\{\Psi^{\text{POD},i}, 1 \leq i \leq R\}$, with respect to the $(\cdot, \cdot)_X$ inner product.

- The set $\mathcal{P}_R = \{\Psi^{\text{POD},i}, 1 \leq i \leq R\}$ is $(\cdot, \cdot)_X$ orthogonal and satisfies the optimality property

$$\mathcal{P}_R = \arg \inf_{X_R \subset \text{span}\{u^k(\mu), 1 \leq k \leq K\}} \left(\frac{1}{K} \sum_{k=1}^K \inf_{v \in X_R} \|u^k(\mu) - v\|_X^2 \right)^{1/2}$$

Evaluation of $\psi^{\text{POD},1} = \text{POD}_X(\{u^k(\mu), 1 \leq k \leq K\})$ (1):

Figur

- 1 Form correlation matrix $\underline{C}^{\text{POD}} \in \mathbb{R}^{K \times K}$ given by POD

$$C_{ij}^{\text{POD}} = \left(\frac{1}{K} \right) (u^i(\mu), u^j(\mu))_X, \quad 1 \leq i, j \leq K.$$

- 2 Solve for eigenpair $(\underline{\psi}^{\text{POD},\max} \in \mathbb{R}^K, \lambda^{\text{POD},\max} \in \mathbb{R}_{+0})$,

$$\underline{C}^{\text{POD}} \underline{\psi}^{\text{POD},k} = \lambda^{\text{POD},k} \underline{\psi}^{\text{POD},k}.$$

- 3 Compute largest POD mode

$$\underline{\psi}^{\text{POD},1} \equiv \sum_{k=1}^K \psi_k^{\text{POD},\max} u^k(\mu)$$

Remarks

- Perform POD on projection error instead of data, i.e.

$$\text{POD}_X \left(\{ e_{N,\text{proj}}^k(\mu), 1 \leq k \leq K \}, 1 \right)$$

where $e_{N,\text{proj}}^k(\mu) = u^k(\mu) - \text{proj}_{X, X_N} u^k(\mu)$, and $\text{proj}_{X, X_N} u^k(\mu)$ is the X -orthogonal projection of $u^k(\mu)$ onto \mathbf{X}_N .

- Algebraic notation

$$\underline{e}_{N,\text{proj}}^k(\mu) = \underline{u}^k(\mu) - \mathbb{Z}_N \left(\mathbb{Z}_N^T \mathbb{X}^N \underline{u}^k(\mu) \right)$$

- In general, we set $R = 1$ and add only one mode at each iteration (offline vs. online effort).

We require a rigorous, sharp, inexpensive error bound:

$$\|u^k(\mu) - u_N^k(\mu)\| \leq \Delta_N^k(\mu), \quad \forall \mu \in \mathcal{D}.$$

Note (see below)

- Effectivities $\bar{\eta}^u$ are $\mathcal{O}(1)$
- Computational cost to evaluate $\Delta_N^k(\mu)$ is $\mathcal{O}(KQ^2N^2)$.

Use Greedy (μ) Idea:

- $\Delta_N^k(\mu)$ is monotonically increasing in time. Find parameter value such that

$$\mu^* = \arg \max_{\mu \in \Xi_{\text{train}}} \Delta_N^K(\mu)$$

\Rightarrow Largest error bound at **final time**.

Figur

Greedy, $L^\infty(\Xi_{\text{train}}, \|\cdot\|)$, space "economization"

$$\boxed{Kn_{\text{train}}} \text{ contestants} \Rightarrow \underbrace{N_{\text{max}}}_{\mu_1^*, \dots, \mu_{N_{\text{max}}}^*} (\ll Kn_{\text{train}}) \text{ winners}$$

$\in \Xi_{\text{train}} \times \mathbb{I}$

in which we never form most snapshots:

$$\|u^k(\mu) - u_N^k(\mu)\|_{n_{\text{train}}} \cdot O(KN^\bullet) \quad \text{replaced by} \quad \frac{\Delta_N^k(\mu)}{n_{\text{train}}} \cdot O(KQ^2N^2)^1$$

note good effectivity of estimator is crucial.

POD(t)-Greedy (μ) Algorithm

Set $X_N = \{0\}, S_N = \{0\}, N = 0, \mu^* = \mu_0^*$

while $\Delta_N^{\max} \geq \varepsilon_{\text{tol}, \min}$

$$e_{N, \text{proj}}^k(\mu^*) = u^k(\mu^*) - \text{proj}_{X, X_N} u^k(\mu^*), 1 \leq k \leq K$$

$$S_{N+1} = S_N \cup \mu^*;$$

$$X_{N+1} = X_N + \text{POD}_X(\{e_{N, \text{proj}}^k(\mu^*), 1 \leq k \leq K\}, 1)$$

$$N = N + 1;$$

$$\mu^* = \arg \max_{\mu \in \Xi_{\text{train}}} \Delta_N^K(\mu) / \|u_N^K(\mu)\|;$$

$$\Delta_N^{\max} = \Delta_N^K(\mu^*) / \|y_N^K(\mu^*)\|;$$

end while

Remarks

Figure

- Spaces X_N are hierarchical.
- Algorithm guarantees that $\|u^k(\mu) - u_N^k(\mu)\| \leq \Delta_N^k(\mu) \leq \varepsilon_{\text{tol}, \min}, \forall \mu \in \Xi_{\text{train}}$.
- We can replace condition on Δ_N^{\max} by a condition on N_{\max} (hp-Reduced Basis).
- No additional Gram-Schmidt orthogonalization required, basis functions are "by construction" \mathbf{X} -orthogonal.
- Computational complexity remains $O(K\mathcal{N}^\bullet) + O(n_{\text{train}})$ not $O(K\mathcal{N}^\bullet n_{\text{train}})$

¹In addition to the offline effort that is required in any event for online rigorous/sharp certification.

- Nonzero initial conditions, $u_0(\mu) \neq 0$.
 - Nonzero (but constant) initial condition

$$\Rightarrow \zeta^1 = u_0(\mu) \neq 0.$$

- Affinely parameter dependent initial condition

$$\underbrace{u_0(\mu)} = \sum_{q=1}^{Q_{u_0}} \Theta_{u_0}^q(\mu) \underbrace{u_0^q}_{\in X}$$

where $u_0^q \in X$, μ -independent and known, and $\Theta_{u_0}^q : \mathcal{D} \rightarrow \mathbb{R}$, μ -dependent functions. We then initialize

$$\Rightarrow \underline{X_N} = \text{span} \{ \underbrace{u_0^q}_{\neq 0}, 1 \leq q \leq \underbrace{Q_{u_0}}_{\neq 0} \}.$$

- No a priori knowledge
 - Series representation of u_0 ;
 - Projection of u_0 onto $X_N(\mathcal{N}$ -dependent cost);
 - Contribution to error & bound.

Handwritten notes:

$$\left. \begin{array}{l} \text{at } t=0 \\ u(t=0) = u_0 \\ u_0 \end{array} \right\}$$

Figure

- Unknown "control" input, $g(t^k)$ (e.g. optimal control). **Duhamel's Principle:** given any control input $g(t^k)$, we can obtain $u^k(\mu)$ from

$$u^k(\mu) = \sum_{j=1}^K h(t^{k-j+1}; \mu) g(t^j), \quad \forall k \in \mathbb{K},$$

where $h(t^k; \mu)$ is the impulse response. We thus train the RB approximation on an **impulse input**

$$\Rightarrow \underline{g(t^k)} = \underline{\delta_{1k}}, \forall k \in \mathbb{K}. \quad \text{linear time invariant}$$

only valid for LTI systems

- Multiple "control" inputs, $g(t^k) \in \mathbb{R}^m$
 \Rightarrow recursive training on each input (LTI).

Temporary page!

\LaTeX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because \LaTeX now knows how many pages to expect for this document.