44. BIRTH

DURATION : 04:46 STUDY SHEET

COURSE SUMMARY

This video explores the profound impacts of birth on our lives and bodies, while offering practical methods to release tensions stored in bodily tissues. Key concepts include the importance of compression and decompression at the cranial, pelvic, and vascular levels, and how this affects the autonomous respiration of newborns. The speaker also highlights the central role of the spleen and oxygen transfer during the first breath, as well as the importance of fluid movement between the parietal and temporal bones to facilitate this crucial transition. Students will learn to better understand the link between bodily movements, spiritual dynamics, and the osteopathic approach, while integrating these principles into their therapeutic practice.

BIRTH: IMPACT AND MECHANISMS

The way you were **born** determines many aspects of your life. Every time you have to **be reborn** into something, like a job change, you can reconnect with that memory. It is sometimes necessary to release tensions in the bodily **tissue**, either by recompressing or decompressing it, in order to re-establish **respiratory disengagement**.

It is not enough to bring these memories to consciousness; they must be reborn in the tissue. In this approach, I will compress from the **parietals**. For a baby, if something seems necessary, I will work at the level of the **pelvis**. The child may need to be compressed or decompressed, and it is the feeling of the moment that guides this decision.

This process leads to a **spiritual** movement and dynamics in the body. At the **osseous-membranous** level, it is also crucial to consider compression at the **vascular** level. Imagine a sponge being compressed; this illustrates well what happens in the body.

The **spleen** plays an essential role during birth. If necessary, it will re-establish a flow, because a **rebalancing** is necessary throughout the vascular system, between the heart, the lungs, and the **umbilical cord**. This passage is fundamental, as it represents a major vascular change: you go from breathing and oxygenation via the cord to respiratory autonomy via your lungs.

The first **cry**, the first **breath**, is crucial. At that moment, you are in an umbilical axis, and blood enters through the **foramen ovale**. This is a major fluidic change. To facilitate this change, the **hepatic** and **splenic axis** intervenes, providing the necessary information. If this transition is not

well performed, some children may be born with **splenic** characteristics, often melancholic, because they did not perform this first breath correctly.

This passage of oxygen, of the global and fluidic compression of the system, as well as membranous compression, is essential. The importance of the **parietals** and the movement of the body should be emphasised.

The fluid movement between **parietal** and **temporal** is a first disengagement. This movement leads to a rotation of the **neck**, essential for releasing the body. The key moment to give the first breath is marked by a slight movement of the neck, followed by a spiral that forms.

This occipital movement, associated with a torsion in the axis, is crucial. Hyperextension of the neck and disengagement of the limbs are also elements to consider. Thus, there is compression in the membranes and compression at the fluidic level that occur simultaneously.