

45. DEMO: VENTRICULAR SYSTEM

DURATION : 03:45

STUDY SHEET

COURSE SUMMARY

This video offers a practical demonstration of the biodynamic approach applied to the ventricular system, focusing on tactile perception and progression through different tissue layers. The instructor guides therapists through a process of body exploration, starting with a delicate contact with the surface of the head and descending to the bony structures, while cultivating an increased sensitivity to changes in texture and temperature. Key concepts discussed include oscillatory interaction in the cerebral hemispheres, as well as the creation of a therapeutic space that promotes better breathing and body engagement. This practice proves enriching for understanding the body's internal and evolutionary dynamics.

DEMONSTRATION: VENTRICULAR SYSTEM

CONTACT AND INITIAL PERCEPTION

I begin by positioning myself from the **periphery** to the **depth**, my thumbs spontaneously orienting along the axis of the **ventricles**. My first point of contact is with the **texture of the hair**.

Without excessive movement, I wait to feel the **substance** of the head and the **warmth** that begins to emanate from my hands.

This is a particular moment where one perceives a **difference**, a clear emergence. Suddenly, one feels the **bone**. My hands, rich in **receptors**, become active.

PROGRESSION THROUGH TISSUE LAYERS

I will use the same **progressive** approach. My hands emit warmth, and I descend delicately, layer by layer: the hair, the skin, the **fascia**, then the bone.

I am still on the bone. But now, the perception changes; it begins to "pass through" to the left. I gently enter into this sensation, this perception, this **frequency**.

The bone no longer seems a **rigid** structure. It becomes like **fluid**, allowing for deeper penetration. It is important to remember that we are passing through **mesoderm** here, a mesodermal tissue that extends to the **brain**.

REACHING THE DEEP SPHERES

I wait for the body to manage a subtle **oscillatory** behaviour. I begin to feel it. I take a step back to better appreciate the **depth**.

We are at the superficial layer, the **cortex**, the shadow of the **arachnoid** with its arachnoid pillars. At this level, I begin to perceive the **hemispheres**.

We will go a little further, playing with the **ventricles**. We will feel the brain as a whole.

THE THERAPEUTIC INTERACTION

She feels a new sensation. I will induce a slight movement, like a **dolphin**, on the right side which is breathing a little less well.

I simply offer her **space** to allow her to breathe better. Then, I slowly withdraw, layer by layer.

RECIPIENT'S PERCEPTION

Here is a description of the perceived sensation:

- **At the bone level:** An initial impact sensation, then great lightness.
- **Below the bone:** The movement of the hemispheres was very pleasant. A clear distinction was perceptible between the left and right, particularly at the level of the ventricles.
- **Finally:** A more **electrical** frequency, a very rapid but not sharp vibration.