

13. PRIMITIVE CRANIO-SACRAL AXIS

DURATION : 03:34

STUDY SHEET

COURSE SUMMARY

In this video, we delve deeply into the primitive cranio-sacral axis, which emerges during the second week of embryonic development with the formation of the notochord. The teaching highlights the importance of this structure for the cranium and sacrum, particularly by emphasising how a free sacrum is essential for ensuring spinal functionality and flexion-extension of the cranial base. Furthermore, the video discusses the implications these embryonic dynamics have on urogenital pathologies, and how the release of certain emotional or energetic blockages between the sigmoid colon and the bladder can contribute to better overall balance. Students will learn to manipulate these concepts to improve their practice in osteopathy and biodynamic embryology.

PRIMITIVE CRANIO-SACRAL AXIS

During the second week of embryonic development, the **S-shape** of the epiblastic process begins to form the **notochord**. The **Hensen's node**, or primitive pit, becomes a fixed fulcrum and progressively moves posteriorly.




FIG. — *Beginning of the epiblastic process*

This structure is essential as it forms the base of the **cranium**, integrating elements such as the **pituitary gland**, the **greater wings of the sphenoid**, the **petrous system**, the **occipital system**, and the **ethmoid**. Inferiorly, the development of the **sacrum** begins, with the S2 process representing the vestige of the primitive streak. This process becomes a crucial fulcrum, highlighting the importance of a **free sacrum** to allow for functional unity.

In this developmental phase, the embryo takes on a racket-like shape, with the notochord and the primitive streak. This growth process generates a suction field to form the **mesoderm**, particularly the precardiac field. The anterior growth of the embryo draws material into this area, pulling **bottle cells** towards the centre. These cells release **hyaluronic acid**, which is essential at the mesenchymal level.

The notochordal axis is fundamental because it influences the freedom of movement of the sacrum. If the sacrum is blocked, the spine cannot function correctly, which affects the base of the cranium. A non-free cranial base prevents proper **flexion-extension**, leading to an imbalance between the **neurocranium** and the **viscerocranium**.

This dynamic is linked to the ascending force of the cranium, where the eye settles, and to the establishment of **mesodermal tissue**. Ectodermal growth pulls upwards, while the primitive node descends towards the sacrum. This creates an area of importance for protein information in

relation to the mother.

Many blocked **ancestral images** are located between the sigmoid colon and the bladder, where it is crucial to release this information. Urogenital problems can be transmitted by this information, and the sacral downward migratory force plays a key role in this process.