

31. PRACTICE: PINEAL AND PITUITARY GLANDS

DURATION : 10:39

STUDY SHEET

COURSE SUMMARY

This video offers an in-depth exploration of the fundamental roles of the pineal and pituitary glands in hormonal regulation and memory. It presents energy rebalancing techniques inspired by the embryonic journey, specifically concerning pituitary motility. Students will learn to identify the interactions between divine and current light, as well as to use clinical observations such as breathing and eye movements to restore balance between these two crucial glands. The course also highlights the importance of the pineal gland in regulating vital functions such as breathing, digestion, and sleep.

PRACTICE: EPIPHYSIS AND HYPOPHYSIS

The epiphysis and hypophysis play a crucial role in **hormonal regulation** and **memory**. The epiphysis, often called the pineal gland, is the **link between neurons** and the hormonal system. It is influenced by light and contains microorganisms that convert **serotonin** into **melatonin**. When light intensity decreases, melatonin production begins.

To rebalance the hypophysis and epiphysis, an energetic rebalancing technique is used, by revisiting the **embryonic journey**. This involves exploring the motility of the hypophysis through the entry points located at **nasion** and **bregma**.

At the time of the formation of the base of the skull, a space called **Rathke's pouch** forms. This invagination of the epiblast epithelium gives rise to the hypophysis, which receives both descending and ascending information. The hypophysis divides into anterior and posterior hypophysis, and its development is linked to the formation of the **prosencephalon**, which divides into **telencephalic** and **diencephalic** parts.

The epiphysis gland moves from front to back, influenced by divine light and the hypophysis. This interaction between divine light and current light is essential for the function of the hypophysis.



FIG. — *Skull base formation*

The hypophysis develops from the surface epithelium, and the flexion movement causes a suction towards the centre, thus forming the anterior hypophysis gland. In parallel, a suction movement in the brain gives rise to the infundibulum, which constitutes the posterior part of the hypophysis.



FIG. — *Pituitary development*

In practice, it is possible to use internal techniques to work on the energetic axis between the hypophysis and the epiphysis. This involves observing the patient's breathing and eye movements, as these elements are indicative of the energetic state.




FIG. — *Infundibulum development*

The **nervous system**, including the hypothalamus and thalamus, plays a key role in this dynamic. By restarting embryonic motility, it is possible to rebalance the anterior and posterior hypophysis. Eye movements can indicate a return to the hara, thus promoting stability.

The epiphysis is essential for the regulation of **respiration**, **digestion**, and **sleep**. It produces melatonin, a hormone that influences brain waves, thus facilitating falling asleep, deep sleep, and dreams. Although the hypophysis controls almost all glands, the epiphysis is considered the master gland, playing a central role in the body's hormonal and energetic balance.