Assignment 1: Aidan Power, Fiona Price, and Emily Burghart

2024-09-18

Setting up the packages and data

Loading in the packages:

```
library(tidyverse)
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
## v dplyr 1.1.4
                      v readr
                                  2.1.5
## v forcats 1.0.0 v stringr 1.5.1
## v ggplot2 3.5.1 v tibble
                                  3.2.1
## v lubridate 1.9.3
                    v tidyr
                                  1.3.1
## v purrr
             1.0.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag() masks stats::lag()
## i Use the conflicted package (<a href="http://conflicted.r-lib.org/">http://conflicted.r-lib.org/</a>) to force all conflicts to become error
library(ggplot2)
library(here)
```

here() starts at /home/guest/Hydrology/Assignment 1

```
library(dplyr)
```

Now, we will load in the Eno data and make separate dataframes for the median outflow, mean outflow, and volume.

```
#Eno data:
eno_outflows <- read.csv(</pre>
 file = here("./Data Raw/Eno_River_monthly_flow_data_through-2022.csv"),
 stringsAsFactors = TRUE)
#Average of median outflow:
eno_average_median <- eno_outflows %>%
  group_by(year) %>%
  summarise(avg_median_flow = mean(median_flow, na.rm = TRUE))
#Average of mean outflow:
eno_average_mean <- eno_outflows %>%
  group_by(year) %>%
  summarise(avg_mean_flow = mean(mean_flow, na.rm = TRUE))
#Average of volume:
eno_volume_mean <- eno_outflows %>%
  group_by(year) %>%
  summarise(avg_vol = mean(total_volume, na.rm = TRUE))
```

Now, we will load in the Flat data and make separate dataframes for the median outflow, mean outflow, and volume.

```
#Flats data:
flat_outflows <- read.csv(</pre>
  file = here("./Data Raw/Flat_River_monthly_flow_data_through-2022.csv"),
  stringsAsFactors = TRUE)
#Average of median outflow:
flat_average_median <- flat_outflows %>%
  group_by(year) %>%
  summarise(avg_median_flow = mean(median_flow, na.rm = TRUE))
#Average of mean outflow:
flat_average_mean <- flat_outflows %>%
  group_by(year) %>%
  summarise(avg_mean_flow = mean(mean_flow, na.rm = TRUE))
#Average of volume:
flat_volume_mean <- flat_outflows %>%
  group_by(year) %>%
  summarise(avg vol = mean(total volume, na.rm = TRUE))
```

Now, we will load in the Neuse data and make separate dataframes for the median inflow, mean inflow, and volume.


```
#Neuse data:
neuse inflows <- read.csv(</pre>
 file = here("./Data Raw/Falls_Lake_outlet_Neuse_River_monthly_flow_data_through-2022.csv"),
  stringsAsFactors = TRUE)
#Average of median inflow:
neuse_average_median <- neuse_inflows %>%
  group_by(year) %>%
  summarise(avg_median_flow = mean(median_flow, na.rm = TRUE))
#Average of mean inflow:
neuse_average_mean <- neuse_inflows %>%
  group_by(year) %>%
  summarise(avg_mean_flow = mean(mean_flow, na.rm = TRUE))
#Average of volume:
neuse_volume_mean <- neuse_inflows %>%
  group_by(year) %>%
  summarise(avg_vol = mean(total_volume, na.rm = TRUE))
```

Plots

Now, we will start making graphs. First, we will look at Eno. We will start with all years.


```
#Median Eno outflow
ggplot(data = (eno_outflows)) +
  geom_col(data = eno_outflows, aes(x = year, y = median_flow)) +
  labs(title = "Median Eno Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Eno Outflow




```
ggplot(data = eno_average_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Eno Outflow", x = "Year", y = "Median Inflow (in Acre-Feet)")
```

Median Eno Outflow




```
#Mean Eno outflow
ggplot(data = eno_outflows) +
  geom_col(data = eno_outflows, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Eno Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Eno Outflow




```
ggplot(data = eno_average_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Eno Outflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Eno Outflow




```
#Mean volume
ggplot(data = eno_volume_mean) +
  geom_col(data = eno_volume_mean, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Eno Volume



```
ggplot(data = eno_volume_mean, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Eno Volume



There is a gap in Eno data from 1971-1985. Now, we will look only at the post-1985 data.


```
#Filter out the data to only include 1985 and beyond.
eno_filtered <- eno_outflows %>%
 filter(year >= 1985)
#Filter out the average median outflow data to only include 1985 and beyond.
eno_avemedian_filtered <- eno_average_median %>%
 filter(year >= 1985)
#Filter out the average mean outflow data to only include 1985 and beyond.
eno_avemean_filtered <- eno_average_mean %>%
 filter(year >= 1985)
#Filter out the average volume data to only include 1985 and beyond.
eno_vol_filtered <- eno_volume_mean %>%
 filter(year >= 1985)
#Median Eno outflow
ggplot(data = (eno_filtered)) +
  geom_col(data = eno_filtered, aes(x = year, y = median_flow)) +
 labs(title = "Median Eno Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Eno Outflow




```
ggplot(data = eno_avemedian_filtered, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Eno Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Eno Outflow




```
#Mean Eno outflow
ggplot(data = eno_filtered) +
  geom_col(data = eno_filtered, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Eno Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Eno Outflow




```
ggplot(data = eno_avemean_filtered, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Eno Outflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Eno Outflow




```
#Mean Eno volume
ggplot(data = eno_vol_filtered) +
  geom_col(data = eno_vol_filtered, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Eno Volume



```
ggplot(data = eno_vol_filtered, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Eno Volume



Lastly, we will look at only years 2000 and beyond to capture recent trends.


```
#Filter out the data to only include 2000 and beyond.
eno_2000 <- eno_outflows %>%
 filter(year >= 2000)
#Filter out the average median outflow data to only include 2000 and beyond.
eno_2000_median <- eno_average_median %>%
 filter(year >= 2000)
#Filter out the average mean outflow data to only include 2000 and beyond.
eno_2000_mean <- eno_average_mean %>%
 filter(year >= 2000)
#Filter out the average volume data to only include 2000 and beyond.
eno_2000_volume <- eno_volume_mean %>%
 filter(year >= 2000)
#Median Eno outflow
ggplot(data = eno_2000) +
 geom_col(data = eno_2000, aes(x = year, y = median_flow)) +
 labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
ggplot(data = eno_2000_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
#Mean Eno outflow
ggplot(data = eno_2000) +
  geom_col(data = eno_2000, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
ggplot(data = eno_2000_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
#Mean Eno volume
ggplot(data = eno_2000_volume) +
  geom_col(data = eno_2000_volume, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Eno Volume



```
ggplot(data = eno_2000_volume, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Eno Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Eno Volume



Now, we will look at Flat data.


```
#Median Flat outflow
ggplot(data = (flat_outflows)) +
  geom_col(data = flat_outflows, aes(x = year, y = median_flow)) +
  labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
ggplot(data = flat_average_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
#Mean Flat outflow
ggplot(data = (flat_outflows)) +
  geom_col(data = flat_outflows, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
ggplot(data = flat_average_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
#Mean volume
ggplot(data = flat_volume_mean) +
  geom_col(data = flat_volume_mean, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Flat Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Flat Volume



```
ggplot(data = flat_volume_mean, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Flat Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Flat Volume



Now we will look at only years 2000 and beyond to capture recent trends.


```
#Filter out the data to only include 2000 and beyond.
flat_2000 <- flat_outflows %>%
 filter(year >= 2000)
#Filter out the average median outflow data to only include 2000 and beyond.
flat_2000_median <- flat_average_median %>%
 filter(year >= 2000)
#Filter out the average mean outflow data to only include 2000 and beyond.
flat_2000_mean <- flat_average_mean %>%
 filter(year >= 2000)
#Filter out the average volume data to only include 2000 and beyond.
flat_2000_volume <- flat_volume_mean %>%
  filter(year >= 2000)
\#Median\ Flat\ outflow
ggplot(data = flat_2000) +
  geom_col(data = flat_2000, aes(x = year, y = median_flow)) +
 labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
ggplot(data = flat_2000_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Flat Outflow", x = "Year", y = "Median Outflow (in Acre-Feet)")
```

Median Flat Outflow




```
#Mean Flat outflow
ggplot(data = flat_2000) +
  geom_col(data = flat_2000, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
ggplot(data = flat_2000_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Flat Outflow", x = "Year", y = "Mean Outflow (in Acre-Feet)")
```

Mean Flat Outflow




```
#Mean Flat volume
ggplot(data = flat_2000_volume) +
  geom_col(data = flat_2000_volume, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Flat Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Flat Volume



```
ggplot(data = flat_2000_volume, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Flat Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Flat Volume



Finally, we will look at Neuse inflows.


```
#Median Neuse inflow
ggplot(data = (neuse_inflows)) +
  geom_col(data = neuse_inflows, aes(x = year, y = median_flow)) +
  labs(title = "Median Neuse Inflow", x = "Year", y = "Median Inflow (in Acre-Feet)")
```

Median Neuse Inflow




```
ggplot(data = neuse_average_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Neuse Inflow", x = "Year", y = "Median Inflow (in Acre-Feet)")
```

Median Neuse Inflow




```
#Mean Neuse inflow
ggplot(data = neuse_inflows) +
  geom_col(data = neuse_inflows, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Neuse Inflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Neuse Inflow




```
ggplot(data = neuse_average_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Neuse Inflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Neuse Inflow




```
#Mean volume
ggplot(data = neuse_volume_mean) +
  geom_col(data = neuse_volume_mean, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Neuse Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Neuse Volume



```
ggplot(data = neuse_volume_mean, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Neuse Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```


Mean Neuse Volume



And now with only 2000 and beyond:


```
#Filter out the data to only include 2000 and beyond.
neuse_2000 <- neuse_inflows %>%
  filter(year >= 2000)
#Filter out the average median outflow data to only include 2000 and beyond.
neuse_2000_median <- neuse_average_median %>%
  filter(year >= 2000)
#Filter out the average mean outflow data to only include 2000 and beyond.
neuse_2000_mean <- neuse_average_mean %>%
  filter(year >= 2000)
#Filter out the average volume data to only include 2000 and beyond.
neuse_2000_volume <- neuse_volume_mean %>%
  filter(year >= 2000)
#Median Neuse inflow
ggplot(data = (neuse_2000)) +
  geom_col(data = neuse_2000, aes(x = year, y = median_flow)) +
  labs(title = "Median Neuse Inflow", x = "Year", y = "Median Inflow (in Acre-Feet)")
```

Median Neuse Inflow




```
ggplot(data = neuse_2000_median, aes(x = year, y = avg_median_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Median Neuse Inflow", x = "Year", y = "Median Inflow (in Acre-Feet)")
```

Median Neuse Inflow




```
#Mean Neuse inflow
ggplot(data = neuse_2000) +
  geom_col(data = neuse_2000, aes(x = year, y = mean_flow)) +
  labs(title = "Mean Neuse Inflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Neuse Inflow




```
ggplot(data = neuse_2000_mean, aes(x = year, y = avg_mean_flow)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Neuse Inflow", x = "Year", y = "Mean Inflow (in Acre-Feet)")
```

Mean Neuse Inflow



```
#Mean Neuse volume
ggplot(data = neuse_2000_volume) +
  geom_col(data = neuse_2000_volume, aes(x = year, y = avg_vol)) +
  labs(title = "Mean Neuse Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Neuse Volume



```
ggplot(data = neuse_2000_volume, aes(x = year, y = avg_vol)) +
  geom_point() +
  geom_smooth(method = "lm") +
  labs(title = "Mean Neuse Volume", x = "Year", y = "Mean Volume (in Acre-Feet)")
```

Mean Neuse Volume

