

# BAM\_RAB v 0.1

RFL, DTW, BPC

2022-11-17

## Preliminary Analyses

This document contains preliminary analyses regarding the BAM\_RAB data set as it stands. These preliminary analyses focus on the following elements: As, Cd, Cu, Fe, Pb, Se and Zn. These elements were selected based on interest in the “Big 5” contaminants of concern plus Fe due to our interest in particle formation. I also added Se given the current interests of our extended research group, as well as some interesting parallels between the behavior of Se and Zn.

### 1. Temporal patterns in metal accumulation in biomass contents (mg/g).

The following figures represent the patterns for all samples over time within the three biomass compartments, all sites included. The code for generation all these figures is located under 1\_code/2\_RFL\_Presentation\_Figures/1\_Barplots.

Arsenic (Fig.1) seems to be the only element that actually decreases over time as the growing season progresses. I think this would be the main suspect when we actually try to trace the relationship between metals in algae and metals in water.




Figure 1: Figure 1.1: Arsenic content of benthic biomass compartments over time.

For most metals, patterns are pretty similar (Figs. 2,3,4,5).




Figure 2: Figure 1.2: Cadmium content of benthic biomass compartments over time.




Figure 3: Figure 1.3: Copper content of benthic biomass compartments over time.




Figure 4: Figure 1.4: Iron content of benthic biomass compartments over time.




Figure 5: Figure 1.5: Lead content of benthic biomass compartments over time.

Now we we get to the 2 elements that actually differ. Selenium seems to peak earlier in the season (when compared to Cd,Cu, Fe and Pb).




Figure 6: Figure 1.6: Selenium content of benthic biomass compartments over time.

And of course, as discussed before, Zn is not really present in filamentous algae. Looking at the other compartments it does not seem to present any temporal trends.




Figure 7: Figure 1.7: Zinc content of benthic biomass compartments over time.

## 2.Temporal patterns in metal “stocks” for individual sites.

## Correlation between metals among compartments

## Principal component analyses of contents and stocks by compartment.

PCA analysis was performed in a two tier fashion. Firstly, I performed a PCA analysis on all samples using the BIG5+ elements (As,Cd,Cu,Fe,Pb,Se and Zn) for both metal contents (mg/g) and stocks (mg/m2). However, due to the high collinearity (next figure, still figuring out figure numbering in Markdown) I performed a Variance Inflation test to eliminate highly collinear variables and retain the elements that comprise the highest amount of variance. I did this independently for contents and stocks. I'm calling this group of variables "LEASTVIF".




Figure 8: Correlation plots of BIG5+ elemental contents for all biomass compartments combined.




Figure 9: Correlation plots of BIG5+ elemental contents for all biomass compartments in separate.

These are the PCA results for contents using the BIG5+ elements. Variables have been logged, scaled and centered.






Figure 10: Principal component analysis of BIG5+ elemental contents for three different biomass compartments. Secondary X and Y axes (top and right) indicate correlation between variables and principal components.

Here is the correlation plot for the LEASTVIF contents dataset. Variables retained were Be,Ca,Co,Mo,Pb,S,Se,Si and Sn.




Figure 11: Correlation plots of LEASTVIF elemental contents for all biomass compartments combined.

These are the results for contents using the LEASTVIF elements. Variables have been logged, scaled and centered.




Figure 12: Principal component analysis of LEASTVIF elemental contents for three different biomass compartments. Secondary X and Y axes (top and right) indicate correlation between variables and principal components.

Here are the results for stocks for BIG5 and LEASTVIF:




Figure 13: Principal component analysis of BIG5+ elemental stocks for three different biomass compartments. Secondary X and Y axes (top and right) indicate correlation between variables and principal components.




Figure 14: Principal component analysis of LEASTVIF elemental stocks for three different biomass compartments. Secondary X and Y axes (top and right) indicate correlation between variables and principal components.

Next steps are to extract PCA scores for samples and see how they vary with time and pigment contents.

## **Methods**

##Study area

##Field Sampling

###Biomass

###Water

##Biomass sample processing

##Nutrient Analysis

##Metal Analysis

##Metabolic measurements

##Other?

## **Results**

##General patterns

Are we in a position to start thinking about these?

## **Discussion**