

Textbook — Notes

Boas: *Mathematical Methods in the Physical Sciences*

May 12, 2026

Contents

1 Infinite Series, Power Series

1.1 Convergence and Divergence

1.2 Convergence Tests

1.3 Power Series

1.4 Taylor and Maclaurin Series

1.5 Binomial Series

2 Complex Numbers

2.1 Complex Algebra

2.2 Polar Form and Euler's Formula

2.3 Powers and Roots

2.4 The Complex Plane; Sets

2.5 Complex Functions

2.6 Cauchy–Riemann Equations

3 Linear Algebra

3.1 Matrices and Determinants

3.2 Linear Systems; Row Reduction

3.3 Vector Spaces and Subspaces

3.4 Eigenvalues and Eigenvectors

3.5 Diagonalization

3.6 Hermitian and Unitary Matrices

3.7 Normal Modes (Application)

4 Partial Differentiation

4.1 Partial Derivatives and Chain Rule

4.2 Total Differentials

4.3 Implicit Differentiation

4.4 Stationary Points; Max/Min

4.5 Lagrange Multipliers

4.6 Change of Variables; Jacobians

5 Multiple Integrals

5.1 Double and Triple Integrals

5.2 Change of Variables

5.3 Surface Area

5.4 Surface Integrals

6 Vector Analysis

6.1 Gradient, Divergence, and Curl

6.2 Line Integrals

6.3 Green's Theorem

6.4 Stokes' Theorem

6.5 Divergence Theorem

6.6 Potentials

7 Fourier Series and Transforms

Abstract

Problems involving vibrations or oscillations occur frequently in physics and engineering. The Fourier series is a tool used to approximate periodic functions with sines and cosines.

7.1 Fourier Series

A function $f(x)$ that is periodic with period $2L$ can be written as

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right)$$

where the **Fourier coefficients** are given by

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx, \quad b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx$$

Remark. The term $a_0/2$ is the average value of f over one period.

Dirichlet Conditions

The Fourier series converges to $f(x)$ at all points where f is continuous, provided:

- (i) f is periodic,
- (ii) f is piecewise continuous on $[-L, L]$,
- (iii) f has finitely many maxima and minima per period.

At a jump discontinuity at x_0 , the series converges to the average *Remark.* Inverse trig functions cannot be represented by a Fourier series b/c they are not single-valued. Use Principle Branch workaround. $\frac{1}{2}[f(x_0^+) + f(x_0^-)]$.

Complex Form


The Fourier series can equivalently be written as

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{in\pi x/L}, \quad c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-in\pi x/L} dx$$

The real and complex coefficients are related by $c_n = \frac{1}{2}(a_n - ib_n)$ and $c_{-n} = \frac{1}{2}(a_n + ib_n)$.

Example 7.1: Square wave

Let $f(x)$ be the 2π -periodic square wave with $f(x) = 1$ for $0 < x < \pi$ and $f(x) = -1$ for $-\pi < x < 0$. Then $a_n = 0$ (odd function) and



$$b_n = \frac{2}{n\pi} (1 - \cos n\pi) = \begin{cases} \frac{4}{n\pi} & n \text{ odd} \\ 0 & n \text{ even} \end{cases}$$

so

$$f(x) = \frac{4}{\pi} \left(\sin x + \frac{\sin 3x}{3} + \frac{\sin 5x}{5} + \cdots \right)$$

7.2 Even and Odd Functions

Definition 7.1: Even and Odd Functions

A function is *even* if $f(-x) = f(x)$ and *odd* if $f(-x) = -f(x)$.

Key consequences for Fourier series on $[-L, L]$:

- **Even** f : all $b_n = 0$; the series contains only cosines.

$$a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi x}{L} dx$$

- **Odd** f : all $a_n = 0$; the series contains only sines.

$$b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi x}{L} dx$$

Useful identities: even $\times$ even = even; odd $\times$ odd = even; even $\times$ odd = odd.

7.3 Half-Range Series

Given $f(x)$ defined only on $[0, L]$, we can extend it to $[-L, L]$ in two ways and obtain a *half-range* series:

- **Half-range cosine series** (even extension):

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{L}, \quad a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi x}{L} dx$$

- **Half-range sine series** (odd extension):

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}, \quad b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi x}{L} dx$$

Remark. The choice of extension affects convergence at the endpoints $x = 0$ and $x = L$. The two series are generally different functions outside $[0, L]$.

7.4 Parseval's Theorem

Theorem 7.1: Parseval's Theorem

If $f(x)$ has Fourier coefficients a_n, b_n on $[-L, L]$, then

$$\frac{1}{L} \int_{-L}^L [f(x)]^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2)$$

Parseval's theorem equates the *energy* (mean-square value) of f to the sum of the squared amplitudes of its frequency components. It is useful for evaluating infinite series.

Example 7.2: A classic sum

The Fourier series of $f(x) = x$ on $[-\pi, \pi]$ gives $b_n = 2(-1)^{n+1}/n$. Applying Parseval's theorem:

$$\frac{1}{\pi} \int_{-\pi}^{\pi} x^2 dx = \sum_{n=1}^{\infty} \frac{4}{n^2} \implies \frac{2\pi^2}{3} = 4 \sum_{n=1}^{\infty} \frac{1}{n^2} \implies \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

7.5 Fourier Transforms

For a non-periodic function $f(t)$ defined on $(-\infty, \infty)$, the Fourier series generalizes to the **Fourier transform pair**:

$$\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt \quad (\text{forward transform})$$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\omega) e^{i\omega t} d\omega \quad (\text{inverse transform})$$

Key Properties

Property	Statement
Linearity	$\widehat{af + bg} = a\hat{f} + b\hat{g}$
Time shift	$\widehat{f(t - t_0)} = e^{-i\omega t_0} \hat{f}(\omega)$
Frequency shift	$\widehat{e^{i\omega_0 t} f(t)} = \hat{f}(\omega - \omega_0)$
Differentiation	$\widehat{f'(t)} = i\omega \hat{f}(\omega)$
Scaling	$\widehat{f(at)} = \frac{1}{ a } \hat{f}(\omega/a)$

Common Transform Pairs

$$e^{-a|t|} \longleftrightarrow \frac{2a}{a^2 + \omega^2}, \quad a > 0$$

$$e^{-at^2} \longleftrightarrow \sqrt{\frac{\pi}{a}} e^{-\omega^2/(4a)}, \quad a > 0$$

$$\delta(t) \longleftrightarrow 1, \quad 1 \longleftrightarrow 2\pi \delta(\omega)$$

Remark. Different textbooks use different conventions for the factor of 2π and the sign in the exponent. Boas uses the convention above.

7.6 Convolution

Definition 7.2: Convolution

The *convolution* of two functions f and g is

$$(f * g)(t) = \int_{-\infty}^{\infty} f(\tau) g(t - \tau) d\tau$$

Theorem 7.2: Convolution Theorem

The Fourier transform of a convolution is the product of the transforms:

$$\widehat{f * g}(\omega) = \hat{f}(\omega) \cdot \hat{g}(\omega)$$

Equivalently, the transform of a product is (up to 2π) the convolution of the transforms:

$$\widehat{f \cdot g}(\omega) = \frac{1}{2\pi} (\hat{f} * \hat{g})(\omega)$$

Properties of Convolution

- Commutative: $f * g = g * f$
- Associative: $f * (g * h) = (f * g) * h$
- Distributive: $f * (g + h) = f * g + f * h$

Remark. Convolution appears naturally in linear systems: if $h(t)$ is the impulse response of a system and $x(t)$ is the input, the output is $y(t) = (h * x)(t)$.

Remark. I will add given examples from the Final Review here after I am finished going through the chapters.

8 Ordinary Differential Equations

8.1 First-Order ODEs

8.2 Second-Order Linear ODEs

8.3 Variation of Parameters

8.4 Power Series Solutions

8.5 Frobenius Method

8.6 Bessel's Equation (intro)

9 Tensor Analysis

- 9.1 Cartesian Tensors; Transformation Law
- 9.2 Tensor Algebra
- 9.3 Symmetric and Antisymmetric Tensors
- 9.4 Kronecker Delta and Levi-Civita Symbol
- 9.5 Isotropic Tensors
- 9.6 Physical Applications

10 Series Solutions and Special Functions

10.1 Legendre's Equation

10.2 Legendre Polynomials

10.3 Orthogonality and Expansion

10.4 Bessel's Equation

10.5 Bessel Functions J_ν and Y_ν

10.6 Orthogonality of Bessel Functions

10.7 Applications

11 Partial Differential Equations

11.1 Separation of Variables

11.2 Heat Equation

11.3 Wave Equation

11.4 Laplace's Equation

11.5 Boundary Conditions

11.6 Sturm–Liouville Theory

11.7 2D and 3D Problems

12 Functions of a Complex Variable

12.1 Analytic Functions (revisited)

12.2 Contour Integrals

12.3 Cauchy's Theorem and Integral Formula

12.4 Laurent Series

12.5 Singularities and Residues

12.6 Residue Theorem

12.7 Evaluation of Real Integrals

A Formula Sheet

Legendre Polynomials

Bessel Function Zeros

Gamma Function

Common Fourier Transforms

Vector Identities