

Alcohol QCM Sensor Dataset Data Set

Download: [Data Folder](#), [Data Set Description](#)

Abstract: Five different QCM gas sensors are used, and five different gas measurements (1-octanol, 1-propanol, 2-butanol, 2-propanol and 1-isobutanol) are conducted in each of these sensors.

Data Set Characteristics:	Multivariate	Number of Instances:	125	Area:	Computer
Attribute Characteristics:	Real	Number of Attributes:	8	Date Donated	2019-07-22
Associated Tasks:	Classification, Regression, Clustering	Missing Values?	N/A	Number of Web Hits:	13903

Source:

Provide the names, email addresses, institutions, and other contact information of the donors and creators of the data set.

M. Fatih ADAK, fatihadak '@' sakarya.edu.tr, Sakarya University, Turkey

Peter Lieberzeit, University of Vienna, Austria

Purim Jarujamrus, Ratchathani University, Thailand

Nejat Yumusak, Sakarya University, Turkey

Data Set Information:

In the dataset there are 5 types of dataset.

QCM3, QCM6, QCM7, QCM10, QCM12

In each of dataset, There is alcohol classification of five types,

1-octanol, 1-propanol, 2-butanol, 2-propanol, 1-isobutanol

Attribute Information:

The gas sample is passed through the sensor in five different concentrations. These concentrations are,

Concentration Air ratio (ml) Gas ratio (ml)

1 0.799 0.201

2 0.700 0.300

3 0.600 0.400

4 0.501 0.499

5 0.400 0.600

There are two channels in the sensor. One of these circles forms channel 1, and the other forms channel 2. MIP and MP ratios used in the QCM sensors are,

Sensor name MIP ratio NP ratio

QCM3 1 1

QCM6 1 0

QCM7 1 0.5

QCM10 1 2

QCM12 0 1