



UC | Chile



Ultracold polarons and impurities in one-dimensional optical lattices

Felipe Isaule

ANID Postdoctoral Fellow
Instituto de Física, PUC, Chile

Universitat de Barcelona

25 November, 2025

Ultracold polarons and impurities


1. Polarons and **impurities** in **ultracold atomic systems**.
2. One-dimensional **harmonically confined lattice polarons and counterflows**.
3. Mobile **impurity** interacting with a **Hubbard chain**.
4. Summary and **future work**.

Ultracold polarons and impurities


1. Polarons and **impurities** in **ultracold atomic systems**.
2. One-dimensional harmonically confined lattice polarons and counterflows.
3. Mobile **impurity** interacting with a **Hubbard chain**.
4. Summary and **future work**.

Polarons and impurities


- The study of **impurities** immersed in a **quantum medium** has a **long history** and is **relevant in many fields** of physics.



Electrons in an ionic crystal
L. Landau and S. Pekar, Zh. Eksp. Teor. Fiz. 18, 419 (1948).



^3He impurities in ^4He
G. Baym and C. Pethick, "Landau Fermi-Liquid Theory: Concepts and Application" (1991).



Impurities in nuclear systems
Tajima et al., AAPPS Bulletin, 34, 9 (2024).


- Such **impurities** often **form** dressed **quasiparticles** referred to as **polarons**.

Polarons in ultracold atomic gases

- The **study of polarons** has been **revitalised** thanks to their **experimental realisation** in **ultracold atomic gases**.
- Impurities are realised with **highly imbalanced atomic mixtures**.


C. Baroni, G. Lamporesi and M. Zaccanti, Nat. Rev. Phys. **6**, 736 (2024).

Fermi polaron



A. Schirotzek et al., PRL **102**, 230402 (2009).

Bose polaron



M.-G. Hu et al., PRL **117**, 055301 (2016).

- **Ultracold atoms offer high controllability**, being an ideal platform for studying impurities and polarons.


P. Massignan et al., Rep. Prog. Phys. **77** 034401 (2014).

F. Grusdt et al., Rep. Prog. Phys. **88**, 066401 (2025).

P. Massignan et al., arXiv:2501.09618 (2025).

Polarons in ultracold atomic gases

- Several properties can be examined:
 - **Polaron energies, residue, effective masses, etc.**
- Impurities can be used to **probe** and **manipulate** quantum systems:
 - Transport properties, **phase transitions, mediated interactions**, etc.
- Theoretical techniques:
 - Variational approaches, QMC, ED, RG, Tensor Networks, etc.



Spectral response of an impurity In a BEC


N. B. Jørgensen, et al., PRL 117, 055302 (2016).

Lattice polarons

- A rich platform for studying ultracold atomic impurities is optical lattices.

M. Lewenstein *et al.*, *Ultracold Atoms in Optical Lattices: Simulating Quantum Many-Body Systems* (Oxford, 2012).
C. Gross and I. Bloch, Science **357**, 995 (2017).

- Such impurities are now referred to as **lattice polarons**.
- Ultracold atoms in **tight lattices** can be modelled with a **Hubbard-based Hamiltonian**.
- Recent theoretical studies have considered both
 - a bosonic medium*,
V. Colussi, C. Menotti C and A. Recati, PRL **130**, 173002 (2023).
M. Santiago-García, S. Castillo-López and A. Camacho-Guardian, NJP **26**, 063015 (2024).
F. Gómez-Lozada, H. Hiyane, T. Busch, T. Fogarty, PRR **7**, 023053 (2025).
 - and a fermionic medium*.
I. Amelio and N. Goldman, Scipost Physics **16**, 056 (2024).
H. Hu, J. Wang, X.-J. Liu, PRA, **110**, 023314 (2024).
G. Pascual, J. Boronat, K. Van Houcke, PRR **7**, L042024 (2025).



* Non-comprehensive list.

Ultracold polarons and impurities

1. Polarons and impurities in ultracold atomic systems.
2. One-dimensional harmonically confined lattice polarons and counterflows.
3. Mobile impurity interacting with a Hubbard chain.
4. Summary and future work.


Phys. Rev. Lett. **135**, 023404 (2025)

Bosons in optical lattices


- Bosons in optical lattices feature a **superfluid-to-Mott insulator (SF-MI) transition.**
- The study of **impurities across this SF-MI transition** has attracted increasing attention.

V. E. Colussi, F. Caleffi, C. Menotti, and A. Recati, Phys. Rev. Lett **130**, 173002 (2023).

R. Alhyder, V. E. Colussi, M. Čufar, J. Brand, A. Recati, G. M. Bruun, SciPost Phys. **19**, 002 (2025).



Density profiles of bosons in a 1D optical lattice with harmonic confinement.



Phase diagram of the 1D BH model.

S. Ejima et al., PRA **85**, 053644 (2012).

- Bosons in tight lattices are described by the **Bose-Hubbard model**:

$$\hat{H} = -t \sum_i \left(\hat{a}_i^\dagger \hat{a}_{i+1} + \text{h.c.} \right) + V_{\text{ho}} \sum_i i^2 \hat{n}_i$$

Tunnelling

$$+ \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1).$$

Harmonic trap

Boson-boson repulsion

Harmonically confined lattice polarons

- We studied a **single impurity** interacting with a **bosonic bath** in a **one-dimensional harmonically confined optical lattice**.

$$\hat{H} = -t \sum_{\sigma=b,I} \sum_i \left(\hat{a}_{i,\sigma}^\dagger \hat{a}_{i+1,\sigma} + \text{h.c.} \right)$$

Tunnelling


$$+ V_{\text{ho}} \sum_i \sum_{\sigma=b,I} i^2 \hat{n}_{i,\sigma}$$

Harmonic trap

$$+ \frac{U_{bb}}{2} \sum_i \hat{n}_{i,b} (\hat{n}_{i,b} - 1) + U_{bI} \sum_i \hat{n}_{i,b} \hat{n}_{i,I}.$$

Boson-boson repulsion

Bath-impurity repulsion




- We employed the **density matrix renormalisation group (DMRG)** for **large lattices** with a **large number of particles**.


S. R. White, Phys. Rev. Lett. **69**, 2863 (1992). U. Schollwöck, Rev. Mod. Phys. **77**, 259 (2005).

Harmonically confined lattice polarons

- We **considered** a bath that is **completely SF** for $t/U_{bb} > 0.155$ and that has a **MI domain of unity filling** for $t/U_{bb} < 0.155$.
- The main **DMRG simulations** considered $N_b = 40$ bosons.
- However, we obtained **identical results** for **other choices**, and even a **qualitative agreement** with ED.



Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$



Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d / \xi, \quad \xi = d \sqrt{t/V_{ho}}$$

- The system shows **well-defined phases**.

Density profiles



$t/U_{bb} = 0.2$

Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$



Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d / \xi, \quad \xi = d \sqrt{t/V_{\text{ho}}}$$

- **Region i** is a **miscible** phase where the **impurity repels** the compressible SF bath.

Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$



Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d / \xi, \quad \xi = d \sqrt{t/V_{\text{ho}}}$$

- **Region ii** is simply a **phase-separated** configuration.

Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$



Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d / \xi, \quad \xi = d \sqrt{t / V_{\text{ho}}}$$

- **Region iii** is another **miscible phase**. The **MI bath** remains **undisturbed** due to its **incompressibility**.

Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$



Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d/\xi, \quad \xi = d \sqrt{t/V_{ho}}$$

- **Region iv** is another **phase-separated** configuration. The MI bath moves further away than an SF bath.

Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

$$x_i = i d / \xi, \quad \xi = d \sqrt{t/V_{ho}}$$

- **Region v** is a non-trivial phase where the **impurity** exhibits an **extended profile**, and the **combined profile** of the bath and impurity displays a domain of **unity filling**.

Counterflow



- The new state corresponds to a **counterflow phase**.
- It shows **anti-pair order**

$$\mathcal{C}_{\text{AP}} = \langle \hat{a}_{0,I} \hat{a}_{0,b}^\dagger \hat{a}_{i,b}^\dagger \hat{a}_{i,I} \rangle.$$

- Counterflows were **realised experimentally** very recently with **binary Mott insulators**.

Y.-G. Zheng *et al.*, Nat. Phys. **21**, 208 (2025).


- Our results show that **counterflows appear for large population imbalances**.
- **Counterflows show properties of insulators.**



Counterflow

- The **impurity forms** this **correlated state** with (almost) the **whole insulating bath**.
- The **residue** abruptly **vanishes** at the **phase transition**.

$$Z(U_{bI}) = |\langle \Psi(U_{bI} = 0) | \Psi(U_{bI}) \rangle|^2.$$



Counterflow


- We have found that the **profile of the impurity** is that of a **free particle in a square well**

$$n_I^{(\text{CF})}(x_i) = n_i^{(0)} \cos^2(\pi(x_i - \langle x_I \rangle)/\ell_{\text{CF}}),$$

where only $n_i^{(0)}$ and $\langle x_I \rangle$ are extracted from the calculations.

- The **width** is given by:

$$\sum_i n_I^{(\text{CF})} = 1 \quad \rightarrow \quad \ell_{\text{CF}} = 2/(\xi n_I^{(0)}).$$



Counterflow

- We have developed a simple **analytical model** to describe the **counterflow** based on **impurity-hole pairs**


$$|\text{MI}\rangle = \prod_i \hat{a}_{i,b}^\dagger |\emptyset\rangle \quad \rightarrow \quad |\Psi_{\text{CF}}\rangle = \sum_i \alpha_i |i_{\text{IH}}\rangle.$$

$$|i_{\text{IH}}\rangle = \hat{a}_{i,I}^\dagger \hat{a}_{i,b} |\text{MI}\rangle$$

- After some algebra, one gets that $n_I(i) = |\alpha_i|^2$.
- From imposing the **square-well solution**, the **correlator** takes the form

$$C_{(\text{AP})} = \sqrt{n_I^{(0)}} \cos(i\pi(x_i - \langle x_I \rangle)/\ell_{\text{CF}}),$$

which **agrees** with the **numerical solution**.



Harmonically confined lattice polarons: Outlook

- An **impurity** can form a correlated **counterflow state** with a **bosonic bath** in harmonically confined optical lattices.
- This shows that **counterflows** can form in **mixtures** with **high population imbalance**.
- Ideas for future work in harmonically confined lattices:
 - Study dynamics.
 - Consider multiple impurities.
 - Bose-Bose and Bose-Fermi mixtures.
 - Bath with spin-1/2 fermions.



Phys. Rev. Lett. 135, 023404 (2025)

Ultracold polarons and impurities

1. Polarons and **impurities** in ultracold atomic systems.
2. One-dimensional harmonically confined lattice polarons and **counterflows**.
3. Mobile **impurity** interacting with a **Hubbard chain**.
4. Summary and **future work**.

Spin ½ fermions in optical lattices

- **Spin ½ fermions** in an **optical lattice** are described by the (Fermi) **Hubbard model**. Thus, a Hubbard chain is described by:

$$\hat{H} = -t \sum_{\sigma=\uparrow,\downarrow} \sum_i \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{i+1,\sigma} + \text{h.c.} \right) + U \sum_i \hat{n}_{i,\uparrow} \hat{n}_{i,\downarrow}.$$

Tunnelling fermion-fermion repulsion

- In **balanced systems**, $N_f=N_\uparrow=N_\downarrow$, the **filling factor** is defined as

$$\nu_f = N_f/M. \quad M : \text{Number of sites}$$

- At **half filling**, $\nu_f=1/2$, and $U>0$, the lattice features an **insulating state** where each site is occupied only by one fermion.

Spin $\frac{1}{2}$ fermions in optical lattices


- An open chain with $\nu_f \neq 1/2$ features Friedel oscillations.

G. Bedürftig B. Brendel H. Frahm R. M. Noack, Phys. Rev. B **58**, 10225 (1998).

$$N_f = N_\uparrow = N_\downarrow$$

$$\nu_f = \frac{N_f}{M} = \frac{1}{4} \quad \longrightarrow$$

$$n_f(i) = \langle \hat{n}_{i,\uparrow} + \hat{n}_{i,\downarrow} \rangle$$



- For $\nu_f < 1/2$, the number of peaks is equal to the number of fermions of each spin N_f .

Impurity interacting with an open Hubbard chain

- We consider a **single impurity** interacting with **spin ½ fermions** in a **one-dimensional open optical lattice**.

$$\hat{H} = -t \sum_{\sigma=\uparrow,\downarrow} \sum_i \left(\hat{c}_{i,\sigma}^\dagger \hat{c}_{i+1,\sigma} + \text{h.c.} \right) + U \sum_i \hat{n}_{i,\uparrow} \hat{n}_{i,\downarrow}$$


Tunnelling fermions fermion-fermion repulsion

$$-t \sum_i \left(\hat{a}_i^\dagger \hat{a}_{i+1} + \text{h.c.} \right) + U_{fI} \sum_{\sigma=\uparrow,\downarrow} \sum_i \hat{n}_{i,\sigma} \hat{n}_{i,I}.$$

Tunnelling impurity fermion-impurity interaction

- We use the **exact diagonalisation** method for small lattices with a few particles.


Density profiles $\nu_f=1/4$



- The system undergoes **phase separations** between the impurity and either the **holes or the fermions**.
- The **position of the impurity** can be influenced by the **Friedel oscillations**.


Spin 1/2 fermions in optical lattices

- For $\nu_f > 1/2$, the Hubbard model shows **Friedel oscillations** with a number of **local minima** equal to the number of holes $1 - N_f$.



- This is due to the **particle-hole symmetry** of the Hubbard model.

Density profiles $\nu_f=3/4$



- The **particle-hole symmetry** imposes:

$$n_f(i; U_{fI}, \nu_f) + n_f(i; -U_{fI}, 1 - \nu_f) = 2, \quad n_I(i; U_{fI}, \nu_f) = n_I(i, -U_{fI}, 1 - \nu_f).$$

Impurity interacting with a Hubbard chain: Outlook

- The system can undergo a **phase separation** between the impurity and either the **fermions or the holes**.
- The **Friedel oscillations** induce an **unusual localisation** of the impurity for intermediate interactions.
- Similar physics can be found at other fillings $\nu_f \neq 1/2$.
- Future work:
 - Larger lattices.
 - Lattice with harmonic confinement.

Ultracold polarons and impurities

1. Polarons and **impurities** in ultracold atomic systems.
2. One-dimensional harmonically confined lattice polarons and **counterflows**.
3. Mobile **impurity** interacting with a **Hubbard chain**.
4. Summary and **future work**.

Summary

- Polarons and **impurities** offer very **rich physics** in **optical lattices**.
- We have found the onset of a correlated **counterflow** bath-impurity state in **harmonically confined optical lattices**.
- **Friedel oscillations** can have an **impact** on the behaviour of **impurities**.
- Future work:
 - Further studies of **impurities** and **mixtures** in **harmonically confined optical lattices**.

Collaborators



Bruno Juliá Díaz
UB



Abel Rojo Francàs
OIST (ex UB)



Luis Morales Molina
PUC Chile



Duc Tuan Hoang
OIST



Thomás Fogarty
OIST



Thomas Busch
OIST

Funding



FONDECYT Postdoctorado
No. 3230023

THANK YOU!