



UC | Chile



Ultracold polarons and impurities

From lattice polarons to effective field theories

Felipe Isaule

ANID Postdoctoral Fellow
Instituto de Física, PUC, Chile

Quantum Systems Unit, OIST
25 July, 2025

Ultracold polarons and impurities


1. Quantum **polarons and impurities** in ultracold atomic systems.
2. One-dimensional **harmonically confined lattice polarons and counterflows**.
3. Polaron physics with the **functional renormalisation group**.
4. Summary and **future work**.

Ultracold polarons and impurities


1. Quantum **polarons and impurities** in ultracold atomic systems.
2. One-dimensional harmonically confined lattice polarons and **counterflows**.
3. Polaron physics with the **functional renormalisation group**.
4. Summary and **future work**.

Polarons and impurities

- The study of **impurities** immersed in **quantum mediums** has a **long history** and is **relevant in many fields** of physics.




Electrons in an ionic crystal
L. Landau and S. Pekar, Zh. Eksp. Teor. Fiz. **18**, 419 (1948).




^3He impurities in ^4He
G. Baym and C. Pethick, "Landau Fermi-Liquid Theory: Concepts and Application" (1991).

Polaronic proton



Polaronic alpha particle



Impurities in nuclear systems
Tajima et al., AAPPS Bulletin, **34**, 9 (2024).

- Such **impurities** often form dressed **quasiparticles** referred to as **polarons**.

Polarons in ultracold atomic gases

- The study of polarons has been revitalised thanks to their experimental realisation in ultracold atomic mixtures with a high population imbalance.

P. Massignan et al., Rep. Prog. Phys. **77**, 034401 (2014). F. Grusdt et al. Rep. Prog. Phys. **88**, 066401 (2025).




Illustration of a Fermi Polaron
A. Schirotzek et al., PRL **102**, 230402 (2009).




Illustration of a Bose Polaron
M.-G. Hu et al., PRL **117**, 055301 (2016).

- Several relevant properties can be examined: quasiparticle or **polaron energies**, **residue** and **orthogonality catastrophe**, coherence, effective masses, etc.
- They could also be used to **probe** and **manipulate** quantum **many-body systems**: transport properties, **phase transitions**, **mediated interactions**, etc.

Polarons in ultracold atomic gases


- Different configurations can be studied:
 - **Statistics:** Fermi and Bose polarons.
 - **Number of impurities:** Bipolarons and multipolarons.
 - **Confinement:** Optical lattices, harmonic traps, dimensionality.
 - **Stationary properties or dynamics.**
- Well-posed problem:
 - Variational approaches, Quantum Monte-Carlo (QMC), Exact Diagonalisation (ED), Tensor Networks, Effective Field Theories (EFT), amongst others.
- **Polarons offer a rich combination of few- and many-body physics**, and they are also a **starting point to study more complicated configurations.**

Ultracold polarons and impurities

1. Quantum **polarons and impurities** in ultracold atomic systems.
2. One-dimensional **harmonically confined lattice polarons and counterflows**.
3. Polaron physics with the **functional renormalisation group**.
4. Summary and **future work**.

Lattice polarons


- A rich platform for studying ultracold atomic impurities is tight optical lattices.
- Such impurities are now referred to as **lattice polarons**.
- The study of **lattice polarons** in **bosonic media**s has gained significant attention in the past few years.
- One relevant question is how the **polaron properties** change across the **superfluid-to-Mott insulator (SF-MI) transition**.
- We have tried to address this question in **one-dimensional lattices** with **numerical techniques**.



V. E. Colussi, F. Caleffi, C. Menotti, and A. Recati, Phys. Rev. Lett **130**, 17, 3002 (2023).

Harmonically confined lattice polarons

- We have recently studied a **single impurity** interacting with a **bosonic bath** in a **one-dimensional harmonically confined optical lattice**.



- We have investigated this system using the **density matrix renormalisation group (DMRG)** for **large lattices** with a **large number of particles**.
S. R. White, Phys. Rev. Lett. **69**, 2863 (1992). U. Schollwöck, Rev. Mod. Phys. **77**, 259 (2005).
 - The latter enables us to examine **sharp phase transitions** instead of the smooth crossovers observed for a few particles with ED.

S. R. White, Phys. Rev. Lett. **69**, 2863 (1992). U. Schollwöck, Rev. Mod. Phys. **77**, 259 (2005).

Harmonically confined Bose-Hubbard model

- But firstly, a reminder on **bosons** in a **harmonically confined optical lattice**.

$$\hat{H} = -t \sum_i \left(\hat{a}_i^\dagger \hat{a}_{i+1} + \text{h.c.} \right) + V_{\text{ho}} \sum_i i^2 \hat{n}_i + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1).$$

Tunnelling Harmonic trap Boson-boson repulsion


- The system shows **SF and MI domains**.
- For **large enough systems**, the properties become **invariant for equal characteristic density**

G. G. Batrouni *et al.*, Phys. Rev. A **78**, 023627 (2008)

$$\tilde{\rho} = dN/\xi, \quad \xi = d\sqrt{t/V_{\text{ho}}}.$$

- The profiles scale with the **rescaled length**


$$x_i = i d/\xi_i.$$




F. Isaule, A. Rojo-Francàs, L. Morales-Molina and B. Juliá-Díaz, Phys. Rev. Lett. **135**, 023404 (2025)

Harmonically confined lattice polarons

- We **considered** a **bath** with a characteristic density that is **completely SF** for $t/U_{bb} > 0.155$ and that has a **MI domain of unity filling** for $t/U_{bb} < 0.155$.
- This choice enables us to nicely **examine the behaviour of the impurity** across the **SF-MI phase transition**.
- We performed **DMRG simulations** for $N_b=40$ bosons. We also obtained **identical results for other choices**, and even a **qualitative agreement** with **ED**.



Harmonically confined lattice polarons: Density profiles



- The system shows **well-defined phases**.

Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$



Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

Harmonically confined lattice polarons: Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:


$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

- Region **i** is a **miscible** phase where the **impurity repels** the **compressible SF bath**.

Harmonically confined lattice polarons: Density profiles



- Region ii is simply a **phase-separated** configuration.

Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$



Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

Harmonically confined lattice polarons: Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:


$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

- Region **iii** is another **miscible phase**. The **MI bath** remains **undisturbed** due to its **incompressibility**.

Harmonically confined lattice polarons: Density profiles



- Region **iv** is another **phase-separated** configuration. The MI bath moves further away than an SF bath.

Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$


Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$



Harmonically confined lattice polarons: Density profiles



- At $U_{bI}=U_{bb}$ the impurity simply behaves as one particle in a **one-component system** with N_b+1 bosons.

F. Isaule, A. Rojo-Francàs, L. Morales-Molina and B. Juliá-Díaz, Phys. Rev. Lett. **135**, 023404 (2025)

Harmonically confined lattice polarons: Density profiles



Density profiles:

$$n_\sigma(i) = \langle \hat{n}_{i,\sigma} \rangle$$

Average position:

$$\langle x_\sigma \rangle = \frac{1}{N_\sigma} \sum_i x_i n_\sigma(i)$$

Impurity cloud:

$$\langle x_I^2 \rangle = \frac{1}{N_\sigma} \sum_i x_i^2 n_I(i)$$

- Region v is a non-trivial phase where the **impurity** shows an **extended profile** and the **combined profile** of bath and impurity display a domain of **unity filling**.

Harmonically confined lattice polarons: Counterflow

- The new state corresponds to a **counterflow phase**, which **only appears** with **harmonic confinement**.
- It shows **long-range anti-pair order**



$$\mathcal{C}_{AP} = \langle \hat{a}_{0,b} \hat{a}_{0,I}^\dagger \hat{a}_{i,b}^\dagger \hat{a}_{i,I} \rangle.$$

- **Supercounterflows** have been **studied theoretically** in the past in **two-component bosonic mixtures**.

A. B. Kuklov and B. V. Svistunov, PRL **90**, 100401 (2003). C. Menotti and S. Stringari, PRA **81**, 045604 (2010).

- They were **realised experimentally** very recently with **binary Mott insulators**.

Y.-G. Zheng et al., Nat. Phys. **21**, 208 (2025).




F. Isaule, A. Rojo-Francàs, L. Morales-Molina and B. Juliá-Díaz, Phys. Rev. Lett. **135**, 023404 (2025)

Harmonically confined lattice polarons: Counterflow

- Counterflows are insulators, explaining the combined insulator domain.
- Counterflows require commensurable filling or a harmonic trap.
- We showed that counterflows appear for large population imbalance.
- The impurity forms a correlated state with the whole insulating bath.
- The residue abruptly vanishes at the phase transition.

$$Z(U_{bI}) = |\langle \Psi(U_{bI} = 0) | \Psi(U_{bI}) \rangle|^2.$$



Harmonically confined lattice polarons: Counterflow


- We have found that the **profile** of the **impurity** is that of a **free particle in a square well**

$$n_I^{(\text{CF})}(x_i) = n_I^{(0)} \cos^2(\pi(x_i - \langle x_I \rangle)/\ell_{\text{CF}}),$$

where only $n_I^{(0)}$ and $\langle x_I \rangle$ are extracted from the calculations.

- The **width** is given by:

$$\sum_i n_I^{(\text{CF})} = 1 \quad \rightarrow \quad \ell_{\text{CF}} = 2/(\xi n_I^{(0)}).$$



Harmonically confined lattice polarons: Counterflow


- We have developed a simple **analytical model** to describe the **counterflow** based on **impurity-hole pairs**

$$|\text{MI}\rangle = \prod_i \hat{a}_{i,b}^\dagger |\emptyset\rangle \quad \longrightarrow \quad |\Psi_{\text{CF}}\rangle = \sum_i \alpha_i |i_{\text{IH}}\rangle.$$
$$|i_{\text{IH}}\rangle = \hat{a}_{i,I}^\dagger \hat{a}_{i,b} |\text{MI}\rangle$$

- After some algebra, one gets that $n_I(i) = |\alpha_i|^2$.
- From imposing the **square-well solution**, the **correlator** takes the form

$$C_{(\text{AP})} = n_I^{(0)} \cos(i\pi(x_i - \langle x_I \rangle)/\ell_{\text{CF}}),$$

which **agrees** with the **numerical solution**.



Harmonically confined lattice polarons: Outlook

- An **impurity** forms a **correlated counterflow state** with an **insulating bath** in **harmonically confined optical lattices**.
- The **impurity** behaves as a **free particle** in a **square well**.
- Ideas for future work in **harmonically confined lattices**:
 - Study **dynamics**.
 - Consider **multiple impurities**.
 - Examine the **attractive branch** (molecule formation?).
 - Study balanced **binary mixtures** (droplets, supercounterflows, etc).

Ultracold polarons and impurities

1. Quantum **polarons and impurities** in ultracold atomic systems.
2. One-dimensional harmonically confined lattice **polarons and counterflows**.
3. Polaron physics with the **functional renormalisation group**.
4. Summary and **future work**.

Bose polarons

- A **Bose polaron** is the quasiparticle formed by an **impurity** interacting with a **weakly-interacting Bose gas** (BEC).


$$\hat{H} = \sum_{k,\sigma=B,I} (\varepsilon_{k,\sigma} - \mu_\sigma) \hat{b}_{k,\sigma}^\dagger \hat{b}_{k,\sigma} + \frac{g_{BB}}{2\mathcal{V}} \sum_{kk'k''} \hat{b}_{k,B}^\dagger \hat{b}_{k',B}^\dagger \hat{b}_{k'-k'',B} \hat{b}_{k'-k'',B}$$

Kinetic energy Chemical potential
Boson-boson repulsion

$$+ \frac{g_{BI}}{\mathcal{V}} \sum_{kk'k''} \hat{b}_{k,B}^\dagger \hat{b}_{k',I}^\dagger \hat{b}_{k'-k'',I} \hat{b}_{k'-k'',B}$$

Bath-impurity interaction

$$\varepsilon_{k,\sigma} = \frac{k^2}{2m_\sigma}$$



- Bose polarons were **achieved experimentally in 2016**.

N. Jørgensen *et al.*, Phys. Rev. Lett. **117**, 055302 (2016), M. Hu *et al.*, Phys. Rev. Lett. **117**, 055301 (2016).

- The branch of **attractive bath-impurity interactions** offers **rich physics**:
 - **Strong coupling regime** (Unitary limit $a_{BI} \rightarrow \infty$), **three- and more-body correlations**, Efimov physics, finite temperatures.

The functional renormalisation group


- **Bose polarons** have been studied with a variety of techniques such as **Quantum MonteCarlo**, variational ansatzë, Frölich model, etc.
- In general, one needs to rely on **non-perturbative methods** to theoretically study the **strongly-interacting regime**.
- One such technique is the **functional renormalisation group (FRG)**.
N. Dupuis *et al.*, Phys. Rep. **910**, 1 (2021).
- The objective of the FRG is to compute an **effective action** Γ (IR) from a microscopic action S (UV).

Wetterich equation:

$$\partial_k \Gamma_k = \frac{1}{2} \text{tr} \left[\partial_k R_k (\Gamma^{(2)} + R_k)^{-1} \right]$$

C. Wetterich, Phys. Lett. B **301**, 90 (1993)

Regulator (“cutoff”)



$$\mathcal{Z}[\varphi] = \int D\varphi e^{-S[\varphi]} \rightarrow \Omega_G = -\frac{1}{\beta} \ln \mathcal{Z} = \frac{1}{\beta} \Gamma[\varphi_{cl,0}]$$

The functional renormalisation group

- The FRG is useful to study **strongly correlated systems** and critical phenomena.
- It has been used in different problems in **ultracold atomic** systems:
 - One-component Bose gases
S. Floerchinger and C. Wetterich, Phys. Rev. A **77**, 053603 (2008). Phys. Rev. A **79**, 013601 (2009).
 - Fermi gases: BCS-BEC crossover
S. Floerchinger *et al.*, Phys. Rev. A **81**, 063619 (2010). I. Boettcher *et al.*, Phys. Rev. A **89**, 053630 (2014).
 - Bose gases in optical lattices
A. Rançon and N. Dupuis, Phys. Rev. A **85**, 063607 (2012), Phys. Rev. A **86**, 043624 (2012).
 - Few bosons: Efimov physics
S. Floerchinger *et al.*, Few-Body Syst. **51**, 153 (2011). R. Schmidt and S. Moroz, Phys. Rev. A **81**, 052709 (2010).
 - Fermi polaron
R. Schmidt and T. Enss, Phys. Rev. A **83**, 063620 (2011). von Milczewski *et al.*, Phys. Rev. A **105**, 013317 (2022).

Field-theory description of Bose Polarons


- The **microscopic model** for the **Bose polaron** problem reads

S. P. Rath and R. Schmidt, Phys. Rev. A **88**, 053632 (2013).

$$\mathcal{S} = \int_x \left[\sum_{\sigma=B,I} \psi_\sigma^\dagger \left(\partial_\tau - \frac{\nabla^2}{2m_\sigma} - \mu_\sigma \right) \psi_\sigma + \nu_\phi \phi^\dagger \phi + \frac{g_{BB}}{2} (\psi_B^\dagger \psi_B)^2 + h_\Lambda (\phi^\dagger \psi_B \psi_I + \phi \psi_B^\dagger \psi_I^\dagger) \right],$$

Kinetic energy Chemical potential Boson-boson repulsion Bath-impurity interaction

where $\phi \sim \psi_B \psi_I$ are **auxiliary dimer fields** that **mediate the bath-impurity interaction**.



Hubbard-Stratonovich transformation

F. Isaule, I. Morera, P. Massignan, and B. Juliá-Díaz, Phys. Rev A. **104**, 023317 (2021).

FRG for Bose polarons

- We employ the following **ansatz** (shortened version):

$$\Gamma_k = \int_x \left[\psi_B^\dagger \left(S_B \partial_\tau - \frac{Z_B}{2m_B} \nabla^2 \right) \psi_B + \psi_I^\dagger \left(S_I \partial_\tau - \frac{Z_I}{2m_I} \nabla^2 + m_I^2 \right) \psi_I \right.$$


$$+ \phi^\dagger \left(S_\phi \partial_\tau - \frac{Z_\phi}{2m_\phi} \nabla^2 + m_\phi^2 \right) \phi + U_B(|\psi_B|^2) + \lambda_{B\phi} |\psi_B|^2 |\phi|^2 + h \left(\phi^\dagger \psi_B \psi_I + \phi \psi_B^\dagger \psi_I^\dagger \right) \right].$$

Boson-boson
repulsion

Three-body
Bath-impurity
interaction



Two-body
Bath-impurity
interaction

- Multi-body correlations** can be added **systematically**.
- We **solved** the RG flow with and without three-body correlations.



F. Isaule, I. Morera, P. Massignan, and B. Juliá-Díaz, Phys. Rev A. **104**, 023317 (2021).

Polaron energy for the three-dimensional Bose Polaron



n : Bath density
 a_{BB} : boson-boson scattering length
 a_{BI} : bath-impurity scattering length

Exp (left): N.B Jørgensen *et al.*, PRL **117**, 055302 (2016).

QMC (left): L. Peña Ardila *et al.*, PRA **99**, 063607 (2019).

QMC (right): L. Peña Ardila and S. Giorgini, PRA **92**, 033612 (2015).

F. Isaule, I. Morera, P. Massignan, and B. Juliá-Díaz, Phys. Rev A. **104**, 023317 (2021).

FRG for Bose polarons

- The **FRG describes** with **high accuracy** the **Bose polaron**, giving a good agreement with Quantum Monte-Carlo.
- It allows us to **quantify** the effect of **three-body correlations** and add them systematically.
- Future work:
 - Polarons across the **BCS-BEC crossover**.
 - Efimov physics in Bose polarons.
 - Bipolarons in BECs.
 - **Bose-Fermi mixtures**.
 - Polarons in **optical lattices**.

Ultracold polarons and impurities

1. Quantum **polarons and impurities** in ultracold atomic systems.
2. One-dimensional harmonically confined lattice **polarons and counterflows**.
3. Polaron physics with the **functional renormalisation group**.
4. Summary and **future work**.

Summary

- We have **numerically** studied **impurities** interacting with **bosonic baths** in **optical lattices** across the **SF-MI transition**.
- We have found the onset of a correlated **counterflow** bath-impurity state in **harmonically confined optical lattices**.
- The **FRG** is a **powerful technique** to study **strongly-interacting polarons**.
- Future work:
 - Further studies of **impurities** and **mixtures** in **harmonically confined optical lattices**.
 - Employ the **FRG** to study **more complicated polaron** configurations.