

Grouping of sugarcane genotypes in function of their genetic dissimilarity with focus for the production of brown sugar

Patrícia Jesus de Melo¹, Alessandro Dal' Col Lúcio², Elisangela Marques Jeronimo³, Tiago Olivoto¹, Maria Inês Diel¹, Bruno Giacomini Sari¹, Darlei Michelski Lambert⁴, João Alberto Zemolin⁴

Abstract: The objective was to group the genotypes according to genetic similarity, intended for the production of brown sugar. Ten genotypes of sugarcane harvested at 15, 17 and 19 months of culture. The IAC genotypes presented grouping among themselves and the genotypes RB966928, RB867515 and SP81-3250 also presented grouping among themselves.

Keywords: *Saccharum* spp; heterotic potential; genetic distance; multivariate analysis.

¹Programa de Pós-graduação em Agronomia (PPG Agronomia), Centro de Ciências Rurais (CCR), Universidade Federal de Santa Maria (UFSM), 97105-900, Santa Maria, RS, Brasil. email: patty_de_melo@yahoo.com.br, tiagoolivoto@gmail.com, brunosari@hotmail.com, mariaines.diel@hotmail.com

²Departamento de Fitotecnia, CCR, UFSM, Santa Maria, RS, Brasil. email: adlucio@ufsm.br

³ Agência Paulista de Tecnologia dos Agronegócios (APTA), Polo Regional Centro Oeste, Avenida Rodrigues Alves, Quadra 27 a Quadra 39, Vila Cardia, 17030-000, Bauru, SP, Brasil. email: elisangela.m.jeronimo@gmail.com

⁴Curso de Agronomia, CCR, UFSM, Santa Maria, RS, Brasil.