Model-Implied Instrumental-Variable and James–Stein Estimation of Structural Equation Models: A Case Study With lavaan

Felipe Fontana Vieira and Yves Rosseel
Department of General and Applied Psychology, Faculty of Psychology and Educational Sciences, Ghent University

Abstract

This article presents model-implied instrumental-variable (MIIV) estimation and scaling (JS) and aggregated (JSA) James–Stein estimation within a common notation, based on their implementation in the R package lavaan. We describe the estimation of factor loadings and latent regression slopes equation by equation, followed by the estimation of variance, covariance, and intercept parameters. We also provide analytical standard errors for JS and JSA using the delta method, accounting for uncertainty across all estimation stages. A simulation study compares the standard error accuracy of these approaches with maximum likelihood (ML) under correct model specification. The standard errors of MIIV, JS, and JSA perform comparably to those of ML under the conditions examined.

Keywords: structural equation modeling, model-implied instrumental variables, James–Stein estimation, delta method, simulation

Structural equation models (SEMs) with latent variables are usually estimated by maximum likelihood (ML), which estimates all parameters jointly. Although ML has desirable asymptotic properties when its assumptions hold, difficulties can arise when it is applied in practice. For instance, a misspecification in one equation can bias estimates in other equations (Bollen et al., 2007; Bollen, 2019; Dhaene & Rosseel, 2023; Kaplan, 1988; see also Jin et al., 2021, for diagonally weighted least squares), and the iterative optimization can fail to converge or end in inadmissible solutions, especially in small samples (Bollen, 2019; Dhaene & Rosseel, 2023). Equation-by-equation estimators offer an alternative (e.g., Bollen, 1996; Burghgraeve et al., 2021). In the approaches considered, factor loadings and latent regression slopes are estimated

---

Felipe Fontana Vieira https://orcid.org/0009-0006-0949-6569
Yves Rosseel https://orcid.org/0000-0002-4129-4477

The authors report there are no competing interests to declare. Generative AI (Claude Opus 5.5, Anthropic) was used to create all figures, with the TikZ and R code based on earlier files written by the first author, and to improve the flow of the Introduction, Results, and Discussion. Commits containing AI generated code or text are marked as such in the commit history of the project repository. All AI output and suggestions were checked by the first author, and the authors take full responsibility for the content of this article. This project was funded by the Special Research Fund (BOF) of Ghent University (BOF/24J/2023/144). Author roles were classified using the Contributor Role Taxonomy (CRediT; https://credit.niso.org/) as follows: Felipe Fontana Vieira: Conceptualization, Methodology, Software, Investigation, Data curation, Formal analysis, Visualization, Writing - original draft, Writing - review & editing; Yves Rosseel: Conceptualization, Methodology, Software, Supervision, Writing - review & editing, Funding acquisition

Correspondence concerning this article should be addressed to Felipe Fontana Vieira, Department of General and Applied Psychology, Faculty of Psychology and Educational Sciences, Ghent University, Henri Dunantlaan 2, Ghent 9000, Belgium, Email: felipe.fontanavieira@ugent.be

one equation at a time and in closed form, which may limit the propagation of misspecification across equations (Bollen et al., 2018; Burghgraeve et al., 2021) and avoids iterative optimization for this part of the estimation. Two such estimators are considered. Model-implied instrumental-variable (MIIV) estimation replaces each latent variable by a scaling indicator and uses instruments implied by the model to correct for the resulting measurement error (Bollen, 1996, 2019; Bollen & Bauer, 2004). James–Stein estimation, proposed for SEMs by Burghgraeve et al. (2021), instead replaces each latent predictor by an estimate of its conditional expectation given observed proxies and then applies least squares. This conditional expectation is estimated using James–Stein type shrinkage (Efron & Morris, 1973, 1975; James & Stein, 1961). The approach follows the principle of regression calibration in errors-in-variables regression (Carroll et al., 2006; Fuller, 1987; Whittemore, 1989). Its scaling version (JS) uses the scaling indicators as proxies, and its aggregated version (JSA) uses reliability weighted sums of indicators. JSA was found to outperform ML and MIIV in terms of mean squared error under the small sample conditions examined (Burghgraeve et al., 2021). Both versions were further evaluated for confirmatory factor models by Dhaene and Rosseel (2024).

Beyond point estimation, the usefulness of these approaches for statistical inference depends on the accuracy of their standard errors. For MIIV, conventional standard errors for factor loadings and latent regression slopes follow from two-stage least squares (Bollen, 1996). Standard errors for these same estimates can also be obtained by the delta method, using derivatives with respect to the sample moments, for which Fisher and Bollen (2020) provided explicit analytical expressions. These derivatives also propagate uncertainty in the estimated loadings and slopes into the standard errors of subsequently estimated variance and covariance parameters. Standard errors for these latter parameters were derived under unweighted least squares by Bollen and Maydeu-Olivares (2007) and extended to general weight matrices by Jin et al. (2021). More recently, Maydeu-Olivares et al. (in press) compared subsequent variance and covariance estimation procedures and the accuracy of their standard errors. For James–Stein estimation, Burghgraeve et al. (2021) derived a sampling variance expression for latent regression estimates that accounts for uncertainty in the estimated reliability adjustment, providing a basis for analytical standard errors for those estimates. However, they did not develop a complete analytical standard error calculation for all model parameters and used bootstrap standard errors for both JS and JSA in their empirical evaluations. The R package lavaan (Rosseel, 2012) now provides analytical standard errors for both estimators by applying the delta method to the complete estimation procedure. In addition, its JS and JSA implementations, initially restricted to confirmatory factor models, have been extended to more general models, and the package also includes MIIV estimation.

Present Study and Contributions

This article makes three minor contributions. First, it describes and conceptually compares MIIV and James–Stein estimation within a common notation, following their implementation in lavaan. Second, it develops and illustrates a complete delta method calculation of standard errors for JS and JSA, accounting for uncertainty across all estimation stages. Third, it evaluates their standard error accuracy in a simulation study comparing JS and JSA with MIIV and ML.

The remainder of the article introduces the model and notation, describes the estimation of factor loadings, latent regression slopes, and the remaining parameters, presents the analytical standard errors, and reports the simulation study.

Model, Notation, and Parameterization

Following the all-y LISREL notation (Jöreskog & Sörbom, 1996, submodel 3B, Chapter 6; Rosseel & Vidal, 2025), let $\mathbf{y} \in \mathbb{R}^p$ contain the continuous observed variables and $\boldsymbol{\eta} \in \mathbb{R}^q$ the latent variables. The measurement and latent variable models are

$$\mathbf{y} = \boldsymbol{\nu} + \boldsymbol{\Lambda}\boldsymbol{\eta} + \boldsymbol{\epsilon}, \qquad \boldsymbol{\eta} = \boldsymbol{\alpha} + \mathbf{B}\boldsymbol{\eta} + \boldsymbol{\zeta},$$

where $\boldsymbol{\nu} \in \mathbb{R}^p$ contains the measurement intercepts, $\boldsymbol{\Lambda} \in \mathbb{R}^{p\times q}$ contains the factor loadings, $\boldsymbol{\alpha} \in \mathbb{R}^q$ contains the structural intercepts, and $\mathbf{B} \in \mathbb{R}^{q\times q}$ contains the latent regression slopes, that is, the regression slopes among the latent variables. The measurement errors $\boldsymbol{\epsilon} \in \mathbb{R}^p$ and structural disturbances $\boldsymbol{\zeta} \in \mathbb{R}^q$ have zero means. Let $\mathbb{S}_+^k$ and $\mathbb{S}_{++}^k$ denote the sets of symmetric positive semidefinite and positive definite $k\times k$ matrices, respectively. Assume $\mathrm{Var}(\boldsymbol{\epsilon}) = \boldsymbol{\Theta} \in \mathbb{S}_+^p$, $\mathrm{Var}(\boldsymbol{\zeta}) = \boldsymbol{\Psi} \in \mathbb{S}_+^q$, and $\mathrm{Cov}(\boldsymbol{\epsilon}, \boldsymbol{\zeta}) = \mathbf{0}_{p\times q}$.

Provided that $\mathbf{I}_q - \mathbf{B}$ is nonsingular, the latent and observed mean vectors are

$$\boldsymbol{\mu}_\eta = (\mathbf{I}_q - \mathbf{B})^{-1}\boldsymbol{\alpha} \in \mathbb{R}^q, \qquad \boldsymbol{\mu}_y = \boldsymbol{\nu} + \boldsymbol{\Lambda}\boldsymbol{\mu}_\eta \in \mathbb{R}^p,$$

where $\mathbf{I}_q$ is the $q \times q$ identity matrix. The implied observed covariance matrix is

$$\boldsymbol{\Sigma} = \boldsymbol{\Lambda}(\mathbf{I}_q - \mathbf{B})^{-1}\boldsymbol{\Psi}(\mathbf{I}_q - \mathbf{B})^{-\top}\boldsymbol{\Lambda}^\top + \boldsymbol{\Theta}.$$

Both estimation procedures considered here require a scaling (marker) indicator parameterization (Bollen, 1996, 2019; Burghgraeve et al., 2021), in which one loading per factor is fixed to one and the corresponding measurement intercept is fixed to zero (Bollen, 1989). Hence, let $\mathbf{m} \in \mathbb{R}^q$ collect the scaling indicators from $\mathbf{y}$, ordered so that the k-th element is the scaling indicator of η_k and loads on η_k only, and let $\boldsymbol{\epsilon}^{(m)} \in \mathbb{R}^q$ collect the corresponding measurement errors. Then $\mathbf{m} = \boldsymbol{\eta} + \boldsymbol{\epsilon}^{(m)}$ and $E(\mathbf{m}) = \boldsymbol{\mu}_\eta$.

Let $\boldsymbol{\theta} \in \mathcal{T} \subseteq \mathbb{R}^d$ collect the free parameters, where $\mathcal{T}$ incorporates identification restrictions. Assume that $\boldsymbol{\Sigma}(\boldsymbol{\theta}) \in \mathbb{S}_{++}^p$. Denote the sample mean vector and covariance matrix by $\overline{\mathbf{y}} \in \mathbb{R}^p$ and $\mathbf{S} \in \mathbb{S}_+^p$, respectively, computed from a single group of independent and identically distributed observations $\mathbf{y}_i$, $i = 1, \ldots, n$, of $\mathbf{y}$, which is the setting of all estimation expressions below.[1] Define

$$\mathbf{s} = \begin{pmatrix} \overline{\mathbf{y}} \\ \mathrm{vech}(\mathbf{S}) \end{pmatrix} \in \mathbb{R}^h, \qquad \boldsymbol{\sigma}(\boldsymbol{\theta}) = \begin{pmatrix} \boldsymbol{\mu}_y(\boldsymbol{\theta}) \\ \mathrm{vech}\{\boldsymbol{\Sigma}(\boldsymbol{\theta})\} \end{pmatrix} \in \mathbb{R}^h, \qquad h = p + \frac{p(p+1)}{2},$$

where vech stacks the lower triangular entries, including the diagonal.

Estimation of Factor Loadings and Latent Regression Slopes

MIIV estimation and James–Stein estimation obtain factor loadings and latent regression slopes equation by equation (Bollen, 1996, 2019; Burghgraeve et al., 2021). Both are estimated from centered observations, with intercepts estimated subsequently. The following expressions assume that the loadings and latent regression slopes being estimated are free and not subject to equality constraints.

Model-Implied Instrumental-Variable Estimation

In MIIV estimation (Bollen, 1996, 2019), the scaling indicator equations are rearranged as

$$\boldsymbol{\eta}_i = \mathbf{m}_i - \boldsymbol{\epsilon}_i^{(m)}$$

and substituted into the remaining measurement and latent variable equations. This latent to observed variable (L2O) transformation produces equations involving observed dependent and explanatory variables, with composite errors containing the original errors or disturbances and scaling indicator measurement errors. The intercepts, factor loadings, and latent regression slopes remain unchanged.

After centering and stacking observations, the j-th L2O equation has the representation

$$\mathbf{d}_j = \mathbf{X}_j\mathbf{a}_j + \mathbf{u}_j,$$

where $\mathbf{d}_j \in \mathbb{R}^n$ contains the centered dependent variable, which is an indicator in a measurement equation or the dependent factor's scaling indicator in a latent variable equation. The matrix $\mathbf{X}_j \in \mathbb{R}^{n\times k_j}$ contains the

[1] Following lavaan, $\mathbf{S}$ uses the divisor n for JS and JSA and $n - 1$ for MIIV estimation.

centered scaling indicators of the k_j latent predictors, $\mathbf{a}_j \in \mathbb{R}^{k_j}$ contains the corresponding loadings or latent regression slopes, and $\mathbf{u}_j \in \mathbb{R}^n$ contains the composite errors. The explanatory variables in $\mathbf{X}_j$ generally correlate with $\mathbf{u}_j$ through their shared scaling indicator measurement errors.

The hypothesized model identifies observed variables that are uncorrelated with the composite error and sufficiently related to the explanatory variables. Let $\mathbf{Z}_j \in \mathbb{R}^{n\times \ell_j}$ contain these centered instruments, with $\ell_j \geq k_j$. In the first stage, the explanatory variables are regressed on the instruments, giving

$$\widehat{\mathbf{X}}_j = \mathbf{P}_j\mathbf{X}_j \in \mathbb{R}^{n\times k_j}, \qquad \mathbf{P}_j = \mathbf{Z}_j(\mathbf{Z}_j^\top\mathbf{Z}_j)^{-1}\mathbf{Z}_j^\top \in \mathbb{R}^{n\times n}.$$

In the second stage, the dependent variable is regressed on these fitted explanatory variables. The resulting two-stage least squares estimator is

$$\widehat{\mathbf{a}}_{j,\mathrm{MIIV}} = (\widehat{\mathbf{X}}_j^\top\widehat{\mathbf{X}}_j)^{-1}\widehat{\mathbf{X}}_j^\top\mathbf{d}_j = (\mathbf{X}_j^\top\mathbf{P}_j\mathbf{X}_j)^{-1}\mathbf{X}_j^\top\mathbf{P}_j\mathbf{d}_j,$$

assuming that $\mathbf{Z}_j$ has full column rank and that $\mathbf{Z}_j^\top\mathbf{X}_j$ has rank k_j.

Scaling James–Stein Estimation

The James–Stein approach replaces the latent predictors in each equation by estimates of their conditional expectations given observed proxies, followed by least squares (Burghgraeve et al., 2021). The implementation considered here is restricted to recursive latent variable models without higher order factors. For the same equation j, let $\mathcal{P}_j$ index its k_j latent predictors. The proxies are indicators that load only on the corresponding latent predictor, with measurement errors $\boldsymbol{\epsilon}_j^{(p)}$ such that $\mathrm{Var}(\boldsymbol{\epsilon}_j^{(p)})$ is diagonal and $\mathrm{Cov}(\boldsymbol{\epsilon}_j^{(p)}, \epsilon_{d_j}) = \mathbf{0}$, where ϵ_{d_j} is the measurement error of the dependent variable. For the latent variable equation of η_k, additionally $\mathrm{Cov}(\zeta_k, \boldsymbol{\eta}_{\mathcal{P}_j}) = \mathbf{0}$. In the scaling version, the proxies are the scaling indicators, whose equations are

$$\mathbf{m}_{i,\mathcal{P}_j} = \boldsymbol{\eta}_{i,\mathcal{P}_j} + \boldsymbol{\epsilon}_{i,\mathcal{P}_j}^{(m)},$$

where all three vectors belong to $\mathbb{R}^{k_j}$. Define

$$\boldsymbol{\Sigma}_{mm,j} = \mathrm{Var}(\mathbf{m}_{i,\mathcal{P}_j}) \in \mathbb{S}_{++}^{k_j}, \qquad \boldsymbol{\Omega}_j = \mathrm{Var}(\boldsymbol{\epsilon}_{i,\mathcal{P}_j}^{(m)}) \in \mathbb{S}_{+}^{k_j}.$$

Under joint normality,

$$E(\boldsymbol{\eta}_{i,\mathcal{P}_j} \mid \mathbf{m}_{i,\mathcal{P}_j}) = \boldsymbol{\mu}_{\eta,\mathcal{P}_j} + \mathbf{R}_j\left(\mathbf{m}_{i,\mathcal{P}_j} - \boldsymbol{\mu}_{\eta,\mathcal{P}_j}\right),$$

where $\boldsymbol{\mu}_{\eta,\mathcal{P}_j} \in \mathbb{R}^{k_j}$ and $\mathbf{R}_j = (\boldsymbol{\Sigma}_{mm,j} - \boldsymbol{\Omega}_j)\boldsymbol{\Sigma}_{mm,j}^{-1} \in \mathbb{R}^{k_j\times k_j}$.

Note that, without normality, the same expression gives the best linear predictor under the stated covariance assumptions (Burghgraeve et al., 2021). Preliminary measurement error variances are obtained using a Spearman type procedure (Burghgraeve et al., 2021). For indicator e with sample variance s_{ee}, the preliminary error variance is $s_{ee}(1 - \widehat{h}_e^2)$, where the communality $\widehat{h}_e^2$ averages $s_{ef}s_{ef'}/(s_{ff'}s_{ee})$ over all pairs (f, f') of other indicators of the same latent variable. Incorporating the small sample correction gives

$$\widehat{\mathbf{R}}_j = \left(\mathbf{S}_{mm,j} - c_n\widehat{\boldsymbol{\Omega}}_j\right)\mathbf{S}_{mm,j}^{-1}, \qquad c_n = \frac{n-3}{n-1},$$

where $n > 3$, $\mathbf{S}_{mm,j} \in \mathbb{S}_{++}^{k_j}$ is the sample covariance matrix of the scaling indicators, and $\widehat{\boldsymbol{\Omega}}_j \in \mathbb{S}_{+}^{k_j}$ is the diagonal matrix of their preliminary measurement error variance estimates. These preliminary error variance estimates are used for prediction and need not equal the final model estimates.

Using the scaling indicator sample means $\overline{\mathbf{m}}_{\mathcal{P}_j} \in \mathbb{R}^{k_j}$, the fitted latent predictors are

$$\widetilde{\boldsymbol{\eta}}_{i,\mathcal{P}_j} = \overline{\mathbf{m}}_{\mathcal{P}_j} + \widehat{\mathbf{R}}_j \left(\mathbf{m}_{i,\mathcal{P}_j} - \overline{\mathbf{m}}_{\mathcal{P}_j} \right) \in \mathbb{R}^{k_j}.$$

Let $\widetilde{\mathbf{X}}_j \in \mathbb{R}^{n \times k_j}$ contain the centered fitted predictors. Using the same dependent variable $\mathbf{d}_j$ as above, the JS estimator of the loadings or latent regression slopes in equation j is

$$\widehat{\mathbf{a}}_{j,\mathrm{JS}} = (\widetilde{\mathbf{X}}_j^\top \widetilde{\mathbf{X}}_j)^{-1} \widetilde{\mathbf{X}}_j^\top \mathbf{d}_j.$$

Aggregated James–Stein Estimation

The JSA estimator replaces each scaling indicator with a weighted aggregate of indicators satisfying the proxy conditions above. For latent predictor η_k in equation j, let $\mathbf{y}_{i,kj} \in \mathbb{R}^{r_{kj}}$ collect these indicators, excluding the dependent variable when estimating a measurement equation. Let $\boldsymbol{\nu}_{kj}$, $\boldsymbol{\lambda}_{kj}$, and $\mathbf{w}_{kj}$, all in $\mathbb{R}^{r_{kj}}$, contain their intercepts, loadings, and weights. Normalizing the aggregate to the latent variable's scale gives

$$g_{ikj} = \frac{\mathbf{w}_{kj}^\top (\mathbf{y}_{i,kj} - \boldsymbol{\nu}_{kj})}{\mathbf{w}_{kj}^\top \boldsymbol{\lambda}_{kj}} = \eta_{ik} + \epsilon_{ikj}^{(A)}, \qquad \mathbf{w}_{kj}^\top \boldsymbol{\lambda}_{kj} \neq 0.$$

The weights are chosen to maximize reliability (Bentler, 1968). In the sample,

$$\widehat{\mathbf{w}}_{kj} \in \arg\max_{\mathbf{w} \neq \mathbf{0}} \left\{ 1 - \frac{\mathbf{w}^\top \widehat{\boldsymbol{\Theta}}_{kj}^{\mathrm{pre}} \mathbf{w}}{\mathbf{w}^\top \mathbf{S}_{kj} \mathbf{w}} \right\},$$

where $\mathbf{S}_{kj} \in \mathbb{S}_{++}^{r_{kj}}$ is the sample covariance matrix of the indicators in the aggregate and $\widehat{\boldsymbol{\Theta}}_{kj}^{\mathrm{pre}} \in \mathbb{S}_{+}^{r_{kj}}$ is the diagonal matrix of their preliminary measurement error variance estimates. Preliminary loading estimates $\widehat{\boldsymbol{\lambda}}_{kj}^{\mathrm{pre}} \in \mathbb{R}^{r_{kj}}$ from JS supply the normalization.

Collect the aggregates and their measurement errors in

$$\mathbf{g}_{ij} = (g_{ikj})_{k \in \mathcal{P}_j} \in \mathbb{R}^{k_j}, \qquad \boldsymbol{\epsilon}_{ij}^{(A)} = (\epsilon_{ikj}^{(A)})_{k \in \mathcal{P}_j} \in \mathbb{R}^{k_j}.$$

Let

$$\boldsymbol{\Sigma}_{gg,j} = \mathrm{Var}(\mathbf{g}_{ij}) \in \mathbb{S}_{++}^{k_j}, \qquad \boldsymbol{\Omega}_{A,j} = \mathrm{Var}(\boldsymbol{\epsilon}_{ij}^{(A)}) \in \mathbb{S}_{+}^{k_j}.$$

The corresponding conditional expectation expression is

$$E(\boldsymbol{\eta}_{i,\mathcal{P}_j} \mid \mathbf{g}_{ij}) = \boldsymbol{\mu}_{\eta,\mathcal{P}_j} + \mathbf{R}_{A,j}(\mathbf{g}_{ij} - \boldsymbol{\mu}_{\eta,\mathcal{P}_j}), \qquad \mathbf{R}_{A,j} = (\boldsymbol{\Sigma}_{gg,j} - \boldsymbol{\Omega}_{A,j}) \boldsymbol{\Sigma}_{gg,j}^{-1}.$$

As in the scaling version, this is an exact conditional expectation under joint normality and otherwise a best linear predictor.

Moreover, because the loadings and latent regression slopes are estimated from centered observations, the empirical aggregates can be calculated without first estimating intercepts:

$$\widehat{g}_{ikj}^{c} = \frac{\widehat{\mathbf{w}}_{kj}^\top (\mathbf{y}_{i,kj} - \overline{\mathbf{y}}_{kj})}{\widehat{\mathbf{w}}_{kj}^\top \widehat{\boldsymbol{\lambda}}_{kj}^{\mathrm{pre}}}.$$

Let $\widehat{\mathbf{g}}_{ij}^{c} \in \mathbb{R}^{k_j}$ collect these centered aggregates, and let $\mathbf{S}_{gg,j} \in \mathbb{S}_{++}^{k_j}$ denote their sample covariance matrix. Under the proxy conditions, the aggregate error covariance estimate is the diagonal matrix

$$\widehat{\boldsymbol{\Omega}}_{A,j} = \operatorname{diag}\left[\left(\frac{\widehat{\mathbf{w}}_{kj}^{\top}\widehat{\boldsymbol{\Theta}}_{kj}^{\text{pre}}\widehat{\mathbf{w}}_{kj}}{\left(\widehat{\mathbf{w}}_{kj}^{\top}\widehat{\boldsymbol{\lambda}}_{kj}^{\text{pre}}\right)^2}\right)_{k\in\mathcal{P}_j}\right].$$

The estimated reliability adjustment and centered fitted predictors are

$$\widehat{\mathbf{R}}_{A,j} = (\mathbf{S}_{gg,j} - c_n\widehat{\boldsymbol{\Omega}}_{A,j})\mathbf{S}_{gg,j}^{-1}, \qquad \widetilde{\boldsymbol{\eta}}_{i,\mathcal{P}_j}^{c,A} = \widehat{\mathbf{R}}_{A,j}\widehat{\mathbf{g}}_{ij}^{c}.$$

Collecting the fitted predictors in $\widetilde{\mathbf{X}}_{A,j} \in \mathbb{R}^{n\times k_j}$ gives

$$\widehat{\mathbf{a}}_{j,\text{JSA}} = (\widetilde{\mathbf{X}}_{A,j}^{\top}\widetilde{\mathbf{X}}_{A,j})^{-1}\widetilde{\mathbf{X}}_{A,j}^{\top}\mathbf{d}_j.$$

The fitted predictor matrices $\widetilde{\mathbf{X}}_j$ and $\widetilde{\mathbf{X}}_{A,j}$ are assumed to have full column rank.

Estimation of Variance, Covariance, and Intercept Parameters

After estimating the factor loadings and latent regression slopes, the remaining variance and covariance parameters can be estimated using several least squares procedures (Maydeu-Olivares et al., in press). For all three estimators, we use reweighted least squares, the default in the implementations considered, holding the estimated loadings and slopes fixed throughout the iterations. Let

$$\widehat{\mathbf{H}} = \widehat{\boldsymbol{\Lambda}}(\mathbf{I}_q - \widehat{\mathbf{B}})^{-1} \in \mathbb{R}^{p\times q},$$

where $\widehat{\boldsymbol{\Lambda}}$ and $\widehat{\mathbf{B}}$ are obtained using the selected estimator, MIIV, JS, or JSA, assuming that $\mathbf{I}_q - \widehat{\mathbf{B}}$ is nonsingular. Conditional on these estimates, the implied covariance matrix is

$$\boldsymbol{\Sigma}(\mathbf{v}) = \widehat{\mathbf{H}}\boldsymbol{\Psi}(\mathbf{v})\widehat{\mathbf{H}}^{\top} + \boldsymbol{\Theta}(\mathbf{v}),$$

where $\mathbf{v} \in \mathbb{R}^{d_v}$ collects the free variance and covariance parameters.

Conditional on the estimated loadings and slopes, the implied covariance moments are linear in the free variance and covariance parameters (Ernst et al., 2023). Incorporating fixed entries and linear equality restrictions gives

$$\operatorname{vech}\{\boldsymbol{\Sigma}(\mathbf{v})\} = \widehat{\mathbf{c}} + \widehat{\mathbf{G}}\mathbf{v},$$

where $t = p(p+1)/2$, $\widehat{\mathbf{c}} \in \mathbb{R}^t$ is the fixed component under the chosen parameterization, and $\widehat{\mathbf{G}} \in \mathbb{R}^{t\times d_v}$ maps the free parameters to the implied covariance moments.

For a weight matrix $\widehat{\mathbf{W}} \in \mathbb{S}_{++}^t$, the weighted least squares estimate is

$$\widehat{\mathbf{v}} = \left(\widehat{\mathbf{G}}^{\top}\widehat{\mathbf{W}}\widehat{\mathbf{G}}\right)^{-1}\widehat{\mathbf{G}}^{\top}\widehat{\mathbf{W}}\left\{\operatorname{vech}(\mathbf{S}) - \widehat{\mathbf{c}}\right\},$$

assuming that $\widehat{\mathbf{G}}$ has full column rank.

Reweighted least squares updates the normal theory weight matrix using the current implied covariance matrix (Browne, 1974; Lee & Jennrich, 1979):

$$\widehat{\mathbf{W}}^{(\tau)} = \frac{1}{2}\mathbf{D}_p^{\top}\left[\{\widehat{\boldsymbol{\Sigma}}^{(\tau)}\}^{-1} \otimes \{\widehat{\boldsymbol{\Sigma}}^{(\tau)}\}^{-1}\right]\mathbf{D}_p, \qquad \widehat{\boldsymbol{\Sigma}}^{(\tau)} = \boldsymbol{\Sigma}(\widehat{\mathbf{v}}^{(\tau)}).$$

Here, $\otimes$ denotes the Kronecker product, and $\mathbf{D}_p \in \mathbb{R}^{p^2\times t}$ is the duplication matrix satisfying $\operatorname{vec}(\mathbf{S}) = \mathbf{D}_p \operatorname{vech}(\mathbf{S})$. At iteration τ, substituting $\widehat{\mathbf{W}}^{(\tau)}$ into the weighted least squares expression gives

$\widehat{\mathbf{v}}^{(\tau+1)}$. The updates are repeated until convergence, assuming that the implied covariance matrices used to construct the weights remain positive definite. The resulting $\widehat{\mathbf{v}}$ determines $\widehat{\boldsymbol{\Theta}}$ and $\widehat{\boldsymbol{\Psi}}$, which are not constrained to be positive semidefinite.

Finally, the intercepts are estimated from the sample means, using the previously estimated factor loadings and latent regression slopes. Under the scaling indicator parameterization, the mean structure is saturated when all nonscaling indicator intercepts and all structural intercepts are free. Let $\overline{\mathbf{m}} \in \mathbb{R}^q$ contain the sample scaling indicator means. Then

$$\widehat{\boldsymbol{\mu}}_\eta = \overline{\mathbf{m}}, \qquad \widehat{\boldsymbol{\nu}} = \overline{\mathbf{y}} - \widehat{\boldsymbol{\Lambda}}\overline{\mathbf{m}} \in \mathbb{R}^p, \qquad \widehat{\boldsymbol{\alpha}} = (\mathbf{I}_q - \widehat{\mathbf{B}})\overline{\mathbf{m}} \in \mathbb{R}^q.$$

These expressions preserve the fixed zero scaling indicator intercepts and reproduce the observed sample means. Additional mean structure restrictions require a corresponding constrained estimation step. Together, these stages estimate the free entries of $\boldsymbol{\Lambda}$, $\mathbf{B}$, $\boldsymbol{\Theta}$, $\boldsymbol{\Psi}$, $\boldsymbol{\nu}$, and $\boldsymbol{\alpha}$. Although the estimated factor loadings and latent regression slopes are held fixed during covariance estimation, their sampling uncertainty is propagated when calculating standard errors.

Analytical Standard Errors

For the MIIV estimates of the factor loadings and latent regression slopes, the conventional homoskedastic covariance estimator is

$$\widehat{\mathbf{V}}_{j,\text{MIIV}} = \widehat{\sigma}^2_{u_j}(\mathbf{X}_j^\top \mathbf{P}_j \mathbf{X}_j)^{-1} \in \mathbb{S}_+^{k_j},$$

where

$$\widehat{\sigma}^2_{u_j} = \frac{\widehat{\mathbf{u}}_j^\top \widehat{\mathbf{u}}_j}{n - k_j - 1}, \qquad \widehat{\mathbf{u}}_j = \mathbf{d}_j - \mathbf{X}_j \widehat{\mathbf{a}}_{j,\text{MIIV}},$$

where the intercept is absorbed by centering and $n > k_j + 1$. Here, $\sigma^2_{u_j}$ is the composite error variance of the L2O equation, rather than an individual measurement error or structural disturbance variance.

This equation specific expression concerns the factor loadings and latent regression slopes. For the variance and covariance parameters $\mathbf{v}$, the delta method is applied to the second stage. Let $\widehat{\mathbf{H}}_2 = (\widehat{\mathbf{G}}^\top \widehat{\mathbf{W}} \widehat{\mathbf{G}})^{-1} \widehat{\mathbf{G}}^\top \widehat{\mathbf{W}}$ be evaluated at the converged weights and treated as fixed, as in Bollen and Maydeu-Olivares (2007) and Jin et al. (2021), let $\widehat{\boldsymbol{\Delta}}_1$ be the Jacobian of $\text{vech}\{\boldsymbol{\Sigma}(\boldsymbol{\theta})\}$ with respect to the factor loadings and latent regression slopes, evaluated at the estimates, and let $\widehat{\mathbf{K}}$ be the Jacobian of their MIIV estimates with respect to $\text{vech}(\mathbf{S})$ (Fisher & Bollen, 2020). Under correct specification, the estimated covariance matrix is

$$\widehat{\mathbf{V}}_{\mathbf{v},\text{MIIV}} = \frac{1}{n} \widehat{\mathbf{H}}_2 (\mathbf{I}_t - \widehat{\boldsymbol{\Delta}}_1 \widehat{\mathbf{K}}) \widehat{\boldsymbol{\Gamma}}_S (\mathbf{I}_t - \widehat{\boldsymbol{\Delta}}_1 \widehat{\mathbf{K}})^\top \widehat{\mathbf{H}}_2^\top \in \mathbb{S}_+^{d_v},$$

where $\widehat{\boldsymbol{\Gamma}}_S \in \mathbb{S}_+^t$ estimates the asymptotic covariance matrix of $\sqrt{n}\,\text{vech}(\mathbf{S})$. The factor $\mathbf{I}_t - \widehat{\boldsymbol{\Delta}}_1 \widehat{\mathbf{K}}$ propagates the sampling uncertainty of the estimated factor loadings and latent regression slopes through the residual moments. The standard errors of the MIIV intercept estimates follow from the same argument applied to the mean structure step.

Burghgraeve et al. (2021) derived a variance expression for the latent regression slopes that accounts for the estimated reliability adjustment and used bootstrap standard errors in their evaluations. Here, analytical standard errors are obtained by differentiating the complete estimation procedure with respect to the sample moments. For $T \in \{\text{JS}, \text{JSA}\}$, write

$$\widehat{\boldsymbol{\theta}}_T = g_{n,T}(\mathbf{s}),$$

where $g_{n,T}$ maps the h sample moments to the d parameter estimates. This mapping includes the preliminary error variance estimates, the estimates of the factor loadings and latent regression slopes, the subsequent variance and covariance estimates, and the intercept estimates.

Suppose

$$\sqrt{n}\{\mathbf{s} - \boldsymbol{\sigma}(\boldsymbol{\theta})\} \xrightarrow{d} N_h(\mathbf{0}, \boldsymbol{\Gamma}), \qquad \boldsymbol{\Gamma} \in \mathbb{S}_+^h,$$

where $\boldsymbol{\Gamma}$ is the joint asymptotic covariance matrix of the sample means and covariances. Under the required moment and differentiability conditions, the delta method covariance estimator is

$$\widehat{\mathbf{V}}_T = \frac{1}{n}\widehat{\mathbf{J}}_T\widehat{\boldsymbol{\Gamma}}\widehat{\mathbf{J}}_T^\top \in \mathbb{S}_+^d, \qquad \widehat{\mathbf{J}}_T = \left.\frac{\partial g_{n,T}(\mathbf{s})}{\partial \mathbf{s}^\top}\right|_{\mathbf{s}=\mathbf{s}_{\text{obs}}} \in \mathbb{R}^{d\times h}.$$

The Jacobian is evaluated analytically by default, including implicit differentiation of the converged reweighted least squares equations. For JSA, it also incorporates derivatives of the estimated aggregation weights and normalization loadings.

The matrix $\boldsymbol{\Gamma}$ can be estimated using normal theory or empirical moments through order four (Browne, 1984). Normal theory estimates may be evaluated at the sample covariance matrix or the fitted model covariance matrix. For parameter $r = 1, \ldots, d$, the analytical standard error is

$$\widehat{\text{SE}}(\widehat{\theta}_{T,r}) = \sqrt{[\widehat{\mathbf{V}}_T]_{rr}} \in [0, \infty).$$

For MIIV factor loadings and latent regression slopes, moment based covariance estimates are also available as $n^{-1}\widehat{\mathbf{K}}\widehat{\boldsymbol{\Gamma}}_S\widehat{\mathbf{K}}^\top$. Together with the second stage expression and the mean structure step above, these provide delta method standard errors calculated separately for each parameter block. A related delta method calculation for instrumental variable regression is presented by Maydeu-Olivares et al. (2019).

Simulation Design and Performance Evaluation

In this section we describe the design of the simulation study and the measures used to evaluate the standard errors.

Population Model and Estimands


We study the model in Figure 1, with three latent variables and four indicators each. The latent variable model is

$$\begin{aligned} \eta_2 &= \alpha_2 + B_{21}\eta_1 + \zeta_2, \\ \eta_3 &= \alpha_3 + B_{31}\eta_1 + B_{32}\eta_2 + \zeta_3. \end{aligned} \tag{1}$$

The design is partly based on previous studies. Burghgraeve et al. (2021) used four indicators per latent variable, a common reliability of .6 or .8 for all indicators. Note also that the model is similar to the empirical illustration in Burghgraeve et al. (2021). Moreover, Jin et al. (2021) used a recursive latent variable model with several equations and standardized slopes between 0.1 and 0.6, which the present model follows partly with $B_{21} = 0.6$, $B_{31} = 0.3$, and $B_{32} = 0.5$. With $\text{Var}(\eta_1) = 1$, $\text{Var}(\zeta_2) = 0.64$, and $\text{Var}(\zeta_3) = 0.48$, the variances of η_2 and η_3 equal 1, so that the slopes are standardized, and the implied correlations are .60 between η_1 and η_2, .60 between η_1 and η_3, and .68 between η_2 and η_3. The scaling indicators have loadings fixed to one and the remaining loadings range from 0.8 to 1.2. The mean structure has $E(\eta_1) = 1$, $\alpha_2 = \alpha_3 = 0.5$, and intercepts of 0.2 for the nonscaling indicators. The main estimands are all free factor loadings, latent regression slopes, residual variances, and latent variances and disturbance variances.

Figure 1

Path diagram of the simulation model with three latent variables and their indicators.



Note. Scaling indicator loadings are fixed to one. The population latent regression slopes are $B_{21} = 0.6$, $B_{31} = 0.3$, and $B_{32} = 0.5$. Intercepts, measurement errors, disturbances, and variances are omitted for clarity.

Simulation Conditions and Estimation Approaches

We employed a fully factorial design with two factors: sample size ($n \in \{100, 200, 300, 400, 500, 600\}$) and indicator reliability ($\rho \in \{.6, .8\}$). This yielded $6 \times 2 = 12$ conditions, each replicated 1,500 times. The sample sizes cover the small to moderate range in which the equation by equation estimators are expected to differ most from ML. The reliability levels match the low and high conditions of Burghgraeve et al. (2021) and determine the size of the reliability adjustment in JS and JSA. As highlighted throughout the article, we investigated the JS, JSA, MIIV, and ML approaches, all with the default options in lavaan (Rosseel, 2012).

Data Generation

Data were generated from the population model with all stochastic components normal and mutually independent. The exogenous latent variable was generated as $\eta_1 \sim \mathcal{N}(1, 1)$ and the structural disturbances as $\zeta_2 \sim \mathcal{N}(0, 0.64)$ and $\zeta_3 \sim \mathcal{N}(0, 0.48)$, from which η_2 and η_3 follow by Equation 1. The measurement error of each indicator was generated as normal with mean zero and variance $\lambda^2(1 - \rho)/\rho$, where λ is the loading of that indicator, so that every indicator has reliability ρ in its condition, and the indicators follow by the measurement equations. The observed variables are therefore multivariate normal with the mean vector and covariance matrix implied by the population values.

Convergence, Outliers, and Inadmissible Solutions

Replications were screened in two steps, and a replication that failed either step for any estimator was removed for all four. A solution was nonconverged when any estimate, standard error, or confidence

Table 1

Performance Measures for Evaluating the Standard Errors

Performance measure	Empirical calculation	MCSE
Relative bias	$(\bar{\theta} - \theta)/\theta$	$S_{\hat{\theta}}/(\sqrt{K}\,\lvert\theta\rvert)$
SD	$S_{\hat{\theta}} = \sqrt{\frac{1}{K-1}\sum_{k=1}^{K}(\hat{\theta}_k - \bar{\theta})^2}$	$\sqrt{\frac{K-1}{K}\sum_{j=1}^{K}(S_{(j)} - S_{\hat{\theta}})^2}$
SE/SD ratio	$\overline{SE}/S_{\hat{\theta}}$	—
SE^2/VAR ratio	$\nu = \overline{SE^2}/S^2_{\hat{\theta}}$	$\sqrt{\frac{K-1}{K}\sum_{j=1}^{K}(\nu_{(j)} - \nu)^2}$
Coverage	$c = \frac{1}{K}\sum_{k=1}^{K} I(L_k \leq \theta \leq U_k)$	$\sqrt{c(1-c)/K}$

Note. $\hat{\theta}_k$ is the estimate of parameter θ in replication k ($k = 1, \ldots, K$) and $\bar{\theta}$ the mean of the K estimates. $\widehat{SE}_k$ is the estimated standard error in replication k, $\overline{SE} = \frac{1}{K}\sum_k \widehat{SE}_k$, $\overline{SE^2} = \frac{1}{K}\sum_k \widehat{SE}_k^2$, and L_k, U_k are the confidence interval limits. The subscript (j) marks a leave-one-out jackknife replicate. The dash marks an MCSE that is not computed.

interval bound of an estimand was missing or non-finite. Among the converged replications, outliers were point estimates more than three interquartile ranges below the first or above the third quartile, per condition, estimator, and estimand. Inadmissible solutions, with a nonpositive variance estimate, were retained and counted.

Performance Measures

Performance was assessed with the measures implemented in the simhelpers package (Joshi & Pustejovsky, 2025), computed for every estimand. These measures, with their empirical and Monte Carlo standard error (MCSE) calculations, are listed in Table 1. The MCSEs of the SD and of the SE^2/VAR ratio are obtained by jackknife and that of coverage in closed form. The SE/SD ratio and the SE^2/VAR ratio compare the same quantities on the standard deviation and on the variance scale, so only the latter, for which an MCSE is available, is shown in the figures. Coverage between .925 and .975 and SE^2/VAR ratios between 0.90 and 1.10 were considered acceptable.

Although the evaluation targets the standard errors, the relative bias of the estimates was also computed, because standard errors of markedly biased estimates are of limited use for inference. It is reported in the text, but not in the figures.

Reproducibility and Computational Details

All analyses were conducted in R (R Core Team, 2026) using the lavaan package (Rosseel, 2012) for estimation, simhelpers (Joshi & Pustejovsky, 2025) for performance evaluation, and ggplot2 (Wickham, 2016) for the figures. The computational environment was pinned with Nix (Dolstra et al., 2004) via the rix package (Rodrigues & Baumann, 2026), and a Dockerfile that builds the same environment inside a container is also provided. The article was written with the apaquarto extension (Schneider, 2026). For an extensive description of all files and how to reproduce the simulation study as well as the manuscript, see the repository.

Results

Convergence, Outliers, and Inadmissible Solutions

All four estimators converged in every replication of every condition. Outlier screening removed 47 of the 18,000 replications (0.26%), concentrated at $n = 100$ with reliability .6, where 1.9% of the replications were removed, flagged mainly through JS (1.7% of the replications in that condition). In every other condition fewer than 0.5% of the replications were removed. Inadmissible solutions occurred in 4 replications, all for JS at $n = 100$ with reliability .6, where a disturbance variance estimate was negative. All of them were also flagged by the outlier screening, so none enters the results below.

Relative Bias

JS, JSA, and ML were essentially unbiased, with absolute relative bias of at most 4.5% for any parameter and condition. MIIV was also nearly unbiased except at reliability .6 and small samples. At $n = 100$ its loadings were slightly underestimated, by 6.2 to 6.9%, whereas its latent and disturbance variances were overestimated by 11.0 to 12.4%. From $n = 300$ onward, all estimators were within 4.6% of the population values.

Standard Error Performance

Each figure presents reliability .6 in the upper half and .8 in the lower half. Figure 2 shows the latent regression slopes, Figure 3 one loading per latent variable, Figure 4 the residual variances of the scaling indicators and one nonscaling indicator per latent variable, and Figure 5 the variance of η_1 and the two disturbance variances. Omitted loadings and residual variances behave similarly to those shown for the same latent variable, with SE^2/VAR ratios that differ by at most 0.15 and coverage by .03 in every condition.

Overall, standard error performance was acceptable. Across all conditions, estimators, and parameters, 96.3% of coverage rates fell between .925 and .975 and 97.9% of SE^2/VAR ratios between 0.90 and 1.10, with no ratio below 0.90. From $n = 300$ onward, the few ratios above 1.10 remained below 1.12. Three departures emerge at $n = 100$ and $n = 200$, mainly at reliability .6.


First, JS standard errors for B_{31} and B_{32} exceed the acceptable range at reliability .6, with SE^2/VAR ratios of 1.5–1.6 at $n = 100$ and 1.13–1.14 at $n = 200$. Coverage nonetheless exceeds .975 only for B_{31} at $n = 100$. Its SD is also the largest, 0.24 versus 0.13–0.17 at $n = 100$. These patterns may reflect instability in the reliability adjustment for the η_3 equation. In some replications, the implied correlation between η_1 and η_2 approaches 1, making the fitted predictors nearly collinear and inflating both estimates and standard errors. Screening removed the most extreme cases, but milder cases remain. JSA, which aggregates indicators within each latent variable, shows no corresponding instability.

Second, MIIV loading intervals fall below the acceptable coverage range at reliability .6, with coverage of .88–.90 at $n = 100$, compared with .93–.96 for the other estimators, despite SE^2/VAR ratios of 0.98–1.14 and the smallest SD. This discrepancy reflects the downward loading bias reported above, consistent with the weak instrument effect described by Jin et al. (2021), and diminishes slowly as sample size increases.

Third, at $n = 100$ MIIV overestimates the standard errors of the latent and disturbance variances, with ratios of 1.11–1.18, but its coverage rates of .96–.97 remain within the acceptable range. ML, in contrast, has ratios close to 1 but coverage rates of .91–.94 for the disturbance variances. JS and JSA fall between them. At reliability .8, the coverage rate for the disturbance variance of η_2 is about .92 for ML, JS, and JSA, just below the acceptable range. Residual variances show less contrast, with coverage rates from .91 to .96 across estimators and a common dip at $n = 100$.

Figure 2


Performance of the Standard Errors of the Latent Regression Slopes



Note. JS = scaling James–Stein, JSA = aggregated James–Stein, IV = MIIV, ML = maximum likelihood. Dotted lines mark the reference values 1 and .95. Error bars represent Monte Carlo standard errors. Values above 1.5 are drawn at 1.5 with an open triangle.

Figure 3


Performance of the Standard Errors of One Factor Loading per Latent Variable



Note. JS = scaling James–Stein, JSA = aggregated James–Stein, IV = MIIV, ML = maximum likelihood. Dotted lines mark the reference values 1 and .95. Error bars represent Monte Carlo standard errors. Values above 1.5 are drawn at 1.5 with an open triangle.

Figure 4


Performance of the Standard Errors of the Residual Variances of the Scaling Indicator and One Nonscaling Indicator per Latent Variable



Note. JS = scaling James–Stein, JSA = aggregated James–Stein, IV = MIIV, ML = maximum likelihood. Dotted lines mark the reference values 1 and .95. Error bars represent Monte Carlo standard errors. Values above 1.5 are drawn at 1.5 with an open triangle.

Figure 5

Performance of the Standard Errors of the Latent and Disturbance Variances



Note. JS = scaling James–Stein, JSA = aggregated James–Stein, IV = MIIV, ML = maximum likelihood. Dotted lines mark the reference values 1 and .95. Error bars represent Monte Carlo standard errors. Values above 1.5 are drawn at 1.5 with an open triangle.

Discussion

This article described MIIV, JS, and JSA estimation as currently implemented in the R package lavaan. It also developed and illustrated analytical standard errors for JS and JSA. A reproducible simulation study evaluated their standard error accuracy against ML under correct model specification.

The findings broadly align with the studies that motivated the simulation design. Bollen (2019), summarizing Bollen et al. (2007), reports few differences between MIIV and ML under correct specification with normally distributed data. Our results agree in terms of bias and standard error accuracy from $n = 300$ onward, although MIIV remained less precise than ML for the latent regression slopes. Jin et al. (2021) found that low scaling indicator reliability can lead to biased MIIV estimates in small samples and that standard errors for variance parameters were the least accurate. Related patterns appeared here at reliability .6 and $n = 100$, with loading bias and undercoverage, alongside inflated standard errors for latent and disturbance variances. Burghgraeve et al. (2021) found JS less efficient than JSA for latent regression estimation and JSA competitive with ML, consistent with the empirical standard deviations observed here. However, their bootstrap standard errors for JS were too small in small samples, whereas our analytical standard errors were too large at reliability .6 and $n \leq 200$. Whether this difference reflects the standard error methods or differences in the simulation conditions requires further investigation.

The simulation was intended as a focused evaluation rather than an exhaustive comparison. Further research should compare MIIV, JS, and JSA across a broader range of conditions, including departures from normality and misspecification of the measurement and latent variable models.

References

Bentler, P. M. (1968). Alpha-maximized factor analysis (alphamax): Its relation to alpha and canonical factor analysis. *Psychometrika*, *33*(3), 335–345. https://doi.org/10.1007/BF02289328

Bollen, K. A. (1989). *Structural equations with latent variables*. Wiley. https://doi.org/10.1002/9781118619179

Bollen, K. A. (1996). An alternative two stage least squares (2SLS) estimator for latent variable equations. *Psychometrika*, *61*(1), 109–121. https://doi.org/10.1007/BF02296961

Bollen, K. A. (2019). Model implied instrumental variables (MIIVs): An alternative orientation to structural equation modeling. *Multivariate Behavioral Research*, *54*(1), 31–46. https://doi.org/10.1080/00273171.2018.1483224

Bollen, K. A., & Bauer, D. J. (2004). Automating the selection of model-implied instrumental variables. *Sociological Methods & Research*, *32*(4), 425–452. https://doi.org/10.1177/0049124103260341

Bollen, K. A., Gates, K. M., & Fisher, Z. F. (2018). Robustness conditions for MIIV-2SLS when the latent variable or measurement model is structurally misspecified. *Structural Equation Modeling: A Multidisciplinary Journal*, *25*(6), 848–859. https://doi.org/10.1080/10705511.2018.1456341

Bollen, K. A., Kirby, J. B., Curran, P. J., Paxton, P. M., & Chen, F. (2007). Latent variable models under misspecification: Two-stage least squares (2SLS) and maximum likelihood (ML) estimators. *Sociological Methods & Research*, *36*(1), 48–86. https://doi.org/10.1177/0049124107301947

Bollen, K. A., & Maydeu-Olivares, A. (2007). A polychoric instrumental variable (PIV) estimator for structural equation models with categorical variables. *Psychometrika*, *72*(3), 309–326. https://doi.org/10.1007/s11336-007-9006-3

Browne, M. W. (1974). Generalized least squares estimators in the analysis of covariance structures. *South African Statistical Journal*, *8*(1), 1–24.

Browne, M. W. (1984). Asymptotically distribution-free methods for the analysis of covariance structures. *British Journal of Mathematical and Statistical Psychology*, *37*(1), 62–83. https://doi.org/10.1111/j.2044-8317.1984.tb00789.x

Burghgraeve, E., De Neve, J., & Rosseel, Y. (2021). Estimating structural equation models using James–Stein type shrinkage estimators. *Psychometrika*, *86*(1), 96–130. https://doi.org/10.1007/s11336-021-09749-2

Carroll, R. J., Ruppert, D., Stefanski, L. A., & Crainiceanu, C. M. (2006). *Measurement error in nonlinear models: A modern perspective* (2nd ed.). Chapman; Hall/CRC. https://doi.org/10.1201/9781420010138

Dhaene, S., & Rosseel, Y. (2023). An evaluation of non-iterative estimators in the structural after measurement (SAM) approach to structural equation modeling (SEM). *Structural Equation Modeling: A Multidisciplinary Journal*, *30*(6), 926–940. https://doi.org/10.1080/10705511.2023.2220135

Dhaene, S., & Rosseel, Y. (2024). An evaluation of non-iterative estimators in confirmatory factor analysis. *Structural Equation Modeling: A Multidisciplinary Journal*, *31*(1), 1–13. https://doi.org/10.1080/10705511.2023.2187285

Dolstra, E., de Jonge, M., & Visser, E. (2004). Nix: A safe and policy-free system for software deployment. *Proceedings of the 18th Large Installation System Administration Conference (LISA '04)*, 79–92.

Efron, B., & Morris, C. (1973). Stein's estimation rule and its competitors: An empirical Bayes approach. *Journal of the American Statistical Association*, *68*(341), 117–130. https://doi.org/10.1080/01621459.1973.10481350

Efron, B., & Morris, C. (1975). Data analysis using Stein's estimator and its generalizations. *Journal of the American Statistical Association*, *70*(350), 311–319. https://doi.org/10.1080/01621459.1975.10479864

Ernst, M. S., Peikert, A., Brandmaier, A. M., & Rosseel, Y. (2023). A note on the connection between

trek rules and separable nonlinear least squares in linear structural equation models. *Psychometrika*, *88*(1), 98–116. https://doi.org/10.1007/s11336-022-09891-5

Fisher, Z. F., & Bollen, K. A. (2020). An instrumental variable estimator for mixed indicators: Analytic derivatives and alternative parameterizations. *Psychometrika*, *85*(3), 660–683. https://doi.org/10.1007/s11336-020-09721-6

Fuller, W. A. (1987). *Measurement error models*. Wiley. https://doi.org/10.1002/9780470316665

James, W., & Stein, C. (1961). Estimation with quadratic loss. *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*, *1*, 361–379.

Jin, S., Yang-Wallentin, F., & Bollen, K. A. (2021). A unified model-implied instrumental variable approach for structural equation modeling with mixed variables. *Psychometrika*, *86*(2), 564–594. https://doi.org/10.1007/s11336-021-09771-4

Jöreskog, K. G., & Sörbom, D. (1996). *LISREL 8: User's reference guide*. Scientific Software International.

Joshi, M., & Pustejovsky, J. E. (2025). *simhelpers: Helper functions for simulation studies*. https://CRAN.R-project.org/package=simhelpers

Kaplan, D. (1988). The impact of specification error on the estimation, testing, and improvement of structural equation models. *Multivariate Behavioral Research*, *23*(1), 69–86. https://doi.org/10.1207/s15327906mbr2301_4

Lee, S.-Y., & Jennrich, R. I. (1979). A study of algorithms for covariance structure analysis with specific comparisons using factor analysis. *Psychometrika*, *44*(1), 99–113. https://doi.org/10.1007/BF02293789

Maydeu-Olivares, A., Rosseel, Y., Fisher, Z. F., & Bollen, K. A. (in press). Getting the second stage right: Evaluating model implied instrumental variable (MIIV) estimators for variances and covariances. *Structural Equation Modeling: A Multidisciplinary Journal*.

Maydeu-Olivares, A., Shi, D., & Rosseel, Y. (2019). Instrumental variables two-stage least squares (2SLS) vs. Maximum likelihood structural equation modeling of causal effects in linear regression models. *Structural Equation Modeling: A Multidisciplinary Journal*, *26*(6), 876–892. https://doi.org/10.1080/10705511.2019.1607740

R Core Team. (2026). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://doi.org/10.32614/R.manuals

Rodrigues, B., & Baumann, P. (2026). *rix: Reproducible data science environments with Nix*. https://docs.ropensci.org/rix/

Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, *48*(2), 1–36. https://doi.org/10.18637/jss.v048.i02

Rosseel, Y., & Vidal, M. (2025). A tutorial for understanding SEM using R: Where do all the numbers come from? *British Journal of Mathematical and Statistical Psychology*. https://doi.org/10.1111/bmsp.70003

Schneider, W. J. (2026). *apaquarto: A Quarto extension for creating APA 7 style documents*. https://github.com/wjschne/apaquarto

Whittemore, A. S. (1989). Errors-in-variables regression using Stein estimates. *The American Statistician*, *43*(4), 226–228. https://doi.org/10.1080/00031305.1989.10475663

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org