

- of lookout on Warm Springs Mountain, elevation 3480', 55 meters above basement complex—San Francisquito Formation contact in Kirby's measured section number 2 (Kirby, 1991, p. 119), Los Angeles County, California. Collected by: M. X. Kirby, 1990. San Francisquito Formation; early Danian.
- 15790 LACMIP: West bank Deer Creek, 160 m N, 240 m E of SW corner sec. 32, T26N, R2E, Panther Spring quadrangle, Tehama County, California. Collected by: John Russell, 1985. Chico Formation, Tenmile Member; early Campanian.
- 15792 LACMIP: East bank Deer Creek 200 m S, 375 m E of NW corner sec. 5, T25N, R2E, Panther Spring quadrangle, Tehama County, California. Collected by: John Russell, 1986. Chico Formation, Tenmile Member; early to early middle Campanian, *Submortoniceras chicoense* Zone.
- 27135 LACMIP [=7135 UCLA]: Fossils collected within 3 m sec-
- 27838 tion in highly fossiliferous zone, W side Horse Canyon, near 1800' contour, 1.125 km N, 1.2 km E of SW corner of topo, T9N, R30W, Bates Canyon quadrangle, Sierra Madre Mountains., Santa Barbara County, California. Collected by: G. Smith, January 1984. Unnamed Formation; late Campanian. CASG: Chico Creek, 3.6 miles from "10 Mile House" on Humboldt Road, U.S.G.S. Chico quadrangle (1895, reprint 1932), Butte County, California. Collected by: H. A. Taff, G D., Hanna, and C. M. Cross, May, 1934. Chico Formation, Tenmile Member; early Campanian.
- 28323 CASG: Gully on N side of Del Puerto Creek, 1000' below conglomerate, SW 1/4, 1/4 of sec. 35, T5S, R6E, Copper Mountain 1:24000 quadrangle, Stanislaus County, California. Collected by: J. A. Taff, 1935. Panoche Formation; early Santonian(?)